



Name : Mr. SINGH JOGINDER S/o UHID : 120385 S No : PID : 33294  
Age/Gender : 53 Year/Male A.S : NP Sample Date : 31-Jul-2024 11:14 AM  
Ref. By Dr. : MEDIWHEEL Report Date : 31-Jul-2024 09:18 PM  
Address : HISAR Status : Pending Sample Type : Inside \*33294\*

Test Name Value Unit Reference Range

HEMATOLOGY

CBC (Complete Blood Count)

Haemoglobin (Hb)	14.1	g/dl	12.0 - 17.4 g/dl
Total RBC Count	4.86	m/cumm	4.70 - 6.10
Haematocrit	44.9	%	35.0 - 50.0 %
Mean Cell Volume	92.4	fL	80.0 - 100 fL
Mean Cell Haemoglobin	31.8	pg	27.0 - 34.0 pg
Mean Cell Haemoglobin Conc	34.4	%	32.0 - 36.0
Red Cell Distribution Width (RDW)-CV	13.6	%	11.0 - 16.0 %
Red Cell Distribution Width (RDW)-SD	50.8	fL	35.0 - 56.0 fL
Total Leucocyte Count	5880	cells/cum m	4000 - 11000
Differential Leucocyte Count	.		
Neutrophils	45	%	32 - 72 %
Lymphocytes	50	%	20 - 50 %
Monocytes	003	%	2 - 11 %
Eosinophils	02	%	1 - 3 %
Basophils	0	%	0 - 2 %
Platelet Count	2,25,000	cells/cunm m	150,000 - 450,000
Platelet Distribution Width	15.2	fL	15.0 - 18.0 fL
Mean Platelet Volume	10.3	fL	7.0 - 13.0 fL

Sample Type : Whole Blood

- Spurious elevation of platelet count may be seen in patients with extensive burns, extreme microcytosis, microangiopathic hemolytic anemia, red cell fragmentation, micro-organisms like bacteria, fungi or yeast, hyperlipidemia, fragments of white blood cell (WBC) cytoplasm in patients with acute leukemia, hairy cell leukemia, lymphomas and in presence of cryoglobulins.
- Spuriously low platelet counts may be seen in cases of platelet clumping (EDTA induced), platelet cold agglutinins, multiple myeloma, platelet satellitism and in giant platelet syndromes.
- Delay in processing due to sample transport may cause a mild time dependent fall in platelet count. It is advisable to repeat the test using a citrate / heparin collection tube to avoid this pitfall.
- Automated platelet counting is subject to 10-15% variation in the result on the same as well as different analysers due to various preanalytic variables like the sampling site, skill in sample collection, anticoagulant used, sample mixing and sample transport etc.

ABO Blood Grouping

Blood Group

O<sup>+</sup>NEGATIVE

Haemaagglutination reaction

A Rh Positive, B Rh Positive, AB Rh Positive, O Rh Positive, A Rh Negative, B Rh Negative, AB Rh Negative, O Rh Negative

Sample Type : Whole Blood

HBA1C

HBA1C 4.9 % 4.27 - 6.00 %



Name : Mr. SINGH JOGINDER S/o      UHID : 120385      PID : 33294  
Age/Gender : 53 Year/Male      Sample Date : 31-Jul-2024      11:14 AM  
Ref. By Dr. : MEDIWHEEL      Report Date : 31-Jul-2024  
Address : HISAR      Sample Type : Inside      \*33294\*

Test Name	Value	Unit	Reference Range
-----------	-------	------	-----------------

## HBA1C

turbidimetric immunoassay

Average Blood Glucose

93.93

mg/dl

90.00 - 120.00 mg/dl

turbidimetric immunoassay

Sample Type : Whole Blood

Remarks :

GLYCOSYLATED HEMOGLOBIN (HbA1c)

Reference Range : Please correlate with clinical conditions.

Bellow 6.0 % Normal value

6.0 %-7.0 % Good control

7.0 %-8.0 % Fair control

8.0 %-10 % Unsatisfactory control

Above10 % Poor control

Technology : Immunoassay and chemistry technology to measure A1C and total HB (A1C now Bayer)

AVERAGE BLOOD GLUCOSE (ABG) CALCULATED

Reference Range: Please correlate with clinical conditions.

90-120 mg/dl Excellent control

121-150 mg/d Good control

151-180 mg/dl Average control

181-210 mg/dl Action suggested

> 211 mg/dl Panic values

NOTE: Average blood glucose value is calculated from HbA1C value and it indicates average blood sugar level over past three months.

Technology: Derived from Hb A1C Values

Sample Type: Sodium heparin:

## ESR

ESR	20	mmHr	0 - 15 mmHr
-----	----	------	-------------

Sample Type : Whole Blood



Name : Mr. SINGH JOGINDER S/o UHID : 120385 S No : PID : 33294  
Age/Gender : 53 Year/Male A.S : NP Sample Date : 31-Jul-2024 11:14 AM  
Ref. By Dr. : MEDIWHEEL Report Date : 31-Jul-2024 09:23 PM  
Address : HISAR Status : Pending Sample Type : Inside \*33294\*

Test Name	Value	Unit	Reference Range
-----------	-------	------	-----------------

#### CLINICAL COMMENTS:

Erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that indirectly measures the degree of inflammation present in the body. Inflammation is part of the body's immune response. It can be acute, developing rapidly after trauma, injury or infection, for example, or can occur over an extended time (chronic) with conditions such as autoimmune diseases or cancer.

Moderately elevated ESR occurs with inflammation but also with anemia, infection, pregnancy, and with aging. A very high ESR usually has an obvious cause, such as a severe infection, marked by an increase in globulins, systemic vasculitis, polymyalgia rheumatica or temporal arteritis. People with multiple myeloma or Waldenstrom's macroglobulinemia (tumors that make large amounts of immunoglobulins) typically have very high ESRs even if they don't have inflammation.

#### Factors increasing ESR:

- Advanced age
- Anemia
- Pregnancy
- High fibrinogen
- Macrocytosis
- Kidney problems
- Thyroid disease
- Some cancers, such as multiple myeloma
- Infection

#### Factors decreasing ESR

- Microcytosis
- Low fibrinogen
- Polycythemia
- Marked leukocytosis

### CLINICAL-CHEMISTRY

#### URIC ACID

Uric acid	6.41	mg/dL	3.5 - 7.2
-----------	------	-------	-----------

Uricase - POD

Sample Type : SERUM

URIC ACID: Increases in case of renal failure, disseminated neoplasms, pregnancy toxemia, psoriasis, liver disease, sarcoidosis etc. Decrease is reported in Wilson's disease, Fanconi's syndrome, xanthinuria.

#### Total Protein

Total Protein	6.64	gm/dl	6.0 - 8.3
BIURET			
Albumin	3.98	g/dl	2.9 - 4.5
BCG			
Globulin	2.66	gm/dl	2.0 - 3.5
Albumin-Globulin Ratio	1.32		1.2 - 2.5



Name : Mr. SINGH JOGINDER S/o      UHID : 120385      S No :      PID : 33294  
Age/Gender : 53 Year/Male      A.S : NP      Sample Date : 31-Jul-2024      11:14 AM  
Ref. By Dr. : MEDIWHEEL      Report Date : 31-Jul-2024      10:20 PM  
Address : HISAR      Status : Pending      Sample Type : Inside      \*33294\*

Test Name	Value	Unit	Reference Range
-----------	-------	------	-----------------

Sample Type : SERUM

### UREA. SERUM

UREA	38.4	mg/dL	14 - 51
------	------	-------	---------

KINETIC METHOD WITH UREASE AND GLDH  
Sample Type : SERUM

UREA: High urea levels suggest poor kidney function, congestive heart failure, shock, stress, recent heart attack or severe burns; bleeding from the gastrointestinal tract; conditions that cause obstruction of urine flow; or dehydration.

Low urea levels can be seen in severe liver disease or malnutrition but are not used to diagnose or monitor these conditions. Low urea levels are also seen in normal pregnancy.

### CREATININE SERUM

CREATININE SERUM	0.9	mg/dL	0.5 - 1.4 mg/dL
------------------	-----	-------	-----------------

Jaffe Kinetic  
Sample Type : SERUM

CREATININE: Increases in any renal functional impairment (intrinsic renal lesions, decreased perfusion of the kidney, or obstruction of the lower urinary tract), acromegaly and hyperthyroidism. Decreases in pregnancy, muscle wasting.

### KIDNEY FUNCTION TEST (KFT Special )

UREA	38.4	mg/dL	14 - 45 mg/dL
------	------	-------	---------------

KINETIC METHOD WITH UREASE AND GLDH

CREATININE SERUM	0.9	mg/dL	0.5 - 1.4
------------------	-----	-------	-----------

Jaffe Kinetic

Uric acid	6.41	mg/dL	3.5 - 7.2
-----------	------	-------	-----------

Uricase - POD

BUN SERUM	17.94	mg/dL	07 - 24
-----------	-------	-------	---------

KINETIC METHOD WITH UREASE & GLDH

SODIUM-SERUM	141.2	mmol/L	135 - 150
--------------	-------	--------	-----------

ISE(DIRECT)

POTASSIUM SERUM	3.6	mmol/L	3.5 - 5.0
-----------------	-----	--------	-----------

ISE(DIRECT)

Chloride	101.3	mmol/L	96 - 106
----------	-------	--------	----------

Ion Selective Electrode (indirect)

Urea / Creatinine Ratio	42.67		40:1 - 100:1
-------------------------	-------	--	--------------

BUN / Creatinine Ratio	19.93		10:1 - 20:1
------------------------	-------	--	-------------

Sample Type : SERUM



Name : Mr. SINGH JOGINDER S/o      UHID : 120385      S No :      PID : 33294  
Age/Gender : 53 Year/Male      A.S : NP      Sample Date : 31-Jul-2024      11:14 AM  
Ref. By Dr. : MEDIWHEEL      Report Date : 31-Jul-2024      09:26 PM  
Address : HISAR      Status : Pending      Sample Type : Inside      \*33294\*

Test Name	Value	Unit	Reference Range
-----------	-------	------	-----------------

### CLINICAL COMMENTS :

UREA: High urea levels suggest poor kidney function, congestive heart failure, shock, stress, recent heart attack or severe burns; bleeding from the gastrointestinal tract; conditions that cause obstruction of urine flow; or dehydration.

Low urea levels can be seen in severe liver disease or malnutrition but are not used to diagnose or monitor these conditions. Low urea levels are also seen in normal pregnancy.

CREATININE: Increases in any renal functional impairment (intrinsic renal lesions, decreased perfusion of the kidney, or obstruction of the lower urinary tract), acromegaly and hyperthyroidism. Decreases in pregnancy, muscle wasting.

URIC ACID: Increases in case of renal failure, disseminated neoplasms, pregnancy toxemia, psoriasis, liver disease, sarcoidosis etc. Decrease is reported in Wilson's disease, Fanconi's syndrome, xanthinuria.

SODIUM: Increases due to water loss (severe diarrhea profuse sweating, polyuria or vomiting), hypergluco- or mineralo-corticoidism, and inadequate water intake. Decreases due to intake of free water or

### LIVER FUNCTION TEST (LFT) (S)

Total Bilirubin-Serum	0.90	mg/dl	0.20 - 1.00 mg/dl
Bilirubin Direct Serum	0.40	mg/dl	0.10 - 0.50 mg/dl
Bilirubin Indirect-Serum	0.50	mg/dl	0.20 - 0.70 mg/dl
<b>SGOT</b> IFCC with Pyridoxal Phosphate	<b>62.9</b>	IU/L	10 - 40 IU/L
<b>SGPT</b> IFCC with Pyridoxal Phosphate	<b>85.4</b>	IU/L	07 - 56 IU/L
Alkaline Phosphatase IFCC PNPP Buffer	110.5	U/L	44 - 147 U/L
Total Protein	6.64	gm/dl	6.0 - 8.3
BIURET Albumin	3.98	g/dl	3.5 - 5.5 g/dl
BCG Globulin	2.66	gm/dl	2.0 - 3.5 gm/dl
AG RATIO	1.32		1.2 - 2.5

Sample Type : SERUM





Name : Mr. SINGH JOGINDER S/o UHID : 120385 S No : PID : 33294  
Age/Gender : 53 Year/Male A.S : NP Sample Date : 31-Jul-2024 11:14 AM  
Ref. By Dr. : MEDIWHEEL Report Date : 31-Jul-2024 10:20 PM  
Address : HISAR Status : Pending Sample Type : Inside \*33294\*

Test Name	Value	Unit	Reference Range
-----------	-------	------	-----------------

#### CLINICAL COMMENT:

Liver function tests can be suggested in case of hepatitis, liver cirrhosis and monitor possible side effects of medications. A variety of diseases and infections can cause acute or chronic damage to the liver, causing inflammation

(hepatitis), scarring (cirrhosis), bile duct obstructions, liver tumors, and liver dysfunction. Alcohol, drugs, some herbal supplements, and toxins can also injure the liver. A significant amount of liver damage may occur before symptoms such as jaundice, dark urine, light-colored stools, itching (pruritus), nausea, fatigue, diarrhea, and unexplained weight loss or gain appear. Early detection of liver injury is essential in order to minimize damage and preserve liver function.

Alanine aminotransferase (ALT) A very high level of ALT is frequently seen with acute hepatitis. Moderate increases may be seen with chronic hepatitis. People with blocked bile ducts, cirrhosis, and liver cancer may have ALT concentrations that are only moderately elevated or close to normal. Aspartate aminotransferase (AST) A very high level of AST is frequently seen with acute hepatitis. AST may be normal to moderately increased with chronic hepatitis. In people with blocked bile ducts, cirrhosis, and liver cancer, AST concentrations may be moderately increased or close to normal. When liver damage is due to alcohol, AST often increases much more than ALT (this is a

pattern seen with few other liver diseases). AST is also increased after heart attacks and with muscle injury.

AST is a less sensitive and less specific marker of liver injury than ALT. AST is more elevated than ALT in alcohol-induced liver injury. AST could be elevated more than ALT like: (i)

#### Lipid Profile

Cholesterol	125.47	mg/dl	<200.0 mg/dl
CHOD - PAP			
Triglycerides	180.1	mg/dl	< 150 mg/dl
GPO - PAP			
HDL Cholesterol	42.1	mg/dl	Adult males >45 mg/dl
Homogeneous Enzymatic Colorimetric test			
LDL Cholesterol	47.35	mg/dl	<100 mg/dl
VLDL Cholesterol	36.02	mg/dl	<30.0 mg/dl
CHO/HDL Ratio	2.98	mg/dl	Low risk 3.3-4.4
Non HDL Cholesterol	83.37	mg/dl	<130 mg/dl

Calculated

Sample Type : SERUM

#### Interpretation

##### Note

- Measurements in the same patient can show physiological & analytical variations. 3 serial samples 1 wk apart are recommended for Total Cholesterol, Triglycerides, HDL & LDL Cholesterol.
- NLA-2014 identifies Non HDL Cholesterol (an indicator of all atherogenic lipoproteins such as LDL, VLDL, IDL, Lp(a), Chylomicron remnants) along with LDL-cholesterol as co-primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL & Non HDL.
- Apolipoprotein B is an optional, secondary lipid target for treatment once LDL & Non HDL goals have been achieved.
- Additional testing for Apolipoprotein B, hsCRP, Lp(a) & LP-PLA2 should be considered among patients with moderate risk for ASCVD for risk refinement.

## CLINICAL PATHOLOGY

### PHYSICAL EXAMINATION



# Lotus Diagnostic & Imaging Centre

A Unit of Lotus Diagnostic & Imaging Solution Pvt. Ltd.

HB से लेकर MRI तक एक ही छत के नीचे

Name : Mr. SINGH JOGINDER S/o UHID : 120385 S No : PID : 33294  
Age/Gender : 53 Year/Male A.S : NP Sample Date : 31-Jul-2024 11:14 AM  
Ref. By Dr. : MEDIWHEEL Report Date : 31-Jul-2024 09:27 PM  
Address : HISAR Status : Pending Sample Type : Inside \*33294\*

Test Name	Value	Unit	Reference Range
Colour Pale-yellow, Yellowish, Colorless, YELLOW	PALE YELLOW		
Quantity	30	ml	
pH	6.2		
Mucus Absent, Present	ABSENT		
Appearance Slightly turbid, Turbid, Clear	CLEAR		
<b>Chemical Examination (Strip)</b>	.		
Specific Gravity	1.025		
Albumin Absent, Present(+), Present(2+), Present(3+)	NEGATIVE		
Sugar Absent, Present(+), Present(2+), Present(3+)	NEGATIVE		
Bilirubin Absent, Present	NEGATIVE		
<b>Microscopic Examination (Microscopy)</b>	.		
Pus Cells	1-3	/HPF	
Epithelial Cells	0-1	/HPF	
RBC	NIL	/HPF	
Casts	ABSENT		
Crystals	ABSENT		
Bacteria	ABSENT		
Others			
Sample Type : Urine			

## Laboratory

Blood Sugar (PP) Blood Sugar PP Sample Type : Others	110.9	mg/dl	70.00 - 140.00 mg/dl
URINE SUGAR FASTING Sample Type : Urine	78.4		
URINE SUGAR PP Sample Type : Urine	78.4		70 - 110

## ENDOCRINE

### Thyroid Hormones (T3 .T4 & TSH)

T3	0.72	ng/ml	0.60 - 1.81 ng/ml
T4	8.39	ng/dl	5.01 - 12.45 ng/dl
TSH Ultrasensitive	4.32	uIU/ml	0.3 - 4.5 uIU/ml

Dr. (Maj.) Guruprasad  
MBBS, DMRD, DNB  
Consultant Radiologist

Dr. Rambaksh Sharma  
MBBS, MD  
Consultant Radiologist

Dr. RAJESH REDDU  
MBBS, DMRD  
Consultant Radiologist

Dr. Amit Verma  
MBBS, MD  
Consultant Physician

Dr. Manish Varshney  
MBBS, MD  
Consultant Pathologist



# Lotus Diagnostic & Imaging Centre

A Unit of Lotus Diagnostic & Imaging Solution Pvt. Ltd.

HB से लेकर MRI तक एक ही छत के नीचे

Name : Mr. SINGH JOGINDER S/o UHID : 120385 S No : PID : 33294  
Age/Gender : 53 Year/Male A.S : NP Sample Date : 31-Jul-2024 11:14 AM  
Ref. By Dr. : MEDIWHEEL Report Date : 31-Jul-2024 08:17 PM  
Address : HISAR Sample Type : Inside \*33294\*

Test Name	Value	Unit	Reference Range
-----------	-------	------	-----------------

Sample Type : SERUM

Remarks :

Note1.TSH levels are subject to circadian variation, reaching peak levels between 2-4.a.m and at a minium between 6-10 pm. The variation is of the 50 %, hence time of the day has influence on the measured serum TSH concentrations.

2. Recommended test for T3 and T4 unbound or free level as it is metabollically active.

3. Physiological rise in Total T3 and T4 level is seen in pregnancy and in patients on steroid therapy.

Clinical Use-

- \* Primary Hypothyroidism
- \* Hperthyroidism
- \* Hypothalamic- Pituitary hypothyroidism
- \* Inappropriate-TSH secretion
- \* Nonthyroidal illness
- \* Autoimmune thyroid disease
- \* Pregnancy associated thyroid disorders
- \* Thyroid dysfunction in infancy and early childhood

## IMMUNOLOGY

Total PSA	1.09	ng/ml	0.00 - 4.0 ng/ml
-----------	------	-------	------------------

Sample Type : SERUM

Summary & Interpretation:

Elevated concentrations of PSA in serum are generally indicative of a patho-logic-condition of the prostate (prostatitis, begin hyperplasia or carcinoma). PSA determinations are employed are the monitoring of progress and efficiency of therapy in patients with prostate carcinoma or receiving hormonal therapy . An inflammation or trauma of the prostate(e.g. In case of urinary retention or following rectal examination, cystoscopy, coloscopy, transurethral biopsy, lasertreatment or ergometry)can lead to PSA elevations of varying duration and magnitu

--End of Report--

Dr. (Maj.)Guruprasad  
MBBS, DMRD, DNB  
Consultant Radiologist

Dr. Rambaksh Sharma  
MBBS, MD  
Consultant Radiologist

Dr. RAJESH REDDU  
MBBS, DMRD  
Consultant Radiologist

Dr. Amit Verma  
MBBS, MD  
Consultant Physician

Dr. Manish Varshney  
MBBS, MD  
Consultant Pathologist





PATIENT NAME: SINGH JOGINDER  
REF. BY: TPA

AGE/SEX: 53 YRS/M  
DATE: JULY 31, 2024

**USG WHOLE ABDOMEN**

**Exaggerated pattern of bowel gases, poor acoustic window.**

**Liver:** normal in size. Parenchymal echotexture is normal and no focal area of altered echogenicity is seen. IHBR not dilated. CBD is normal in diameter.

**GB:** is contracted however appears normal. Wall thickness is normal.

**Pancreas:** head and body shows normal size and parenchymal attenuation.

**Spleen:** normal in size and normal echotexture.

**Right Kidney:** is not seen in right renal fossa and lying in right side of pelvis.

**Right kidney measures 9.2 x 3.4 cm.—Ectopic kidney.** No evidence of any calculus detected. Pelvi calyceal system is normal. CMD is maintained.

**Left Kidney:** measures 12.3 x 5.4 cm, is normal in position, size and morphology. No evidence of any calculus detected. Pelvi calyceal system is normal. CMD is maintained.

**Urinary Bladder:** appears normal.

**Prostate:** normal in size and echotexture. No significant median lobe bulge seen.

No obvious abnormal bowel dilatation or wall thickening is seen in present scan.

No free fluid seen.

**IMPRESSION:** - Right kidney is not seen in right renal fossa and lying in right side of pelvis.—Ectopic kidney.

Clinical correlation and further evaluation is suggested.

**Dr. Ram Baksh Sharma**  
Radiologist

**Dr. Rambaksh Sharma**  
Consultant Radiologist

**Dr. Anshul Jain**  
Consultant Radiologist

**Dr. Rajesh Reddu**  
MBBS, DMRD  
Consultant Radiologist

**Dr. Amit Verma**  
Echocardiography Specialist

**Dr. Sonam Aneja**  
Consultant Pathologist






# GEETANJALI HOSPITAL HISAR HARYANA I

Age / Gender: 53/Male  
Patient ID: 0000327150

Date and Time: 31st Jul 24 12:41 PM

*Ms. Singh Jagindes*



AR: 74bpm    VR: 74bpm    QRSD: 108ms    QT: 373ms    QTcB: 415ms    PR: 148ms    P-R-T: 47° 53° 35°

ECG Within Normal Limits: Sinus Rhythm. Please correlate clinically.

REPORTED BY



Disclaimer: Analysis in this report is based on ECG alone and should only be used as an adjunct to clinical history, symptoms and results of other invasive and non-invasive tests and must be interpreted by a qualified physician.

KMC 95602