



Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 2024-10-26 15:48:50
UHID/MR NO	: ALDP.0000152994	Received	: 2024-10-26 15:48:50
Visit ID	: ALDP0284052425	Reported	: 27/Oct/2024 10:08:45
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

# DEPARTMENT OF CARDIOLOGY-ECG MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

## ECG/EKG

1. Machnism, Rhythm	Sinus, Regular	
2. Atrial Rate	81	/mt
3. Ventricular Rate	81	/mt
4. P - Wave	Normal	
5. P R Interval	Normal	
6. Q R S Axis : R/S Ratio : Configuration :	Normal Normal Normal	
7. Q T c Interval	Normal	
8. S - T Segment	Normal	
9. T – Wave	Normal	

9. 1 – wa <u>FINAL IMPRESSION</u>

Abnormal: Sinus Rhythm, Abnormally Long QT Interval observed.Hyperacute T waves in leads V2. LVH by votage criteria noted. Please correlate clinically.





View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:11
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 15:30:25
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing), Blood				
Blood Group	В			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Rh ( Anti-D)	POSTIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Complete Blood Count (CBC), Whole Blood				
Haemoglobin	14.10	g/ dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRICMETHOD (CYANIDE-FREE REAGENT)
TLC (WBC) DLC	7,600.00	/Qumm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils )	60.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	33.00	%	20-40	FLOW CYTOMETRY
Monocytes	4.00	%	2-10	FLOW CYTOMETRY
Eosinophils	3.00	%	1-6	FLOW CYTOMETRY
Basophils <b>ESR</b>	0.00	%	<1-2	FLOW CYTOMETRY
Observed	4.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5 80-91 Yr 15.8	



Page 2 of 13







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:11
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 15:30:25
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

## DEPARTMENT OF HAEM ATOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	-	Mm for 1st hr.	,	
PCV (HCT)	43.00	%	40-54	
Platelet count				
Platelet Count	1.45	LACS cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.50	fL	9-17	<b>ELECTRONIC IMPEDANCE</b>
P-LOR (Platelet Large Cell Patio)	-	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.20	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	14.50	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBCCount				
RBCCount	5.14	Mill./cumm	4.2-5.5	<b>ELECTRONIC IMPEDANCE</b>
Blood Indices (MCV, MCH, MCHC)				
MCV	85.30	fl	80-100	CALCULATED PARAMETER
МСН	27.50	pg	27-32	CALCULATED PARAMETER
МОНС	32.20	%	30-38	CALCULATED PARAMETER
RDW-CV	13.30	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	45.40	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	4,560.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	228.00	/cu mm	40-440	

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965.0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D /M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 16:25:21
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
GLUCOSE FASTING, Pasma Gucose Fasting	86.70	Ŭ 1	: 100 Normal 00-125 Pre-diabetes : 126 Diabetes	GOD POD

#### Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions. b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential. c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body . Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP	116.70	mg/ dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

#### **Interpretation:**

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions. b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential. c) I.G.T = Impaired Glucose Tolerance.

### GLYCOSYLATED HAEM OGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	6.00	%NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	41.80	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	125	mg/dl	

#### Interpretation:

#### NOTE:-

• eAG is directly related to A1c.





View Reports on







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 16:25:21
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Test Name	nesuit	Unit	DIU. HEI. IIILEI VAI	method

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	<b>Degree of Glucose Control Unit</b>
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

\*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. \*\*Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

#### **Clinical Implications:**

\*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

\*With optimal control, the HbA 1c moves toward normal levels.

\*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated \*Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

\*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

\*Pregnancy d. chronic renal failure. Interfering Factors:

\*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (	(Blood	Urea	Nitroger	1)
Sampl	e:Serum		-	

7.75

mg/dL 7.0-23.0

CALCULATED





Page 5 of 13





Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

			<b>–</b> • •			
Patient Name	: Mr.ABHAY SINGH		Registered	d On	: 26/Oct/2024 1	
Age/Gender UHID/MR NO	: 34 Y 10 M 27 D /M : ALDP.0000152994		Collected Received		: 26/Oct/2024 1 : 26/Oct/2024 1	
Visit ID	: ALDP:0000152994 : ALDP0284052425		Reported		: 26/Oct/2024 1	
	: Dr. MEDIWHEEL-ARCO	FEMI HEALTH	-			
Ref Doctor	CARE LTD -		Status		: Final Report	
		DEPARTMEN	T OF BIOCH	HEMIST	RY	
	MEDIWH	IEEL BANK OF	-			
Test Name		Result	ι	Jnit	Bio. Ref. Interva	l Method
Interpretation:						
Note: Elevated B	UN levels can be seen in th	e following:				
High-protein diet, D	Dehydration, Aging, Certain m	edications, Burns,	, Gastrointesti	mal (GI) t	leeding.	
Low BUN levels c	an be seen in the following	:				
Low-protein diet, o	verhydration, Liver disease.					
reatinine ample:Serum Interpretation: The significance of	single creatinine value must be	<b>1.32</b> e interpreted in lig	mg/dl ht of the patie:	0.7-1.3		MODIFIED JAFFES
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m	single creatinine value must be gher creatinine concentration. 7 concentration. Serum creatinin ildly and may result in anomal	e interpreted in lig The trend of serum ne concentrations	ht of the patien n creatinine co may increase	nts muscle oncentration when an a	e mass. A patient wi ons over time is mo ACE inhibitor (AC	ith a greater muscle re important than E) is taken. The assay
ample:Serum Interpretation: The significance of mass will have a hig absolute creatinine of	gher creatinine concentration.	e interpreted in lig The trend of serum ne concentrations	ht of the patien n creatinine co may increase	nts muscle oncentration when an a	e mass. A patient wi ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay
Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Interpretation: Note:- Elevated uric acid	gher creatinine concentration. The concentration of	e interpreted in lig The trend of serun he concentrations lous values if seru 5.94 5.94	ht of the patie n creatinine co may increase m samples ha mg/dl	nts muscle oncentratie when an A ve heterop 3.4-7.6	e mass. A patient wi ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or
Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Fric Acid Ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p	gher creatinine concentration. The concentration of	e interpreted in lig The trend of serun he concentrations lous values if seru 5.94 5.94	ht of the patie n creatinine co may increase m samples ha mg/dl	nts muscle oncentratie when an A ve heterop 3.4-7.6	e mass. A patient wi ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or
Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM	gher creatinine concentration. The concentration of	e interpreted in lig The trend of serum ne concentrations lous values if seru 5.94 <b>5.94</b> <b>bllowing:</b> kidney disease, H	ht of the patien n creatinine co may increase m samples ha mg/ dl	nts muscle oncentration when an a ve heterop 3.4-7.0 Dbesity.	e mass. A patient wi ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or URICASE
ample:Serum Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A	gher creatinine concentration. The concentration. Serum creatining ildly and may result in anomal <b>Hevels can be seen in the for</b> protein diet, alcohol), Chronic <b>HAGT</b> ), <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serum ne concentrations lous values if seru 5.94 5.94 bllowing: kidney disease, H	ht of the patien n creatinine co may increase m samples har mg/ dl	nts muscle oncentration when an ve heterop 3.4-7.0 Obesity.	e mass. A patient wa ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or URICASE
Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Fric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am	gher creatinine concentration. The concentration of	e interpreted in lig The trend of serum ne concentrations lous values if seru 5.94 <b>5.94</b> <b>5.94</b> <b>bllowing:</b> kidney disease, H <b>43.40</b> <b>59.70</b>	ht of the patie n creatinine co may increase m samples ha mg/dl	nts muscle oncentration when an a ve heterop 3.4-7.0 Desity.	e mass. A patient wa ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or URICASE
ample:Serum Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A	gher creatinine concentration. The concentration. Serum creatining ildly and may result in anomal <b>Hevels can be seen in the for</b> protein diet, alcohol), Chronic <b>HAGT</b> ), <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serum ne concentrations lous values if seru 5.94 5.94 bllowing: kidney disease, H	ht of the patie n creatinine co may increase m samples ha mg/dl ypertension, C U/L U/L IU/L	nts muscle oncentration when an ve heterop 3.4-7.0 Obesity.	e mass. A patient wo ons over time is mo ACE inhibitor (ACI hilic antibodies, he	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or URICASE
Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT)	gher creatinine concentration. The concentration. Serum creatining ildly and may result in anomal <b>Hevels can be seen in the for</b> protein diet, alcohol), Chronic <b>HAGT</b> ), <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serum ne concentrations lous values if seru 5.94 <b>5.94</b> <b>bllowing:</b> kidney disease, H <b>43.40</b> <b>59.70</b> 16.20	ht of the patie n creatinine co may increase m samples ha mg/dl	nts muscle oncentration when an a ve heterop 3.4-7.0 Desity. 2005 210 235 240 11-50	e mass. A patient with ons over time is mo ACE inhibitor (ACI hilic antibodies, he )	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or URICASE
ample:Serum Interpretation: The significance of mass will have a hig absolute creatinine of could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	gher creatinine concentration. The concentration. Serum creatining ildly and may result in anomal <b>Hevels can be seen in the for</b> protein diet, alcohol), Chronic <b>HAGT</b> ), <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serum te concentrations lous values if seru 5.94 <b>5.94</b> <b>5.94</b> <b>5.94</b> <b>5.94</b> <b>6.00</b> <b>59.70</b> 16.20 <b>7.37</b>	ht of the patien n creatinine co may increase m samples har mg/ dl Typertension, C U/L U/L U/L IU/L gm/ dl	nts muscle oncentration when an a ve heterop 3.4-7.0 Desity. Obesity. <35 <40 11-50 6.2-8.0	e mass. A patient wa ons over time is mo ACE inhibitor (ACI hilic antibodies, he )	ith a greater muscle re important than E) is taken. The assay molyzed, icteric or URICASE IFICCASE IFICCWITHOUT PSP IFICCWITHOUT PSP IFICCWITHOUT PSP OPTIMIZED SZAZING BURET









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 16:25:21
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	U	Init Bio. Ref. Int	erval Method
Alkaline Phosphatase (Total)	79.00	U/L	42.0-165.0	PNP/ AMP KINETIC
Bilirubin (Total)	0.83	mg/ dl	0.3-1.2	JENDRASSIK & GROF
Bilirubin (Direct)	0.31	mg/ dl	< 0.30	JENDRASSIK & GROF
Bilirubin (Indirect)	0.52	mg/ dl	<0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI), Serum				
Cholesterol (Total)	181.00	mg/ dl	<200 Desirable 200-239 Borderline I >240 High	CHOD-PAP High
HDL Cholesterol (Good Cholesterol)	51.50	mg/ dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	97	mg/ dl	< 100 Optimal 100-129 Nr. Optimal/ Above Opt 130-159 Borderline I 160-189 High > 190 Very High	
VLDL	32.24	mg/ dl	10-33	CALCULATED
Triglycerides	161.20	mg/dl	< 150 Normal 150-199 Borderline I 200-499 High >500 Very High	GPO-PAP High

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 17:01:45
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 17:36:35
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 19:15:01
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE, Urine	2			
Color	PALEYELOW			
Specific Gravity	1.020			
Reaction PH	Acidic (6.0)			DIPSTICK
Appearance	CLEAR			
Protein	TRACE	mg %	<10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) >500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	<0.5 (+) 0.5-1.0 (++) 1-2 (+++) >2 (++++)	DIPSTICK
Ketone	ABSENT	mg/ dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial œlls	0-2/h.p.f			MICROSCOPIC EXAMINATION
Puscells	0-2/h.p.f			
RBCs	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			

Urine Microscopy is done on centrifuged urine sediment.



View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 17:01:45
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 17:36:35
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 19:15:01
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

# DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
SUGAR, FASTING STAGE, Urine Sugar, Fasting stage	ABSENT	gms%		
Interpretation:   (+) < 0.5				
SUGAR, PP STAGE, Urine				
Sugar, PP Stage Interpretation: (+) < 0.5 gms% (++) 0.5-1.0 gms% (+++) 1-2 gms% (++++) > 2 gms%	ABSENT			

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 15:44:03
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF IMMUNOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
PSA (Prostate Specific Antigen), Total Sample:Serum	0.43	ng/mL	<4.1	ALD

#### Interpretation:

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone<sup>-</sup>
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

### THYROID PROFILE- TOTAL, Serum

T3, Total (tri-iodothyronine)	149.00	ng/ dl	84.61–201.7	CLIA
T4, Total (Thyroxine)	9.32	ug/ dl	3.2-12.6	CLIA
TSH (Thyroid Stimulating Hormone)	4.190	μlU/mL	0.27 - 5.5	ala

### Interpretation:

0.3-4.5	µIU/mL	First Trimest	er
0.5-4.6	µIU/mL	Second Trim	ester
0.8-5.2	µIU/mL	Third Trimes	ter
0.5-8.9	µIU/mL	Adults	55-87 Years
0.7-27	µIU/mL	Premature	28-36 Week
2.3-13.2	µIU/mL	Cord Blood	> 37Week
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)
1-39	µIU/mL	Child	0-4 Days
1.7-9.1	µIU/mL	Child	2-20 Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or





Page 10 of 13







Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 26/Oct/2024 12:51:24
UHID/MR NO	: ALDP.0000152994	Received	: 26/Oct/2024 14:05:48
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 15:44:03
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF IMMUNOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

**3**) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

**4**) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

**6)** In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

**8**) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 2024-10-26 12:51:05
UHID/MR NO	: ALDP.0000152994	Received	: 2024-10-26 12:51:05
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 15:50:10
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF X-RAY

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

### X-RAY DIGITAL CHEST PA

### <u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Poor inspiratory effort.
- Both lung field did not reveal any significant lesion.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Soft tissue shadow appears normal.
- Bony cage is normal.

Please correlare clinically.



Dr. Aishwarya Neha (MD Radiodiagnosis



()







Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.ABHAY SINGH	Registered On	: 26/Oct/2024 12:21:12
Age/Gender	: 34 Y 10 M 27 D / M	Collected	: 2024-10-26 13:45:43
UHID/MR NO	: ALDP.0000152994	Received	: 2024-10-26 13:45:43
Visit ID	: ALDP0284052425	Reported	: 26/Oct/2024 13:48:04
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

### DEPARTMENT OF ULTRASOUND

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

### ULTRASOUND WHOLE ABDOM EN (UPPER & LOWER)

**LIVER:** - Enlaregd in size (16.0 cm), with normal shape and shows diffusely raised echotexture. No focal lesion is seen. No intra hepatic biliary radicle dilation is seen.

GALL BLADDER :- Is not visualized (Post op).

**CBD** :- Normal in calibre at porta.

PORTAL VEIN: - Normal in calibre and colour uptake at porta.

**PANCREAS:** - Head is visualised, normal in size & echopattern. No evidence of ductal dilatation or calcification is seen. Rest of the pancreas is obscured by bowel gases.

SPLEEN: - Normal in size (10.0 cm), shape and echogenicity. No evidence of mass lesion is seen.

**RIGHT KIDNEY**: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

**LEFT KIDNEY**: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

**URINARY BLADDER :-** Is adequately distended. No evidence of calculus is seen. **Wall is thickened** (Maximum thickness 4.0 mm) and irregular.


PROSTATE :- Normal in size (2.3 x 2.9 x 3.1 cm vol - 11.4 cc), shape and echo pattern.

**HIGH RESOLUTION** :- No evidence of bowel loop dilatation or abnormal wall thickening is seen. No significant retroperitoneal lymphadenopathy is seen. No free fluid is seen in the abdomen/pelvis.

### **IMPRESSION**:

- Mild hepatomegaly with grade II fatty changes.
- Chronic cystitis.

### Please correlate clinically





08069366666

Chandan 24x7 App