



सत्यमेव जयते

भारत सरकार

Government of India



कोमल गुप्ता

Komal Gupta


जन्म तिथि / DOB : 05/10/1993

महिला / Female



8001 0566 8104

मेरा **आधार**, मेरी पहचान



भारत सरकार

Government of India



कोमल गुप्ता

Komal Gupta


जन्म तिथि / DOB : 05/10/1993

महिला / Female

Komal

DR. PIYUSH GOYAL
MBBS, DMR (Radiologist)
RMC No. 637041

~~80001~~ ~~05668~~ 8104



सेरा आधार, सेरी पहचान



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
- ☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



General Physical Examination

Date of Examination: 27/01/24

Name: KOMAL GUPTA Age: 30yrs DOB: 05/01/1993 Sex: Male

Referred By: BANK of BARODA

Photo ID: AADHAR CARD ID #: 8104

Ht: 167 (cm)

Wt: 75 (Kg)

Chest (Expiration): 95 (cm)

Abdomen Circumference: 107 (cm)

Blood Pressure: 125/85 mm Hg

PR: 79 / min

RR: 18 / min

Temp: Afebrile

BMI 26.9

Eye Examination:

RIE 6/6, N16, NCB
LE 6/6, N16, NCB

Other:

NO

On examination he/she appears physically and mentally fit: Yes / No

Signature Of Examinee : Komal Name of Examinee: KOMAL GUPTA

Signature Medical Examiner : Dr. PIYUSH GOYAL Name Medical Examiner Dr. PIYUSH GOYAL
MBBS, DMRD (Radiologist)
RMC No.-037041



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 27/01/2024 17:43:52

HAEMOGARAM

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
FULL BODY HEALTH CHECKUP BELOW 40 FEMAL			
HAEMOGLOBIN (Hb)	10.3 L	g/dL	12.0 - 15.0
TOTAL LEUCOCYTE COUNT	8.90	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	51.0	%	40.0 - 80.0
LYMPHOCYTE	44.0 H	%	20.0 - 40.0
EOSINOPHIL	2.0	%	1.0 - 6.0
MONOCYTE	3.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	4.42	$\times 10^6/\mu\text{L}$	3.80 - 4.80
HEMATOCRIT (HCT)	32.90 L	%	36.00 - 46.00
MEAN CORP VOLUME (MCV)	74.0 L	fL	83.0 - 101.0
MEAN CORP HB (MCH)	23.2 L	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	31.3 L	g/dL	31.5 - 34.5
PLATELET COUNT	184	$\times 10^3/\mu\text{L}$	150 - 410
RDW-CV	15.1 H	%	11.6 - 14.0

Technologist
VIKARANISJ
Page No: 1 of 17

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 27/01/2024 10:10:51

Final Authentication : 27/01/2024 17:43:52

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

Method:- Westergreen

16

mm in 1st hr

00 - 20

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases. ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein. ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as



Technologist
VIKARANTSIJ
Page No: 2 of 17

Tanu

DR. TANU RUNGTA

MD (Pathology)

RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-42234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance. and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan





P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA	Patient ID :-12234503	Date :- 27/01/2024	10:10:51
Age :- 30 Yrs 3 Mon 24 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Female	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 27/01/2024 17:43:52

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------


FASTING BLOOD SUGAR (Plasma) Method:- GOD POD	82.0	mg/dl	70.0 - 115.0
--	------	-------	--------------

Impaired glucose tolerance (IGT)	111 - 125 mg/dL
Diabetes Mellitus (DM)	> 126 mg/dL

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .

BLOOD SUGAR PP (Plasma) Method:- GOD PAP	90.6	mg/dl	70.0 - 140.0
---	------	-------	--------------

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .



Technologist
VIKARANTSIJ
Page No: 4 of 17

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA	Patient ID :-12234503	Date :- 27/01/2024	10:10:51
Age :- 30 Yrs 3 Mon 24 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Female	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 27/01/2024 17:43:52

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

GLYCOSYLATED HEMOGLOBIN (HbA1C)

Method:- CAPILLARY with EDTA

5.3 mg%

Non-Diabetic < 6.0
Good Control 6.0-7.0
Weak Control 7.0-8.0
Poor control > 8.0

MEAN PLASMA GLUCOSE

Method:- Calculated Parameter

104 mg/dL

68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in %

Non diabetic adults >=18 years < 5.7

At risk (Prediabetes) 5.7 - 6.4

Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings.

Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al]

1. Erythropoiesis

- Increased HbA1c: iron, vitamin B12 deficiency, decreased erythropoiesis.
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease.

2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.

3. Glycation

- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c: certain hemoglobinopathies, increased intra-erythrocyte pH

4. Erythrocyte destruction

- Increased HbA1c: increased erythrocyte life span: Splenectomy.
- Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

5. Others

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
- Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs

Technologist
VIKARANTSIJ
Page No: 5 of 17

DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axis Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 27/01/2024

10:10:51

Final Authentication : 27/01/2024 17:43:52

HAEMATOLOGY

BLOOD GROUP ABO

Method:- Haemagglutination reaction

"B" POSITIVE



Technologist
VIKARANTSI
Page No: 6 of 17

Tanu

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎️ +91 141 4824885 ✉️ maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA	Patient ID :-12234503	Date :- 27/01/2024	10:10:51
Age :- 30 Yrs 3 Mon 24 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Female	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 27/01/2024 17:43:52

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

LIPID PROFILE

TOTAL CHOLESTEROL Method:- CHOD-PAP methodology	131.00	mg/dl	Desirable <200 Borderline 200-239 High > 240
--	--------	-------	--

InstrumentName:MISPA PLUS Interpretation: Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES Method:- GPO-PAP	173.00 H	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
-----------------------------------	----------	-------	--

InstrumentName:Radox Rx Imola Interpretation : Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	36.50	mg/dl	MALE- 30-70 FEMALE - 30-85
--	-------	-------	-------------------------------

Instrument Name: Rx Daytona plus Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

LDL CHOLESTEROL Method:- Calculated Method	65.67	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
---	-------	-------	--

VLDL CHOLESTEROL Method:- Calculated	34.60	mg/dl	0.00 - 80.00
---	-------	-------	--------------

T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	3.59		0.00 - 4.90
--	------	--	-------------

LDL / HDL CHOLESTEROL RATIO Method:- Calculated	1.80		0.00 - 3.50
--	------	--	-------------

TOTAL LIPID Method:- CALCULATED	487.81	mg/dl	400.00 - 1000.00
------------------------------------	--------	-------	------------------

- 1 Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol
- 2 As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is

Technologist
VIKARAN JI
Page No: 7 of 17

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 27/01/2024 17:43:52

BIOCHEMISTRY

recommended

3 Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.



Technologist
VIKARANTSI
Page No: 8 of 17

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 27/01/2024 17:43:52

BIOCHEMISTRY

LIVER PROFILE WITH GGT

SERUM BILIRUBIN (TOTAL)

Method:- DMSO/Diazo

0.89 mg/dL

Infants : 0.2-8.0 mg/dL
Adult - Up to - 1.2 mg/dL

SERUM BILIRUBIN (DIRECT)

Method:- DMSO/Diazo

0.21 mg/dL

Up to 0.40 mg/dL

SERUM BILIRUBIN (INDIRECT)

Method:- Calculated

0.68 mg/dl

0.30-0.70

SGOT

Method:- IFCC

23.2 U/L

0.0 - 40.0

SGPT

Method:- IFCC

25.2 U/L

0.0 - 35.0

SERUM ALKALINE PHOSPHATASE

Method:- DGKC - SCE

100.20 U/L

42.00 - 110.00

SERUM GAMMA GT

Method:- Szasz methodology

Instrument Name Randox Rx Imola

Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and

metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

SERUM TOTAL PROTEIN

Method:- Direct Biuret Reagent

7.25 g/dl

6.00 - 8.40

SERUM ALBUMIN

Method:- Bromocresol Green

4.21 g/dl

3.50 - 5.50

SERUM GLOBULIN

Method:- CALCULATION

3.04 gm/dl

2.20 - 3.50

A/G RATIO

1.38

1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

Note :- These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B ,C ,paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.

Technologist
VIKARANTSIJ
Page No: 9 of 17


DR. TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 27/01/2024 17:43:52

BIOCHEMISTRY

RFT / KFT WITH ELECTROLYTES

SERUM UREA 26.30 mg/dl 10.00 - 50.00
Method:- Urease/GLDH

InstrumentName: HORIBA CA 60 Interpretation : Urea measurements are used in the diagnosis and treatment of certain renal and metabolic diseases.

SERUM CREATININE 0.99 mg/dl Males : 0.6-1.50 mg/dl
Method:- Jaffe's Method Females : 0.6 -1.40 mg/dl

Interpretation :

Creatinine is measured primarily to assess kidney function and has certain advantages over the measurement of urea. The plasma level of creatinine is relatively independent of protein ingestion, water intake, rate of urine production and exercise. Depressed levels of plasma creatinine are rare and not clinically significant.

SERUM URIC ACID 5.62 mg/dl 2.40 - 7.00

InstrumentName:HORIBA YUMIZEN CA60 Daytona plus Interpretation: Elevated Urate:High purine diet,Alcohol• Renal insufficiency,Drugs , Polycythaemia vera, Malignancies,Hypothyroidism,Rare enzyme defects ,Downs syndrome,Metabolic syndrome, Pregnancy,Gout.

SODIUM 140.2 mmol/L 135.0 - 150.0
Method:- ISE

POTASSIUM 4.21 mmol/L 3.50 - 5.50
Method:- ISE

CHLORIDE 98.9 mmol/L 94.0 - 110.0
Method:- ISE

SERUM CALCIUM 9.65 mg/dL 8.80 - 10.20
Method:- Arsenazo III Method

InstrumentName:MISPA PLUS Interpretation: Serum calcium levels are believed to be controlled by parathyroid hormone and vitamin D. Increases in serum PTH or vitamin D are usually associated with hypercalcemia .Hypocalcemia may be observed in hypoparathyroidism, nephrosis and pancreatitis.

SERUM TOTAL PROTEIN 7.25 g/dl 6.00 - 8.40
Method:- Direct Biuret Reagent

SERUM ALBUMIN 4.21 g/dl 3.50 - 5.50
Method:- Bromocresol Green

SERUM GLOBULIN 3.04 gm/dl 2.20 - 3.50
Method:- CALCULATION

A/G RATIO 1.38 1.30 - 2.50

Interpretation : Measurements obtained by this method are used in the diagnosis and treatment of a variety of dis liver, kidney and

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226

Technologist
VIKARANTSI
Page No: 10 of 17



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 27/01/2024

10:10:51

Final Authentication : 27/01/2024 17:43:52

BIOCHEMISTRY

bone marrow as well as other metabolic or nutritional disorders.

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR. In urine, it can remove the need for 24-hour collections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection. Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the blood increases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

Apart from renal failure Blood Urea can increase in dehydration and GI bleed



Technologist
VIKARANTSI
Page No: 11 of 17

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
+91 141 4824885 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 27/01/2024 17:43:52

TOTAL THYROID PROFILE

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

THYROID-TRIODOXYTHYRONINE T3

1.07

ng/mL

0.70 - 2.04

Method:- ECLIA

NOTE: In pregnancy total T3,T4 increase to 1.5 times the normal range.

Reference Range (T3): Premature Infants 26-30 Weeks ,3-4 days

0.24 - 1.32 ng/ml

Full-Term Infants 1-3 days

0.89 - 4.05 ng/ml

1 Week

0.91 - 3.00 ng/ml

1- 11 Months

0.85 - 2.50 ng/ml

Prepubertal Children

1.19 - 2.18 ng/ml

Reference Ranges (T4): Premature Infants 26-30 weeks ,3-4 days

2.60 - 14.0 ug/dl

Full -Term Infants 1-3 days

8.20 - 19.9 ug/dl

1 weeks 6.00 - 15.9 ug/dl 1-11 Months

6.10 - 14.9 ug/dl

Prepubertal children 12 months-2yrs

6.80 - 13.5 ug/dl

Prepubertal children 3-9 yrs

5.50 - 12.8 ug/dl

Reference Ranges (TSH): Premature Infants 26-32 weeks ,3-4 Days

0.80 - 6.9 uIU/ml

Full Term Infants 4 Days

1.36 - 16 uIU/ml

1 - 11 Months:0.90 - 7.70 | Prepubertal children:0.60 - 5.50.Primary malfunction of the thyroid gland may result in hyper or low release of T3 or T4

In additional as TSH directly affect thyroid function malfunction of the pituitary or the hypothalamus influences the thyroid gland activity. Disease in

any portion of the thyroid pituitary hypothalamus system may influence the level of T3 and T4 in the blood in Primary hypo thyroidism TSH levels

are significantly elevated while in secondary and tertiary hypothyroidism TSH levels may be low

THYROID THYROXINE (T4)

8.54

ug/dl

5.10 - 14.10

Method:- ECLIA

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration.Dose and time of drug intake also influence the test result. Transient increase in TSH levels or abnormal TSH levels can be seen in some non thyroidal conditions,simultaneous measurement of TSH with free T4 is useful in evaluating differential diagnosis

INTERPRETATION-Ultra Sensitive 4th generation assay 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with * TSH level.2.Low TSH,high FT4 and TSH receptor antibody(TRAb)

*ve seen in patients with Graves disease 3.Low TSH,high FT4 and TSH receptor antibody(TRAb)-ve seen in patients with Toxic adenoma/Toxic Multinodular goiter 4.HighTSH,Low FT4 and Thyroid microsomal

antibody increased seen in patients with Hashimotos thyroiditis 5.HighTSH,Low FT4 and Thyroid microsomal antibody normal seen in patients with Iodine deficiency/Congenital T4 syntheses deficiency 6.Low

TSH,Low FT4 and TRH stimulation test -Delayed response seen in patients with Tertiary hypothyroidism

7.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & 'serum TSH levels8.Normal T4 levels accompanied by ' T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis9.Normal or ' T3 & ' T

10.Normal T3 & T4 along with * TSH indicate mild / Subclinical Hyperthyroidism .11.Normal T3 & * T4 along with ' TSH is seen in Hypothyroidism .12.Normal T3 & T4 levels with ' TSH indicate Mild / Subclinical Hypoth

DURING PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association) 1st Trimester : 0.10-2.50 uIU/mL 2nd Trimester : 0.20-3.00 uIU/mL 3rd Trimester : 0.30-3.00

uIU/mL. The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

REMARK-Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while

thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test. Abnormal thyroid test findings often found in critically ill patients should be

repeated after the critical nature of the condition is resolved.TSH is an important marker for the diagnosis of thyroid dysfunction.Recent studies have shown that the TSH distribution progressively shifts to a higher

concentration with age ,and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly.

TSH

1.644

uIU/mL

0.350 - 5.500

Method:- ECLIA

DR.TANU RUNGTA

MD (Pathology)

RMC No. 17226

Technologist
VIKARAN TSI
Page No: 15 of 17



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

- 📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Date :- 27/01/2024

10:10:51

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Final Authentication : 27/01/2024 17:43:52

IMMUNOASSAY

4th Generation Assay,Reference ranges vary between laboratories

. PREGNANCY - REFERENCE RANGE for TSH IN uIU/mL (As per American Thyroid Association)

1st Trimester : 0.10-2.50 uIU/mL

2nd Trimester : 0.20-3.00 uIU/mL

3rd Trimester : 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circadian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- 1.Primary hyperthyroidism is accompanied by ↑serum T3 & T4 values along with ↓ TSH level.
- 2.Primary hypothyroidism is accompanied by ↓ serum T3 and T4 values & ↑serum TSH levels
- 3.Normal T4 levels accompanied by ↑ T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or ↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with ↓ TSH indicate mild / Subclinical Hyperthyroidism

. COMMENTS: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.

. Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction.Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age ,and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly

. Reference ranges are from Teitz fundamental of clinical chemistry 8th ed (2018)

Test performed by Instrument : Beckman coulter Dxi 800

. Note : The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease, it has to be correlated with clinical data for interpretation.

Technologist
VIKARANTSI
Page No: 16 of 17


DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 ✉ maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 27/01/2024

10:10:51

Final Authentication : 27/01/2024 17:43:52

PAP SMEAR

PAP SMEAR FOR CYTOLOGY EXAMINATION

Microscopic & diagnosis,

Smears are

No endocervical cells seen.

No atypical or malignant cells seen.

IMPRESSION :Inflammatory smears, (Negative for intraepithelial lesion or malignancy).

Adv: Clinical correlation.

Note: Please note papanicolaou smear study is a screening procedure for cervical cancer with inherent false negative result, hence should be interpreted with caution.

*** End of Report ***

Technologist
VIKARAN TSI
Page No. 17 of 17

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA	Patient ID :-12234503	Date :- 27/01/2024	10:10:51
Age :- 30 Yrs 3 Mon 24 Days	Ref. By Doctor:-BANK OF BARODA		
Sex :- Female	Lab/Hosp :-		
	Company :-	Mr.MEDIWHEEL	

Final Authentication : 27/01/2024 17:43:52

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
<u>PHYSICAL EXAMINATION</u>			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
<u>CHEMICAL EXAMINATION</u>			
REACTION(PH)	6.0		5.0 - 7.5
SPECIFIC GRAVITY	1.015		1.010 - 1.030
PROTEIN	NIL		NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL		NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
<u>MICROSCOPY EXAMINATION</u>			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT



Technologist
VIKARANTSI
Page No: 12 of 17


DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME :- Mrs. KOMAL GUPTA

Age :- 30 Yrs 3 Mon 24 Days

Sex :- Female

Patient ID :-12234503

Ref. By Doctor:-BANK OF BARODA

Lab/Hosp :-

Company :- Mr.MEDIWHEEL

Date :- 27/01/2024 10:10:51

Final Authentication : 27/01/2024 17:43:52

CLINICAL PATHOLOGY

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil



Technologist
VIKARANTSI
Page No: 13 of 17

DR.TANU RUNGTA
MD (Pathology)
RMC No. 17226



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



NAME:	MRS. KOMAL GUPTA	AGE	30 YRS/F
REF.BY	BANK OF BARODA	DATE	27/01/2024

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected

DR.SHALINI GOEL

M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954



P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

📍 B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
☎ +91 141 4824885 📧 maxcarediagnostics1@gmail.com



MRS. KOMAL GUPTA	Age : 30 Y/F
Registration Date: 27/01/2024	Ref. by: BANK OF BARODA

ULTRASOUND OF WHOLE ABDOMEN

Liver is of normal size (144 mm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intrahepatic biliary channels are not dilated. Portal vein diameter is normal.

Gall bladder is well distended. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

Kidneys are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

Right kidney is measuring approx. 103 mm.

Left kidney is measuring approx. 100 mm.

Urinary bladder does not show any calculus or mass lesion.

Uterus is anteverted and normal in size (measuring approx. 88 x 41 mm).

Myometrium shows normal echo -pattern. No focal space occupying lesion is seen. Endometrial echo is normal. Endometrial thickness is 3.6 mm.

Both ovaries are visualized and are normal. No adnexal mass lesion is seen.

No enlarged nodes are visualized. No retro-peritoneal lesion is identified.

No significant free fluid is seen in pouch of Douglas.

IMPRESSION: No significant abnormality is detected.

Dr. Mukesh Sharma
M.B.B.S; M.D. (Radiodiagnosis)
RMC No. 43418/17437

Dr. MUKESH SHARMA
M.B.B.S., M.D. (Radiodiagnosis)
RMC No. : 43418/17437
P3 Health Solutions LLP







P3 HEALTH SOLUTIONS LLP

(ASSOCIATES OF MAXCARE DIAGNOSTICS)

● B-14, Vidhyadhar Enclave-II, Near Axix Bank
Central Spine, Vidhyadhar Nagar, Jaipur - 302023
● +91 141 4824885 ● maxcarediagnostics1@gmail.com



MRS. KOMAL GUPTA	Age : 30 Y/F
Registration Date: 27/01/2024	Ref. by: BANK OF BARODA

2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY: FAIR TRANSTHORACIC ECHOCARDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL	TRICUSPID VALVE	NORMAL
AORTIC VALVE	NORMAL	PULMONARY VALVE	NORMAL

M.MODE EXAMINATION:

AO	2.4	Cm	LA	2.6	cm	IVS-D	0.9	cm
IVS-S	1.2	cm	LVID	4.5	cm	LVSD	2.7	cm
LVPW-D	0.9	cm	LVPW-S	1.2	cm	RV		cm
RVWT		cm	EDV		ml	LVVS		ml
LVEF	55-60%		RWMA			ABSENT		

CHAMBERS:


LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM		NORMAL	


COLOUR DOPPLER:

MITRAL VALVE				
E VELOCITY	0.90	m/sec	PEAK GRADIENT	Mm/hg
A VELOCITY	0.61	m/sec	MEAN GRADIENT	Mm/hg
MVA BY PHT		Cm2	MVA BY PLANIMETRY	Cm2
MITRAL REGURGITATION	ABSENT			
AORTIC VALVE				
PEAK VELOCITY	1.43	m/sec	PEAK GRADIENT	mm/hg
AR VMAX		m/sec	MEAN GRADIENT	mm/hg
AORTIC REGURGITATION	ABSENT			
TRICUSPID VALVE				
PEAK VELOCITY		m/sec	PEAK GRADIENT	mm/hg
MEAN VELOCITY		m/sec	MEAN GRADIENT	mm/hg
VMax VELOCITY				
TRICUSPID REGURGITATION	MILD			
PULMONARY VALVE				
PEAK VELOCITY	0.65	M/sec.	PEAK GRADIENT	Mm/hg
MEAN VELOCITY			MEAN GRADIENT	Mm/hg
PULMONARY REGURGITATION	ABSENT			

Impression—

- NORMAL LV SIZE & CONTRACTILITY.
- NO RWMA, LVEF 55-60%.
- MILD TR/ PAH (RVSP 27 MMHG+ RAP).
- NORMAL DIASTOLIC FUNCTION.
- NO CLOT, NO VEGETATION, NO PERICARDIAL EFFUSION.

(Cardiologist)

DR. JYOTI AGARWAL
 M.B.B.S, PGDCC (Cardiologist)
 RMC No.- 27255




Tems (P) Ltd

#P3 HEALTH SOLUTIONS LLP B-14, Vidhyadhar nahar , Jaipur
128541925460671/Mrs Komal Gupta 30Yrs/Female Kgs/31 Cms BP: ___/___ mmHg

Ref.: BANK OF BARODA Test Date: 27-Jan-2024(1:49:54 P) Notch: 50Hz 0.05Hz : 35Hz 10mm/mV 25mm/Sec

HR: 94 bpm

PR Interval: 114 ms
QRS Duration: 124 ms
QT/QTc: 377/473ms
P-QRS-T Axis: 59 - 56 - 11 (Deg)



FINDINGS: Normal Sinus Rhythm with Abnormal QTc Interval
Vent Rate : 94 bpm; PR Interval : 114 ms; QRS Duration : 124 ms; QT/QTc Int : 377/473 ms
P-QRS-T axis : 59 • 56 • 11 • (Deg)
Comments :

Sinus Rhythm with small q in lead I and aVL


Komal

Dr. Nareesh Kumar Mohanka
RMC NO: 35703
MBBS, DIP. RADIOLOGISTS
@NAREESH MOHANKA

This is informed to every staff member.

A summary of 2000 will be available in the office with below the sign.



 **GPS Map Camera**



Jaipur, Rajasthan, India

CANTEEN, SHANKARA EYE HOSPITAL, 6/168, Sector 2, Sector 6, Vidyadhar Nagar,
Jaipur, Rajasthan 302032, India

Lat 26.964543°

Long 75.782551°

27/01/24 11:16 AM GMT +05:30



R



12234503 KOMAL GUPTA 30 YRS , BOB F
27.JAN.2024
MAXCARE DIAGNOSTIC (ASSOCIATES OF P3 HEALTH SOLUTIONS LLP)

