Name	: Mr. BASKARAN K			
PID No.	: MED121778424	Register On : 3	1/03/2023 8:31 AM	~
SID No.	: 128623000926	Collection On : :	31/03/2023 10:20 AM	
Age / Sex	: 59 Year(s) / Male	Report On :	31/03/2023 5:05 PM	medall
Туре	: OP	Printed On : (	01/04/2023 12:22 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			
Investiga	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
BLOOD TYPINC	GROUPING AND Rh	'A' 'Positive'		
	ood/Agglutination)			
	<b>RETATION:</b> Reconfirm the Blood g	roup and Typing befor	e blood transfusion	
<u>Complet</u>	e Blood Count With - ESR			
Haemog (EDTA Bl	lobin 00d/Spectrophotometry)	14.3	g/dL	13.5 - 18.0
	Cell Volume(PCV)/Haematocrit ood/Derived from Impedance)	41.5	%	42 - 52
RBC Co (EDTA Bl	unt ood/Impedance Variation)	3.96	mill/cu.mm	4.7 - 6.0
	orpuscular Volume(MCV) ood/Derived from Impedance)	104.6	fL	78 - 100
	orpuscular Haemoglobin(MCH) ood/Derived from Impedance)	36.1	pg	27 - 32
concentr	orpuscular Haemoglobin ation(MCHC) ood/Derived from Impedance)	34.5	g/dL	32 - 36
RDW-C (EDTA Bl	V ood/Derived from Impedance)	13.9	%	11.5 - 16.0
RDW-SI (EDTA Bl	D ood/Derived from Impedance)	50.89	fL	39 - 46
	ukocyte Count (TC) ood/Impedance Variation)	6800	cells/cu.mm	4000 - 11000
Neutropl (EDTA Bl <i>Cytometry</i>	ood/Impedance Variation & Flow	63.8	%	40 - 75
Lympho (EDTA Bl <i>Cytometry</i> )	ood/Impedance Variation & Flow	24.5	%	20 - 45







APPROVED BY

The results pertain to sample tested.

Page 1 of 9

Name	: Mr. BASKARAN K			
PID No.	: MED121778424	Register On	: 31/03/2023 8:31 AM	
SID No.	: 128623000926	<b>Collection On</b>	: 31/03/2023 10:20 AM	
Age / Sex	: 59 Year(s) / Male	Report On	: 31/03/2023 5:05 PM	medall
Туре	: OP	Printed On	: 01/04/2023 12:22 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	4.4	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	6.9	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.4	%	00 - 02
INTERPRETATION: Tests done on Automated	Five Part cell count	er. All abnormal results	are reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	4.34	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.67	10^3 / µl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.30	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.47	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.03	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	218	10^3 / µl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	8.3	fL	7.9 - 13.7
PCT (EDTA Blood/Automated Blood cell Counter)	0.18	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	20	mm/hr	< 20







APPROVED BY

The results pertain to sample tested.

Page 2 of 9

Name	: Mr. BASKARAN K		
PID No.	: MED121778424	Register On : 31/03/2023 8:31 AM	
SID No.	: 128623000926	Collection On : 31/03/2023 10:20 AM	$\mathbf{O}$
Age / Sex	: 59 Year(s) / Male	Report On : 31/03/2023 5:05 PM	medall
Туре	: OP	Printed On : 01/04/2023 12:22 PM	DIAGNOSTICS

### Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
BUN / Creatinine Ratio	11.07		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	91.2	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	94.2	mg/dL	70 - 140

## **INTERPRETATION:**

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV/derived)	11.3	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	1.02	mg/dL	0.9 - 1.3

**INTERPRETATION:** Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/ <i>Enzymatic</i> )	5.5	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	1.20	mg/dL	0.1 - 1.2
Dr ANITHA Consultant Pathologist MCI -112788 VERIFIED BY			DR.SUNDAR ELAYAPERUMAL MD, CIC CONSULTANT MICROBIOLOGIST REG NO. 41854 APPROVED BY

The results pertain to sample tested.

Page 3 of 9

Name	: Mr. BASKARAN K	
PID No.	: MED121778424	Register On : 31/03/2023 8:31 AM
SID No.	: 128623000926	Collection On : 31/03/2023 10:20 AM
Age / Sex	: 59 Year(s) / Male	Report On : 31/03/2023 5:05 PM
Туре	: OP	Printed On : 01/04/2023 12:22 PM DIAG



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.25	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.95	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/ <i>Modified IFCC</i> )	15.6	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/ <i>Modified IFCC</i> )	16.1	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	18.0	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/ <i>Modified IFCC</i> )	42.3	U/L	56 - 119
Remark: Please correlate clinically.			
Total Protein (Serum/ <i>Biuret</i> )	7.49	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.20	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	3.29	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.28		1.1 - 2.2
<u>Lipid Profile</u>			

Cholesterol Total (Serum/CHOD-PAP with ATCS)

Ref. Dr

: MediWheel

mg/dL

Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240





198.4

DR.SUNDAR ELAYAPERUMAL MD, CIC CONSULTANT MICROBIOLOGIST REG NO. 41854

APPROVED BY

The results pertain to sample tested.

Page 4 of 9

Name	: Mr. BASKARAN K			
PID No.	: MED121778424	Register On	: 31/03/2023 8:31 AM	
SID No.	: 128623000926	<b>Collection On</b>	: 31/03/2023 10:20 AM	
Age / Sex	: 59 Year(s) / Male	Report On	31/03/2023 5:05 PM	medall
Туре	: OP	Printed On	: 01/04/2023 12:22 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			
<u>Investiga</u>	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Triglyce	rides	70.2	mg/dL	Optimal: < 150 Porderline: 150 100

(Serum/GPO-PAP with ATCS)

Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the õusualö"circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	46.9	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	137.5	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	14	mg/dL	< 30
Non HDL Cholesterol (Serum/ <i>Calculated</i> )	151.5	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.







APPROVED BY

The results pertain to sample tested.

Page 5 of 9

Name	: Mr. BASKARAN K		
PID No.	: MED121778424	Register On : 31/03/2023 8:31 AM	
SID No.	: 128623000926	Collection On : 31/03/2023 10:20 AM	$\mathbf{C}$
Age / Sex	: 59 Year(s) / Male	Report On : 31/03/2023 5:05 PM	edall
Туре	: OP		GNOSTICS
Ref. Dr	: MediWheel		

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Total Cholesterol/HDL Cholesterol Ratio (Serum/ <i>Calculated</i> )	4.2		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/ <i>Calculated</i> )	1.5		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.9		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
<u>Glycosylated Haemoglobin (HbA1c)</u>			
HbA1C (Whole Blood/ <i>HPLC</i> )	4.9	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control : 6.1 - 7.0 % , Fair control : 7.1 - 8.0 % , Poor control >= 8.1 %

Estimated Average Glucose 93.93 mg/dL

(Whole Blood)

## **INTERPRETATION:** Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.







APPROVED BY

The results pertain to sample tested.

Page 6 of 9

Name	: Mr. BASKARAN K					
PID No.	: MED121778424	Register On : 3	31/03/2023 8:31 AM			
SID No.	: 128623000926	-	31/03/2023 10:20 AM	$\mathbf{C}$		
Age / Sex	: 59 Year(s) / Male		31/03/2023 5:05 PM	medall		
Туре	: OP		01/04/2023 12:22 PM	DIAGNOSTICS		
Ref. Dr	: MediWheel	-				
Investiga	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval		
Prostate specific antigen - Total(PSA) (Serum/Manometric method)		0.29	ng/mL	Normal: 0.0 - 4.0 Inflammatory & Non Malignant conditions of Prostate & genitourinary system: 4.01 - 10.0 Suspicious of Malignant disease of Prostate: > 10.0		
INTERPI	RETATION: REMARK : PSA alone	e should not be used as	an absolute indicator of m	nalignancy.		
	ID PROFILE / TFT					
	odothyronine) - Total nemiluminescent Immunometric Assay	1.09	ng/ml	0.4 - 1.81		
(CLIA)) <b>INTERPRETATION:</b> <b>Comment :</b> Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.						
T4 (Tyre	oxine) - Total nemiluminescent Immunometric Assay	8.19	µg/dl	4.2 - 12.0		
<b>INTERPRETATION:</b> <b>Comment :</b> Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.						
	nyroid Stimulating Hormone) nemiluminescent Immunometric Assay	2.09	µIU/mL	0.35 - 5.50		
	Dr ANITHA sultant Pathologist MCI -112788 ERIFIED BY	■ <i>37時</i> で <i>変や</i> 3 いた (2月23) ■ 1155		DR.SUNDAR ELAYAPERUMAL MD, CIC CONSULTANT MICROBIOLOGIST REG NO. 41854		

APPROVED BY

The results pertain to sample tested.

Page 7 of 9

Name	: Mr. BASKARAN K			
PID No.	: MED121778424	Register On	: 31/03/2023 8:31 AM	~
SID No.	: 128623000926	<b>Collection On</b>	: 31/03/2023 10:20 AM	
Age / Sex	: 59 Year(s) / Male	Report On	: 31/03/2023 5:05 PM	medall
Туре	: OP	Printed On	: 01/04/2023 12:22 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			

**Investigation** <u>Observed</u> <u>Unit</u> **Biological** Value Reference Interval **INTERPRETATION:** 

Reference range for cord blood - upto 20 1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0 (Indian Thyroid Society Guidelines) **Comment :** 

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI. 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3.Values&amplt0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

# Urine Analysis - Routine

COLOUR (Urine)	Pale yellow		Yellow to Amber
APPEARANCE (Urine)	Clear		Clear
Protein (Urine/Protein error of indicator)	Negative		Negative
Glucose (Urine/GOD - POD)	Negative		Negative
Pus Cells (Urine/Automated ó"Flow cytometry)	Occasional	/hpf	NIL
Epithelial Cells (Urine/Automated ó"Flow cytometry )	Occasional	/hpf	NIL
RBCs (Urine/Automated ó"Flow cytometry )	NIL	/HPF	NIL
Casts (Urine/Automated ó"Flow cytometry )	NIL	/hpf	NIL
Crystals (Urine/Automated ó"Flow cytometry )	NIL	/hpf	NIL
Others (Urine)	NIL		







APPROVED BY

The results pertain to sample tested.

Page 8 of 9

Name	: Mr. BASKARAN K			
PID No.	: MED121778424	Register On	: 31/03/2023 8:31 AM	
SID No.	: 128623000926	<b>Collection On</b>	: 31/03/2023 10:20 AM	
Age / Sex	: 59 Year(s) / Male	Report On	: 31/03/2023 5:05 PM	medall
Туре	: OP	Printed On	: 01/04/2023 12:22 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			

Investigation

<u>Observed</u> <u>Unit</u> <u>Value</u> Biological Reference Interval

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.







APPROVED BY

-- End of Report --

The results pertain to sample tested.

Page 9 of 9