Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 09:46 AM
 Sample Receiving DATE
 : 23-Mar-2024 10:21 AM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 12:20 PM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 12:27 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

### **DEPARTMENT OF HAEMATOLOGY**

#### BLOOD GROUPING (ABO AND RH) (Specimen: EDTA)

Date	Status	23/Mar/24 12:27PM			Unit	<b>Bio Ref Interval</b>
Blood Group (aggultination method)		"B"				-
Rh Type (aggultination method)		POSITIVE				-

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 09:46 AM
 Sample Receiving DATE
 : 23-Mar-2024 10:21 AM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 11:17 AM

IPD No. / Ward : /

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

### DEPARTMENT OF BIOCHEMISTRY

Approved DATE

: 23-Mar-2024 11:35 AM

Blood Sugar Fasting\* (Specimen: FLUORIDE)

Date	Status	23/Mar/24 11:35AM			Unit	Bio Ref Interval
Blood Sugar Fasting		87.0			mg/dl	70-100

Patient NAME : Ms. POOJA GOENKA

IPD No. / Ward : / Approved DATE : 23-Mar-2024 11:42 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF HAEMATOLOGY

#### Complete Haemogram\* (Specimen : EDTA)

Date	Status	23/Mar/24 12:20PM			Unit	<b>Bio Ref Interval</b>
Haemoglobin (whole blood/photometric method)	L	11.1			g/dl	13.0-17
Total Leucocyte Count (TLC) (whole blood/impedence method)		4900			cells/c.mm	4000-10000
Neutrophil		58.8			%	45-70
Lymphocyte		30.9			%	20-40
Eosinophils	н	5.3			%	1.0-5.0
Monocytes		5.0			%	2.0-10.0
Basophils		0.0			%	0.0-1.0
Packed Cell Volume (PCV) (whole blood,calculation)	L	32.2			%	36-46
Red Blood Cell Count (whole blood,impedence method)		3.90			million/c.mm	3.8-4.8
Mean Cell Volume (MCV) (whole blood,calculated)		83.5			fl	83-101
Mean Cell Haemoglobin (MCH) (whole blood,calculated)		28.8			pg	27-32
MCHC (whole blood,calculated)		34.5			g/dl	31.5-34.5
RDW - CV		14.0			%	11.0-16.0
Platelet Count (whole blood,impedence method)		1.50			lakh/c.mm	1.5-4.0
MPV (Mean Platelet Volume)		11.4			fL	6.5-12.0
ESR	н	16			mm/Hr	0-15

### Interpretation:

Complete Haemogram\*: EDTA Whole Blood-Tests done on Automated Five Part Cell Counter.( Hb is performed by photometric method,WBC,RBC,Platelet Count by impedence method,WBC differential by Flow Cytometry technology other parameters calculated) All Abnormal Haemograms are reviewed confirmed microscopically.

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 09:46 AM
 Sample Receiving DATE
 : 23-Mar-2024 10:21 AM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 11:21 AM

 IPD No. / Ward
 : /
 Approved DATE
 : 23-Mar-2024 11:25 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF IMMUNOLOGY

#### Free Thyroid Profile (FT3, FT4, TSH) (Specimen: SERUM)

Date	Status	23/Mar/24 11:25AM			Unit	Bio Ref Interval
FT3		3.33			pg/ml	1.4-5.6
FT4		1.00			ng/dL	0.67-1.71
TSH		2.89			μIU/ml	0.25-5.00

Interpretation:

Free Thyroid Profile (FT3, FT4, TSH):

#### Interpretation:-

TSH	T3 / FT3	T4 / FT4	Suggested Interpretation for the Thyroid Function Tests Pattern
Within Range	Decreased	Within Range	. Isolated Low T3-often seen in elderly & associated Non-
Raised	Within Range	Within Range	Thyroidal illness. In elderly the drop in T3 level can be upto 25%.  .Isolated High TSH especially in the range of 4.7 to 15 mlU/ml is commonly associated with Physiological & Biological TSH Variability.  .Subclinical Autoimmune Hypothyroidism .Intermittent T4 therapy for hypothyroidism
Raised	Decreased	Decreased	.Recovery phase after Non-Thyroidal illness .Chronic Autoimmune Thyroiditis .Post thyroidectomy,Post radioiodine .Hypothyroid phase of transient thyroiditis
Raised or within Range	Raised	Raised or within Range	Interfering antibodies to thyroid hormones (anti-TPO antibodies) Intermittent T4 therapy or T4 overdose Drug interference- Amiodarone, Heparin,Beta blockers,steroids, anti-epileptics
Decreased	Raised or within Range	Raised or within Range	.Isolated Low TSH -especially in the range of 0.1 to 0.4 often seen in elderly & associated with Non-Thyroidal illness .Subclinical Hyperthyroidism .Thyroxine ingestion
Decreased	Decreased	Decreased	.Central Hypothyroidism .Non-Thyroidal illness .Recent treatment for Hyperthyroidism (TSH remains suppressed)
Decreased	Raised	Raised	.Primary Hyperthyroidism (Graves disease),Multinodular goitre, Toxic nodule .Transient thyroiditis:Postpartum, Silent (lymphocytic), Postviral (granulomatous,subacute, DeQuervains),Gestational thyrotoxicosis with hyperemesis gravidarum

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Ms. POOJA GOENKA

Sample Coll. DATE : 23-Mar-2024 09:46 AM Sample Receiving DATE : 23-Mar-2024 10:21 AM

UHID : 284776 Reporting DATE : 23-Mar-2024 11:21 AM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 11:25 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

### DEPARTMENT OF IMMUNOLOGY

Ì	Decreased or	Raised	Within Range	.T3 toxicosis
	within Range		9	.Non-Thyroidal illness

Patient NAME : Ms. POOJA GOENKA

Sample Coll. DATE : 23-Mar-2024 09:46 AM Sample Receiving DATE : 23-Mar-2024 10:21 AM

UHID : 284776 Reporting DATE : 23-Mar-2024 11:21 AM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 11:35 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

#### DEPARTMENT OF BIOCHEMISTRY

HbA1c (Specimen: EDTA)

Date	Status	23/Mar/24 11:35AM			Unit	Bio Ref Interval	
HbA1c		5.0			%	-<5.7	
AVERAGE BLOOD SUGAR		97.0			MG/DL	-<117	

Interpretation : HbA1c : Hba1c:

As per American Diabetes Association (ADA)							
Reference Group	HbA1c in %						
Non- diabetic adults	<5.7%						
Pre- diabetic	5.7-6.4 %						
Diabetic	>or = 6.5%						
ADA Target	>7.0						
Action suggested	>8.0						

Glycation is nonenzymatic addition of sugar residue to amino groups of proteins. HbA1C is formed by condensation of glucose with n-terminal valine residue of each beta chain of hb a to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of red blood cells(120 days) and the blood glucose concentration. the GHB concentration represents the integrated values for glucose over a period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasma glucose concentration in GHb depends on the time interval, with the most recent values providing a larger contribution than earlier values. The interpretation of GHb depends on RBC having normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb is been reported in iron deficiency anaemia.

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically. (\*) Test conducted under NABL scope MC-3302, Neo Hospital Laboratory, Noida.

Barcode No. : M316626 : M316626 : 32.0 YRS / Female

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 09:46 AM
 Sample Receiving DATE
 : 23-Mar-2024 10:21 AM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 11:17 AM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 11:35 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF BIOCHEMISTRY

#### KFT (Kidney Function Test)\* (Specimen: SERUM)

Date	Status	23/Mar/24 11:35AM		Unit	Bio Ref Interval
Blood Urea (urease with indicator dye)	L	9.0		mg/dl	15.0-37.0
Serum Creatinine (enzymatic(creatinine amidohydrolase))	L	0.5		mg/dl	0.52-1.04
Uric Acid (uricase/peroxidase)		3.7		mg/dl	2.5-6.2
Sodium (Na+) (direct ion selective mode)		139.0		mmol/L	137.0-145.0
Potassium (K+) (direct ion selective mode)		4.3		mmol/L	3.5-5.1
Chloride (CI-) (direct ion selective mode)		107.0		mmol/L	98.0-107.0
Serum Calcium (arsenazo dye)		9.1		mg/dl	8.4-10.2
Phosphorus Serum (phosphomolybdate reduction)		3.7		mg/dl	2.5-4.5
Alkaline Phosphatase (ALP) (4-nitrophenyl phosphate(pnpp)/amp)		76.0		U/L	38.0-126.0
Total protein (biuret(alkaline cupric sulphate))		7.3		gm/dl	6.3-8.2
Albumin (bromocresol green dye binding)		4.0		gm/dl	3.5-5.0
Albumin/Globulin Ratio (Calculated) (calculated)		1.2		Ratio	1.0-2.1
eGFR (calculated)		143.0		mL/min	-

### Lipid Profile\* (Specimen : SERUM)

Date	Status	23/Mar/24 11:35AM			Unit	<b>Bio Ref Interval</b>
Total Cholesterol (serum/enzymatic(che,cho/pod))		146.0			mg/dl	<200
Triglyceride (serum/enzymatic(lipase/gk/gpo/pod)without correction for free glycerol)		39.0			mg/dl	<150.0
HDL Cholesterol (serum/phosphotungstic acid/mgcl2+enzymatic)		52.0			mg/dl	>40.0
LDL		86.2			mg/dl	<100.0

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Ms. POOJA GOENKA

Sample Coll. DATE : 23-Mar-2024 09:46 AM Sample Receiving DATE : 23-Mar-2024 10:21 AM

UHID : 284776 Reporting DATE : 23-Mar-2024 11:17 AM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 11:35 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

#### DEPARTMENT OF BIOCHEMISTRY

(calculation)			
VLDL (calculation)	7.8	mg/dl <3	30
LDL/HDL Ratio (calculation)	1.66	<3	3.6
Total Cholesterol : HDL Ratio	2.81	</td <td>5.0</td>	5.0

# Interpretation :

NATIONAL LIPID ASSOCIATION RECOMMENDATIONS (NLA-2014)	TOTAL CHOLESTEROL in mg/dL	TRIGLYCERIDE in mg/dL	LDL CHOLESTEROL in mg/dL	NON HDL CHOLESTEROL in mg/dL
Optimal	<200	<150	<100	<130
Above Optimal	-	-	100-129	130 - 159
Borderline High	200-239	150-199	130-159	160 - 189
High	>=240	200-499	160-189	190 - 219
Very High		>=500	>=190	>=220

#### Note:

- 1. Measurements in the same patient can show physiological& analytical variations. Three serial samples 1 week apart are recommended for Total Cholesterol, Triglycerides, HDL& LDL Cholesterol.
- 2. As per NLA-2014 guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended.
- Low HDL levels are associated with increased risk forAtherosclerotic Cardiovascular disease (ASCVD) due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated from peripheral tissues.
- 4. NLA-2014identifies Non HDL Cholesterol(an indicator of all atherogeniclipoproteins such as LDL, VLDL, IDL, Lpa, Chylomicron remnants)along with LDL-cholesterol as co- primary target for cholesterol lowering therapy. Note that major risk factors can modify treatment goals for LDL &Non HDL.

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 09:46 AM
 Sample Receiving DATE
 : 23-Mar-2024 10:21 AM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 02:32 PM

IPD No. / Ward : /

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF CLINICAL PATHOLOGY

Approved DATE

: 23-Mar-2024 02:45 PM

Urine for Sugar Fasting\* (Specimen : URINE)

Date Status 23/Mar/24 Unit Bio Ref Interval 02:45PM Unit Fasting NIL -

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 09:46 AM
 Sample Receiving DATE
 : 23-Mar-2024 10:21 AM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 11:17 AM

IPD No. / Ward : /

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF BIOCHEMISTRY

Approved DATE

: 23-Mar-2024 11:35 AM

#### LFT PANEL (LIVER FUNCTION TEST) (Specimen : SERUM)

Date	Status	23/Mar/24 11:35AM			Unit	Bio Ref Interval
Bilirubin Total		0.6			mg/dl	0.2-1.3
Bilirubin Direct		0.2			mg/dl	0.0-0.3
Bilirubin Indirect		0.4			mg/dl	0.0-1.1
Aspartate Transaminase (SGOT, AST)		25.0			U/I	14.0-36.0
SGPT, ALT (Alanine Transaminase)		13.0			U/L	<35.0
Alkaline Phosphatase (ALP)		76.0			U/L	38.0-126.0
Total protein		7.3			gm/dl	6.3-8.2
Albumin		4.0			gm/dl	3.5-5.0
Globulin (Calculated)		3.3			gm/dl	2.0-3.5
Albumin/Globulin Ratio (Calculated)		1.2			Ratio	1.0-2.1
GGT (Gamma Glutamyl Transpeptidase)	L	10.0			U/L	12.0-43.0

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 12:26 PM
 Sample Receiving DATE
 : 23-Mar-2024 12:27 PM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 08:31 PM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 08:31 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

### DEPARTMENT OF CYTOLOGY

### PAP SMEAR REPORT

Smears are adequate for evaluation.

Endocervical cells are seen.

Benign reactive cellular changes associated with inflammation are not seen.

No protozoal or fungal elements are noted.

Background shows moderate acute inflammatory cells and numerous red blood cells.

Impression: Negative for intraepithelial lesion/malignancy

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Ms. POOJA GOENKA

 Sample Coll. DATE
 : 23-Mar-2024 11:51 AM
 Sample Receiving DATE
 : 23-Mar-2024 12:22 PM

 UHID
 : 284776
 Reporting DATE
 : 23-Mar-2024 05:23 PM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 06:18 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF CLINICAL PATHOLOGY

# **URINE ROUTINE**

SAMPLE: URINE

	OBSERVED VALUE	UNIT	REFERENCE RANGE	
PHYSICAL EXAMINATION		•		
VOLUME(visual observation)	30 mL		N/A	
COLOUR(visual observation)	PALE YELLOW		PALE YELLOW	
TRANSPARENCY (APPEARANCE)(visual observation)	S.TURBID		CLEAR	
SPECIFIC GRAVITY(automated multistrips,colour reaction/Pka change)	1.020		1.005 TO 1.030	
pH(automated multistrips double indicator method)	6.5		5-7	
CHEMICAL EXAMINATION				
PROTEIN (ALBUMIN)automated multistrips)protein error of pH),sulphosalicylic acid method.	NIL		NIL	
GLUCOSE(automated multistrips,(enzyme reaction) benedicts method	NIL		NIL	
KETONE BODIES(automated multistrips,rotheras method)	NEGATIVE		NEGATIVE	
BILIRUBIN(automated multistrips, fouchets method)	NEGATIVE		NEGATIVE	
UROBILINOGEN(automated multistrips,ehrlichs aldehyde method)	NORMAL		NORMAL (1mg/dL )	
BLOOD(automated multistrips ,bencidine method)	PRESENT		ABSENT	
MICROSCOPIC EXAMINATION				
PUS CELLS(light microscopy)	1-2	/hpf	0-5	
RED BLOOD CELLS(light microscopy)	0	/hpf	0-3	
EPITHELIAL CELLS(light microscopy)	15-20	/hpf	0-5	
CASTS(light microscopy)	ABSENT		ABSENT	
CRYSTALS(light microscopy)	ABSENT		ABSENT	

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Patient NAME : Ms. POOJA GOENKA

Sample Coll. DATE : 23-Mar-2024 11:51 AM Sample Receiving DATE : 23-Mar-2024 12:22 PM

UHID : 284776 Reporting DATE : 23-Mar-2024 05:23 PM

IPD No. / Ward : / Approved DATE : 23-Mar-2024 06:18 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

#### DEPARTMENT OF CLINICAL PATHOLOGY

OTHERS(light microscopy)	YEAST CELL PRESENT	-
--------------------------	--------------------	---

Note: 1. Chemical examination through Dipstick includes test methods as Protein(Protein Error Principle), Glucose (GOD-POD), Ketone(Legals Test), Bilirubin(Azo-Diazo reaction), Urobilinogen (Diazonium ion Reaction). All abnormal results of chemical examination are confirmed by manual methods.

- 2.Pre-test conditions to be observed while submitting the sample-First void,mid-stream urine, collect in a clean, dry, sterile container is recommended for routine urine analysis., avoid contamination with any discharge from vaginal ,urethra, perineum, as applicable , avoid prolonged transist time&undue exposure to sunlight.
- 3. During interpretation, Trace proteinuria can be seen with many physiological conditions like prolonged recumbency, excercise, high protein diet. False positive reactions for bile pigments, proteins, glucose can be caused by peroxidase like activity by disinfectants, therapeutic dyes, ascorbic acid and certain drugs.
- 4.All urine samples are checked for adequacy and suitability before examination.

Prepared By: Mr. Arvind Printed By: Mrs. Mala

These values are only indicative not confirmatory of diagnosis; Kindly correlate clinically.

Barcode No. : M316626 Age / Sex : 32.0 YRS / Female

Patient Name : Ms. POOJA GOENKA Registration Date : 23-Mar-2024 09:17 AM

IPD No. : Reporting Date : 23-Mar-2024 02:02 PM

UHID : 284776 Approved Date : 26-Mar-2024 10:26 AM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

#### DEPARTMENT OF CARDIOLOGY

# MITDAL VALVE

**ECHOCARDIOGRAPHY REPORT** 

MITRAL VALVE

 $Morphology \quad AML\textbf{-Normal/} Thickening/Calcification/Flutter/Vegetation/Prolapse/SAM/Doming. \\$ 

PML-Normal/Thickening/Calcification/Prolapes/Paradoxical motion/Fixed.

Subvalvular deformity Present/Absent. Score: \_\_\_\_\_

Doppler Normal/Abnormal E/A=99/56, E>A A>E S>D Mitral Stenosis Present/Absent RR Interval\_\_\_\_msec

EDG\_\_\_mmHg MDG\_\_\_mmHg MVA\_\_\_cm<sup>2</sup>

Mitral Regurgitation Absent/**Trivial**/Mild/Moderate/Severe.

TRICUSPID VALVE

Morphology **Normal**/Atresia/Thickening/Calcification/Prolapse/Vegetation/Doming.

Doppler Normal/Abnormal TRICSPID VALVE= 152cm/s.

Tricuspid Stenosis Present/**Absent** RR Interval\_\_\_\_msec.

EDG\_\_\_\_mmHg MDG\_\_\_\_mmHg

Tricuspid regurgitation Absent/**Trivial**/Mild/Moderate/Severe Fragmented Signals

Velocity\_\_\_\_msec Pred.RVSP =21+10mmHg

**PULMONARY VALVE** 

Morphology **Normal**/Atresia/Thickening/Doming/Vegetation

Doppler Normal/Abnormal PULMONARY VALVE= 64cm/s.

Pulmonary Stenosis Present/**Absent** Level

PSG\_\_\_\_mmHg Pulmonary annulus\_\_\_mm

Pulmonary regurgitation Present/**Absent** 

AORTIC VALVE

Morphology Normal/**Thickening**/Calcification/Restricted opening/Flutter/Vegetation

No. of cusps 1/2/3/4

Doppler Normal/Abnormal AORTIC VALVE= 132cm/s.

Aortic Stenosis Present/**Absent** Level

 $\begin{array}{lll} {\sf PSG}\underline{\qquad} {\sf mmHg} & {\sf Aortic\ annulus}\underline{\qquad} {\sf mm} \\ {\sf Aortic\ regurgitation} & {\sf Absent/Trivial/{\it Mild/Moderate/Severe}}. \end{array}$ 

Barcode No. Age / Sex : 32.0 YRS / Female : M316626

: Ms. POOJA GOENKA Patient Name Registration Date : 23-Mar-2024 09:17 AM

IPD No. Reporting Date : 23-Mar-2024 02:02 PM

**UHID** : 284776 Approved Date : 26-Mar-2024 10:26 AM

: Dr. Rakesh Malhotra (H) Referring Doctor

Passport No.

#### DEPARTMENT OF CARDIOLOGY

**Normal Valves Normal Valves** <u>Measurements</u> **Measurements** Aorta 2.4 (2.0-3.7 cm) LA es 3.3 (1.9-4.0 cm) LV es LV ed (3.7-5.6 cm) 3.5 (2.2-4.0 cm) 5.2 **IVSed** 1.2/1.8 (0.6-1.1 cm) PW (LV) 1.2/1.9 (0.6-1.1 cm) (0.7-2.6 cm) RV Anterior Wall **RVed** (upto 5 cm)

LVVd (ml) LVVs (ml)

EF 60% (54%-76%) IVS motion Normal/Flat/Paradoxical **IVS** Any Other

**CHAMBERS** 

Normal/Enlarged/Clear/Thrombus/Hypertrophy, Contraction,

Normal/Reduced/Regional wall motion abnormality: nil,

LA Normal/Enlarged/Clear/Thrombus Normal/Enlarged/Clear/Thrombus RA RV Normal/Enlarged/Clear/Thrombus **PERICARDIUM** Normal/Thickening/Calcification/Effusion

**COMMENTS & SUMMARY** 

No RWMA, LVEF-60% Normal LV systolic function

Mild AR/No AS

Trivial MR/TR(Normal PASP)

MIP=Normal Intact IAS/IVS No LA/LV clot

No pericardial effusion.

**IMPRESSION** 

Normal LV/RV systolic function

Mild AR

Barcode No. : M316626 Age / Sex : 32.0 YRS / Female

Patient Name : Ms. POOJA GOENKA Registration Date : 23-Mar-2024 09:17 AM

IPD No. : Reporting Date : 23-Mar-2024 12:19 PM

UHID : 284776 Approved Date : 23-Mar-2024 12:19 PM

Referring Doctor : **Dr. Rakesh Malhotra** (**H**)

Passport No. :

# DEPARTMENT OF RADIOLOGY

#### **USG WHOLE ABDOMEN**

Liver is normal in size, shape and echotexture. No focal SOL noted. Vascular channels are clear. No evidence of IHBR dilatation.

Gall Bladder is well distended and reveals normal walls. No evidence of calculus or mass lesion. CBD & PV are normal.

Spleen is normal in size, shape and echotexture.

Pancreas is normal in size, shape & echotexture.

Both Kidneys are normal in size, shape, position & echogenicity. CMD is maintained. No evidence of calculus or hydronephrosis.

Right kidney -10.0 x 4.3 cm

Left kidney - 11.3 x 4.4 cm

<u>Urinary Bladder</u> is well distended with normal wall thickness. No calculi / mass lesion noted. No diverticulum noted.

<u>Uterus</u> is normal in size, shape and echotexture. No focal lesion noted. Endometrial echo is normal (8.1 mm). Cervix is normal.

Both adnexa are clear.

No free fluid noted in peritoneal cavity.

#### IMPRESSION:

• NO SIGNIFICANT ABNORMALITY.

Please correlate clinically

Barcode No. : M316626 Age / Sex : 32.0 YRS / Female

Patient Name : Ms. POOJA GOENKA Registration Date : 23-Mar-2024 09:17 AM

IPD No. : Reporting Date : 23-Mar-2024 05:06 PM

UHID : 284776 Approved Date : 23-Mar-2024 05:06 PM

Referring Doctor : Dr. Rakesh Malhotra (H)

Passport No. :

# DEPARTMENT OF RADIOLOGY

#### X- RAY CHEST PA VIEW

Both lung fields are clear.

Hilar shadows are normal.

Both costophrenic angles are clear.

Cardiac silhouette is normal.

Bony thorax is normal.

Please correlate clinically

\*\*\* End Of Report \*\*\*

Dr. Vijay Singh Rawat DMRD,MD Radiodiagnosis Consultant Radiologist

Prepared By: Mr. Arvind

Dr. Sagar Tomar MD Radiodiagnosis, Fellow MSK MRI (Consultant Radiologist)

Dr. Rohit Kundra MD Radiodiagnosis (Consultant Radiologist) Dr. Harshita Tripathi MD Radiodiagnosis (Consultant Radiologist)

Printed By: Mrs. Mala