



Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:53
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 12:50:01
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 16:17:33
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

### DEPARTMENT OF HAEMATOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
i est ivallie	Kesuit	Unit	DIO. REI. IIILEI VAI	IVIELIIUU
Blood Group (ABO & Rh typing) **	, Blood			
Blood Group	0			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Rh ( Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Complete Blood Count (CBC) ** , u	Whole Blood			
Haemoglobin	14.00	g/dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRIC METHOD (CYANIDE-FREE REAGENT)
TLC (WBC)	6,400.00	/Cu mm	4000-10000	IMPEDANCE METHOD
DLC				
Polymorphs (Neutrophils )	57.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	33.00	%	20-40	FLOW CYTOMETRY
Monocytes	4.00	%	2-10	FLOW CYTOMETRY
Eosinophils	6.00	%	1-6	FLOW CYTOMETRY
Basophils	0.00	%	< 1-2	FLOW CYTOMETRY
ESR				
Observed	10.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5	



80-91 Yr 15.8








Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:53
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 12:50:01
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 16:17:33
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

### DEPARTMENT OF HAEMATOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	8.00	Mm for 1st hr.	< 9	
PCV (HCT)	45.00	%	40-54	
Platelet count				
Platelet Count	3.40	LACS/cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	14.00	fL	9-17	ELECTRONIC IMPEDANCE
P-LCR (Platelet Large Cell Ratio)	18.20	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.30	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	8.80	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBC Count				
RBC Count	4.38	Mill./cu mm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	97.70	fl	80-100	CALCULATED PARAMETER
MCH	32.10	pg	27-32	CALCULATED PARAMETER
MCHC	32.80	%	30-38	CALCULATED PARAMETER
RDW-CV	14.90	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	52.00	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	3,648.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	384.00	/cu mm	40-440	



View Reports on

Chandan 24x7 App









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 13:19:39
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 14:39:34
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

## DEPARTMENT OF BIOCHEMISTRY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
<b>GLUCOSE FASTING **</b> , <i>Plasma</i> Glucose Fasting	100.60	100-	0 Normal 125 Pre-diabetes 6 Diabetes	GOD POD

#### Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

**CLINICAL SIGNIFICANCE:-** Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP **	143.60	mg/dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

#### Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

#### GLYCOSYLATED HAEMOGLOBIN (HBA1C) \*\* , EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	6.20	% NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	44.00	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	131	mg/dl	

#### Interpretation:

#### NOTE:-

• eAG is directly related to A1c.



View Reports on

Chandan 24x7 App





T



# CHANDAN DIAGNOSTIC CENTRE

Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 13:19:39
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 14:39:34
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

### DEPARTMENT OF BIOCHEMISTRY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

est Name	Result	Unit	Bio. Ref. Interval	Method

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	<b>Degree of Glucose Control Unit</b>
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

\*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. \*\*Some danger of hypoglycemic reaction in Type 1 diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

#### **Clinical Implications:**

\*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

\*With optimal control, the HbA 1c moves toward normal levels.

\*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated \*Increases in glycosylated hemoglobin occur in the following nondiabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

\*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

\*Pregnancy d. chronic renal failure. Interfering Factors:

\*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood Urea Nitrogen) **	10.40	mg/dL	7.0-23.0	CALCULATED
Sample:Serum				



08069366666

Page 4 of 13







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name Age/Gender UHID/MR NO	: Mr.ADITYA PRAKASH : 29 Y 0 M 0 D /M : CALI.0000059174		Registered On Collected Received	: 28/Sep/2024 : 28/Sep/2024 : 28/Sep/2024	11:23:15 13:19:39
Visit ID Ref Doctor	: CALI0133872425 : Dr.Mediwheel - Arcofer	ni Health Care Ltd.	Reported Status	: 28/Sep/2024 : Final Report	14:39:34
Rei Doctoi	-		Status		
		DEPARTMENT ( IEEL BANK OF BA			
Test Name		Result	Unit	Bio. Ref. Interva	I Method
Interpretation: Note: Elevated BU	N levels can be seen in th	e following:			
High-protein diet, De	hydration, Aging, Certain m	edications, Burns, Ga	astrointestimal (Gl	) bleeding.	
Low BUN levels ca	n be seen in the following	:			
Low-protein diet, ove	erhydration, Liver disease.				
rootining **					
Creatinine **		0.74	mg/dl 0.7-	1.30	MODIFIED JAFFES
ample:Serum		0.74	mg/dl 0.7-	1.30	MODIFIED JAFFES
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic.	ingle creatinine value must b ner creatinine concentration. oncentration. Serum creatinin dly and may result in anoma	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s	of the patients mus reatinine concentra y increase when a samples have heter	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine cc could be affected mile	ner creatinine concentration.	e interpreted in light o The trend of serum cr ne concentrations ma	of the patients mus reatinine concentra y increase when a	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he	ith a greater muscle bre important than E) is taken. The assay
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ** Sample:Serum Interpretation:	ner creatinine concentration.	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s	of the patients mus reatinine concentra y increase when a samples have heter	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ** Sample:Serum Interpretation: Note:-	her creatinine concentration. oncentration. Serum creatinin dly and may result in anoma	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51	of the patients mus reatinine concentra y increase when a samples have heter	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or
<i>The significance of si mass will have a high absolute creatinine co could be affected mile lipemic.</i> <i>Jric Acid ** cample:Serum</i> <i>Interpretation:</i> <i>Note:-</i> <i>Elevated uric acid based and acid based and acid based acid bacid bacid based acid bacid based acid based acid ba</i>	her creatinine concentration. To oncentration. Serum creatining dly and may result in anoma dly and may result in anoma description of the seen in the features of the second sec	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 4.51	of the patients mus reatinine concentra y increase when a samples have heter mg/dl 3.4-	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid b	her creatinine concentration. oncentration. Serum creatinin dly and may result in anoma	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 4.51	of the patients mus reatinine concentra y increase when a samples have heter mg/dl 3.4-	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMMA	her creatinine concentration. To oncentration. Serum creatining dly and may result in anoma levels can be seen in the for otein diet, alcohol), Chronic A GT) ** , Serum	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 4.51	of the patients mus reatinine concentra y increase when a samples have heter mg/dl 3.4-	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or URICASE
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMMA SGOT / Aspartate Ar	her creatinine concentration. To oncentration. Serum creatinin dly and may result in anoma levels can be seen in the for otein diet, alcohol), Chronic A GT) ** , <i>Serum</i> minotransferase (AST)	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 <b>bllowing:</b> kidney disease, Hype 31.80	of the patients mus reatinine concentra y increase when a samples have heter mg/dl 3.4- ertension, Obesity U/L < 35	cle mass. A patient w ations over time is mo n ACE inhibitor (AC ophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or URICASE
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMIMA SGOT / Aspartate Ar SGPT / Alanine Amir	her creatinine concentration. To oncentration. Serum creatinin dly and may result in anoma levels can be seen in the for otein diet, alcohol), Chronic A GT) ** , <i>Serum</i> minotransferase (AST)	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 <b>ollowing:</b> kidney disease, Hype 31.80 <b>54.20</b>	of the patients must reatinine concentra y increase when a samples have heter mg/dl 3.4- ertension, Obesity U/L < 35 U/L < 40	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-proc FT (WITH GAMIMA SGOT / Aspartate Ar SGPT / Alanine Amir Gamma GT (GGT)	her creatinine concentration. To oncentration. Serum creatinin dly and may result in anoma levels can be seen in the for otein diet, alcohol), Chronic A GT) ** , <i>Serum</i> minotransferase (AST)	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 <b>bllowing:</b> kidney disease, Hype 31.80 <b>54.20</b> <b>57.70</b>	of the patients mus reatinine concentra y increase when a samples have heter mg/dl 3.4- ertension, Obesity U/L < 35 U/L < 40 IU/L 11-5	cle mass. A patient w ations over time is mo n ACE inhibitor (AC ophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mile lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-pro FT (WITH GAMIMA SGOT / Aspartate Ar SGPT / Alanine Amir Gamma GT (GGT) Protein	her creatinine concentration. To oncentration. Serum creatinin dly and may result in anoma levels can be seen in the fe otein diet, alcohol), Chronic A GT) ** , <i>Serum</i> minotransferase (AST)	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 <b>bllowing:</b> kidney disease, Hype 31.80 <b>54.20</b> <b>57.70</b> 6.82	of the patients must reatinine concentra y increase when a samples have heter mg/dl 3.4- ertension, Obesity U/L < 35 U/L < 40 IU/L 11-5 gm/dl 6.2-	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he 7.0	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING BIURET
Sample:Serum Interpretation: The significance of si mass will have a high absolute creatinine co could be affected mild lipemic. Jric Acid ** Sample:Serum Interpretation: Note:- Elevated uric acid I Drugs, Diet (high-proc FT (WITH GAMIMA SGOT / Aspartate Ar SGPT / Alanine Amir Gamma GT (GGT)	her creatinine concentration. To oncentration. Serum creatinin dly and may result in anoma levels can be seen in the fe otein diet, alcohol), Chronic A GT) ** , <i>Serum</i> minotransferase (AST)	e interpreted in light of The trend of serum cr ne concentrations ma lous values if serum s 4.51 <b>bllowing:</b> kidney disease, Hype 31.80 <b>54.20</b> <b>57.70</b>	of the patients mus reatinine concentra y increase when a samples have heter mg/dl 3.4- ertension, Obesity U/L < 35 U/L < 40 IU/L 11-5	cle mass. A patient w ations over time is mo n ACE inhibitor (AC rophilic antibodies, he 7.0 7.0 60 8.0 5.4	ith a greater muscle ore important than E) is taken. The assay emolyzed, icteric or URICASE IFCC WITHOUT P5P IFCC WITHOUT P5P OPTIMIZED SZAZING









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 13:19:39
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 14:39:34
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

# DEPARTMENT OF BIOCHEMISTRY

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	U	nit Bio. Ref. Int	erval Method
Alkaline Phosphatase (Total)	72.94	U/L	42.0-165.0	PNP/AMP KINETIC
Bilirubin (Total)	0.97	mg/dl	0.3-1.2	JENDRASSIK & GROF
Bilirubin (Direct)	0.41	mg/dl	< 0.30	JENDRASSIK & GROF
Bilirubin (Indirect)	0.56	mg/dl	< 0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI) ** , Serum				
Cholesterol (Total)	131.00	mg/dl	<200 Desirable 200-239 Borderline > 240 High	CHOD-PAP High
HDL Cholesterol (Good Cholesterol)	36.20	mg/dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	81	mg/dl	< 100 Optimal 100-129 Nr. Optimal/Above Opt 130-159 Borderline 160-189 High > 190 Very High	
VLDL	13.12	mg/dl	10-33	CALCULATED
Triglycerides	65.60	mg/dl	< 150 Normal 150-199 Borderline 200-499 High >500 Very High	GPO-PAP High

ISO 9001:2015

## Dr. Anupam Singh (MBBS MD Pathology)









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:53
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 15:13:01
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 18:57:19
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 19:49:36
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

# DEPARTMENT OF CLINICAL PATHOLOGY

### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE ** , $\iota$	Irine			
Color	PALE YELLOW			
Specific Gravity	1.010			
Reaction PH	Acidic (6.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	< 10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) > 500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	< 0.5 (+) 0.5-1.0 (++) 1-2 (+++) > 2 (++++)	DIPSTICK
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial cells	0-1/h.p.f			MICROSCOPIC EXAMINATION
Pus cells	ABSENT			
RBCs	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			
SUGAR, FASTING STAGE ** , Urine				
Sugar, Fasting stage	ABSENT	gms%		









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:53
Age/Gender	: 29 Y 0 M 0 D /M	Collected	: 28/Sep/2024 15:13:01
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 18:57:19
Visit ID	: CALI0133872425		: 28/Sep/2024 19:49:36
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

# DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

1112					
Test Name	Result	Unit	Bio. Ref. Interval	Method	
Interpretation:					

 $\begin{array}{ll} (+) & < 0.5 \\ (++) & 0.5\text{-}1.0 \\ (+++) & 1\text{-}2 \\ (++++) & > 2 \end{array}$ 

### SUGAR, PP STAGE \*\* , Urine

Sugar, PP Stage

ABSENT

#### Interpretation:

(+) < 0.5 gms% (++) 0.5-1.0 gms% (+++) 1-2 gms% (++++) > 2 gms%



Page 8 of 13

View Reports on

Chandan 24x7 App

Dr. Mamta Barthwal MD(Micro-Biology)









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 12:50:25
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 14:22:44
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

### DEPARTMENT OF IMMUNOLOGY

#### MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
<b>PSA (Prostate Specific Antigen)</b> , <b>Total</b> ** Sample:Serum	1.08	ng/mL	<4.1	CLIA

#### **Interpretation:**

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone<sup>-</sup>
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

#### THYROID PROFILE - TOTAL \*\* , Serum

T3, Total (tri-iodothyronine)	110.20	ng/dl	84.61–201.7	CLIA
T4, Total (Thyroxine)	8.30	ug/dl	3.2-12.6	CLIA
TSH (Thyroid Stimulating Hormone)	3.100	μlU/mL	0.27 - 5.5	CLIA

#### Interpretation:

0.3-4.5	µIU/mL	First Trimest	er
0.5-4.6	µIU/mL	Second Trim	ester
0.8-5.2	µIU/mL	Third Trimes	ter
0.5-8.9	µIU/mL	Adults	55-87 Years
0.7-27	µIU/mL	Premature	28-36 Week
2.3-13.2	µIU/mL	Cord Blood	> 37Week
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)
1-39	µIU/mL	Child	0-4 Days
1.7-9.1	µIU/mL	Child	2-20 Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or





Page 9 of 13







Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 28/Sep/2024 11:23:15
UHID/MR NO	: CALI.0000059174	Received	: 28/Sep/2024 12:50:25
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 14:22:44
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

### DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

**3**) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

**4**) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

**6)** In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

**8**) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr. Anupam Singh (MBBS MD Pathology)



Home Sample Collection 08069366666









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 2024-09-28 13:56:22
UHID/MR NO	: CALI.0000059174	Received	: 2024-09-28 13:56:22
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 13:56:37
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

# **DEPARTMENT OF X-RAY**

# MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

# X-RAY DIGITAL CHEST PA (500 mA COMPUTERISED UNIT SPOT FILM DEVICE)

# **DIGITAL CHEST P-A VIEW**

- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Pulmonary vascularity & distribution are normal.
- Pulmonary parenchyma did not reveal any significant lesion.

# IMPRESSION :

• NO SIGNIFICANT DIAGNOSTIC ABNORMALITY SEEN.

Dr. Pankaj Kumar Gupta (M.B.B.S D.M.R.D)









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
Age/Gender	: 29 Y O M O D /M	Collected	: 2024-09-28 12:27:11
UHID/MR NO	: CALI.0000059174	Received	: 2024-09-28 12:27:11
Visit ID	: CALI0133872425	Reported	: 28/Sep/2024 12:31:44
Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

## DEPARTMENT OF ULTRASOUND

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

## **ULTRASOUND WHOLE ABDOMEN (UPPER & LOWER)**

### WHOLE ABDOMEN ULTRASONOGRAPHY REPORT

### LIVER

• Liver is normal in size ~ 14.2 cm in longitudinal span and shows diffused raised echogenicity of hepatic parenchyma with loss of periportal echoes ..... S/O grade II fatty liver. No focal lesion is seen.

### PORTAL SYSTEM

- The intra hepatic portal channels are normal.
- The portal vein is not dilated.
- Porta hepatis is normal.

### BILIARY SYSTEM

- The intra-hepatic biliary radicles are normal.
- Common duct is not dilated.
- The gall bladder is normal in size and has regular walls. Lumen of the gall bladder is anechoic.

## PANCREAS

• The pancreas is normal in size and shape and has a normal homogenous echotexture. Pancreatic duct is not dilated.

## KIDNEYS

- Right kidney is normal in size ~ 9.6 x 4.2 cm position and cortical echotexture. Corticomedullary demarcation is maintained.
- Left kidney is normal in size ~ 9.8 x 4.6 cm position and cortical echotexture. Corticomedullary demarcation is maintained.
- A welldefined thinwalled simple cortical cyst measuring ~ 1.0 x 1.1 cm seen at upper pole region of left kidney.
- The collecting system of both the kidneys are not dilated.

## SPLEEN

• The spleen is normal in size ~ 9.9 cm and has a normal homogenous echo-texture.

## ILIAC FOSSAE & PERITONEUM

- Scan over the iliac fossae does not reveal any fluid collection or mass.
- No free fluid is noted in peritoneal cavity.
- Visualized bowel loops are gaseous and grossly appear normal in caliber,









Add: B 1/2, Sector J, Near Sangam Chauraha, LDA Stadium Road, ALIGANJ Ph: 9235432681 CIN: U85110UP2003PLC193493

1				
	Patient Name	: Mr.ADITYA PRAKASH	Registered On	: 28/Sep/2024 11:21:54
	Age/Gender	: 29 Y O M O D /M	Collected	: 2024-09-28 12:27:11
	UHID/MR NO	: CALI.0000059174	Received	: 2024-09-28 12:27:11
		: CALI0133872425	Reported	: 28/Sep/2024 12:31:44
	Ref Doctor	: Dr.Mediwheel - Arcofemi Health Care Ltd.	Status	: Final Report

## **DEPARTMENT OF ULTRASOUND**

## MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

peristalsis and wall thickness.

#### URINARY BLADDER

• The urinary bladder is normal. Bladder wall is normal in thickness and is regular. No calculus is seen.

#### PROSTATE

The prostate gland is normal in size with smooth outline (volume ~ 18.6 cc).

#### FINAL IMPRESSION

- LEFT RENAL SIMPLE CORTICAL CYST.
- GRADE II FATTY INFILTRATION OF LIVER.

#### Adv: Clinico-pathological correlation and follow-up.

\*\*\* End Of Report \*\*\*

(\*\*) Test Performed at Chandan Speciality Lab.

Result/s to Follow:

STOOL, ROUTINE EXAMINATION, ECG / EKG, Tread Mill Test (TMT)





This report is not for medico legal purpose. If clinical correlation is not established, kindly repeat the test at no additional cost within seven days

Facilities: MRI, CT scan, DR X-ray, Ultrasound, Sonomammography, Digital Mammography, ECG (Bedside also), 2D Echo, TMT, Holter, OPG, EEG, NCV, EMG & BERA, Audiometry, BMD, PFT, Fibroscan, Bronchoscopy, Colonoscopy and Endoscopy, Allergy Testing, Biochemistry & Immunoassay, Hematology, Microbiology & Serology, Histopathology & Immunohistochemistry, Cytogenetics and Molecular Diagnostics and Health Checkups 365 Days Open

\*Facilities Available at Select Location Page 13 of 13



**Home Sample Collection** View Reports on 08069366666 Chandan 24x7 App

