

Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

**Reg. Location**: Khar West (Main Centre)



R

E

Use a QR Code Scanner Application To Scan the Code

Collected : 21-Jun-2023 / 08:04 Reported : 21-Jun-2023 / 11:10

### MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO

CBC (Complete Blood Count), Blood				
<u>PARAMETER</u>	<u>RESULTS</u>	<b>BIOLOGICAL REF RANGE</b>	<u>METHOD</u>	
RBC PARAMETERS				
Haemoglobin	14.3	13.0-17.0 g/dL	Spectrophotometric	
RBC	5.23	4.5-5.5 mil/cmm	Elect. Impedance	
PCV	43.3	40-50 %	Calculated	
MCV	82.6	81-101 fl	Measured	
MCH	27.3	27-32 pg	Calculated	
MCHC	33.0	31.5-34.5 g/dL	Calculated	
RDW	13.6	11.6-14.0 %	Calculated	
WBC PARAMETERS				
WBC Total Count	5720	4000-10000 /cmm	Elect. Impedance	
WBC DIFFERENTIAL AND ABS	OLUTE COUNTS			
Lymphocytes	33.9	20-40 %		
Absolute Lymphocytes	1940	1000-3000 /cmm	Calculated	
Monocytes	9.8	2-10 %		
Absolute Monocytes	560	200-1000 /cmm	Calculated	
Neutrophils	50.5	40-80 %		
Absolute Neutrophils	2880	2000-7000 /cmm	Calculated	
Eosinophils	5.3	1-6 %		
Absolute Eosinophils	300	20-500 /cmm	Calculated	
Basophils	0.5	0.1-2 %		
Absolute Basophils	30	20-100 /cmm	Calculated	
Immature Leukocytes	-			

WBC Differential Count by Absorbance & Impedance method/Microscopy.

### PLATELET PARAMETERS

Platelet Count	237000	150000-410000 /cmm	Elect. Impedance		
MPV	9.1	6-11 fl	Measured		
PDW	16.7	11-18 %	Calculated		

### **RBC MORPHOLOGY**

Hypochromia Microcytosis -



Macrocytosis

CID : 2317209024

Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : Reg. Location : Khar West (Main Centre)

-

Anisocytosis -

Poikilocytosis -

Polychromasia -

Target Cells -

Basophilic Stippling -

Normoblasts -

Others Normocytic, Normochromic

WBC MORPHOLOGY

PLATELET MORPHOLOGY -

COMMENT -

Specimen: EDTA Whole Blood

ESR, EDTA WB-ESR 5 2-20 mm at 1 hr. Sedimentation

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*

lac-MRA



Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist & Lab Director

Anto

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

:21-Jun-2023 / 08:04

:21-Jun-2023 / 11:32

Collected

Reported

Page 2 of 13



CID : 2317209024

Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr.

Reg. Location : Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

: 21-Jun-2023 / 08:04 :21-Jun-2023 / 16:31

Hexokinase

Hexokinase

### MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO

**PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD** 

GLUCOSE (SUGAR) FASTING. 92 Non-Diabetic: < 100 mg/dl Fluoride Plasma Impaired Fasting Glucose:

100-125 mg/dl

Diabetic: >/= 126 mg/dl

Collected

Reported

GLUCOSE (SUGAR) PP, Fluoride 107 Non-Diabetic: < 140 mg/dl

Plasma PP/R Impaired Glucose Tolerance:

140-199 mg/dl

Diabetic: >/= 200 mg/dl

Urine Sugar (Fasting) **Absent Absent** Urine Ketones (Fasting) Absent Absent

Urine Sugar (PP) Absent **Absent** Urine Ketones (PP) Absent Absent

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab \*\*\* End Of Report \*\*\*





Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist & Lab Director

Anto

Page 3 of 13



CID : 2317209024

Name : MR.KISHOR BENDORI

: 60 Years / Male Age / Gender

Consulting Dr.

Reg. Location : Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected :21-Jun-2023 / 08:04 Reported

:21-Jun-2023 / 14:10

### MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO **KIDNEY FUNCTION TESTS**

<u>PARAMETER</u>	RESULTS	BIOLOGICAL REF RANGE	<u>METHOD</u>
BLOOD UREA, Serum	45.2	19.29-49.28 mg/dl	Calculated
BUN, Serum	21.1	9.0-23.0 mg/dl	Urease with GLDH
CREATININE, Serum	0.98	0.60-1.10 mg/dl	Enzymatic
eGFR, Serum	83	>60 ml/min/1.73sqm	Calculated
Note: eGFR estimation is calculated	using MDRD (Modification of die	et in renal disease study group) equ	ation
TOTAL PROTEINS, Serum	6.7	5.7-8.2 g/dL	Biuret
ALBUMIN, Serum	4.2	3.2-4.8 g/dL	BCG
GLOBULIN, Serum	2.5	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	1.7	1 - 2	Calculated
URIC ACID, Serum	5.6	3.7-9.2 mg/dl	Uricase/ Peroxidase
PHOSPHORUS, Serum	3.6	2.4-5.1 mg/dl	Phosphomolybdate
CALCIUM, Serum	8.7	8.7-10.4 mg/dl	Arsenazo
SODIUM, Serum	139	136-145 mmol/l	IMT
POTASSIUM, Serum	4.3	3.5-5.1 mmol/l	IMT
CHLORIDE, Serum	106	98-107 mmol/l	IMT

<sup>\*</sup>Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab \*\*\* End Of Report \*\*\*









Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

**Reg. Location**: Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

:21-Jun-2023 / 08:04

**Reported** :21-Jun-2023 / 13:03

## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO GLYCOSYLATED HEMOGLOBIN (HbA1c)

# PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD Glycosylated Hemoglobin 5.7 Non-Diabetic Level: < 5.7 %</td> HPLC

Glycosylated Hemoglobin 5.7 Non-Diabetic Level: < 5.7 % H (HbA1c), EDTA WB - CC Prediabetic Level: 5.7-6.4 %

Diabetic Level: >/= 6.5 %

Collected

Estimated Average Glucose 116.9 mg/dl Calculated (eAG), EDTA WB - CC

Intended use:

- In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year
- In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly
- For microvascular disease prevention, the HbA1C goal for non pregnant adults in general is Less than 7%.

Clinical Significance:

- HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.
- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

Factors affecting HbA1c results:

**Increased in:** High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, Splenectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.

Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies

Reflex tests: Blood glucose levels, CGM (Continuous Glucose monitoring)

References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*





Dr.JYOT THAKKER M.D. (PATH), DPB Pathologist & AVP( Medical Services)

Page 5 of 13



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

**Reg. Location**: Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

: 21-Jun-2023 / 08:04

**Reported** :21-Jun-2023 / 13:18

Collected

## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO PROSTATE SPECIFIC ANTIGEN (PSA)

PARAMETER RESULTS BIOLOGICAL REF RANGE METHOD

TOTAL PSA, Serum 1.675 <4.0 ng/ml CLIA

#### Clinical Significance:

- PSA is detected in the serum of males with normal, benign hyper-plastic, and malignant prostate tissue.
- Monitoring patients with a history of prostate cancer as an early indicator of recurrence and response to treatment.
- Prostate cancer screening 4.The percentage of Free PSA (FPSA) in serum is described as being significantly higher in patients with BPH
  than in patients with prostate cancer. 5.Calculation of % free PSA (ie. FPSA/TPSA x 100), has been suggested as way of improving the
  differentiation of BPH and Prostate cancer.

#### Interpretation:

Increased In- Prostate diseases, Cancer, Prostatitis, Benign prostatic hyperplasia, Prostatic ischemia, Acute urinary retention, Manipulations like Prostatic massage, Cystoscopy, Needle biopsy, Transurethral resection, Digital rectal examination, Radiation therapy, Indwelling catheter, Vigorous bicycle exercise, Drugs (e.g., testosterone), Physiologic fluctuations. Also found in small amounts in other cancers (sweat and salivary glands, breast, colon, lung, ovary) and in Skene glands of female urethra and in term placenta, Acute renal failure, Acute myocardial infarction.

Decreased In- Ejaculation within 24-48 hours, Castration, Antiandrogen drugs (e.g., finasteride), Radiation therapy, Prostatectomy, PSA falls 17% in 3 days after lying in hospital, Artifactual (e.g., improper specimen collection; very high PSA levels). Finasteride (5-α reductase inhibitor) reduces PSA by 50% after 6 months in men without cancer.

Reflex Tests: % FREE PSA, USG Prostate

#### Limitations:

- tPSA values determined on patient samples by different testing procedures cannot be directly compared with one another and could be the cause of erroneous medical interpretations. If there is a change in the tPSA assay procedure used while monitoring therapy, then the tPSA values obtained upon changing over to the new procedure must be confirmed by parallelmeasurements with both methods. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels.
- Patients who have been regularly exposed to animals or have received immunotherapy or diagnostic procedures utilizing immunoglobulins or immunoglobulin fragments may produce antibodies, e.g. HAMA, that interferes with immunoassays.
- PSA results should be interpreted in light of the total clinical presentation of the patient, including: symptoms, clinical history, data from additional tests, and other appropriate information.
- Serum PSA concentrations should not be interpreted as absolute evidence for the presence or absence of prostate cancer.

#### Reference:

- Wallach's Interpretation of diagnostic tests
- Total PSA Pack insert

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*





Dr.ANUPA DIXIT
M.D.(PATH)
Consultant Pathologist & Lab Director

Annha

Page 6 of 13



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

**Reg. Location**: Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected

Reported

pplication To Scan the Code: 21-Jun-2023 / 08:04

:22-Jun-2023 / 13:17

R

E

## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO EXAMINATION OF FAECES

### <u>PARAMETER</u> <u>RESULTS</u> <u>BIOLOGICAL REF RANGE</u>

### PHYSICAL EXAMINATION

ColourBrownBrownForm and ConsistencySemi SolidSemi SolidMucusAbsentAbsentBloodAbsentAbsent

**CHEMICAL EXAMINATION** 

Reaction (pH) Acidic (6.5) -

Occult Blood Absent Absent

#### MICROSCOPIC EXAMINATION

Protozoa Absent Absent Flagellates Absent Absent Ciliates Absent Absent **Parasites** Absent Absent Macrophages Absent Absent Mucus Strands Absent Absent Fat Globules Absent Absent RBC/hpf Absent Absent WBC/hpf Absent Absent Yeast Cells Absent **Absent Undigested Particles** Present ++ Concentration Method (for ova) No ova detected Absent Reducing Substances Absent





Dr.TRUPTI SHETTY M. D. (PATH) Pathologist

Page 7 of 13

<sup>\*</sup>Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : Reg. Location : Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected : 21-Jun-2023 / 08:04 Reported : 21-Jun-2023 / 12:43

## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO URINE EXAMINATION REPORT

<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Yellow	Pale Yellow	-
5.0	4.5 - 8.0	Chemical Indicator
1.025	1.001-1.030	Chemical Indicator
Clear	Clear	-
10	-	-
Absent	Absent	pH Indicator
Absent	Absent	GOD-POD
Absent	Absent	Legals Test
Absent	Absent	Peroxidase
Absent	Absent	Diazonium Salt
Normal	Normal	Diazonium Salt
Absent	Absent	Griess Test
<u>N</u>		
1-2	0-5/hpf	
Absent	0-2/hpf	
0-1		
Absent	Absent	
Absent	Absent	
Absent	Absent	
5-6	Less than 20/hpf	
-		
	Yellow 5.0 1.025 Clear 10  Absent Absent Absent Absent Normal Absent  V  1-2 Absent 0-1 Absent Absent Absent Absent	Yellow 5.0 4.5 - 8.0 1.025 Clear Clear Clear 10 - Absent

Interpretation: The concentration values of Chemical analytes corresponding to the grading given in the report are as follows:

- Protein: (1+ ~25 mg/dl, 2+ ~75 mg/dl, 3+ ~ 150 mg/dl, 4+ ~ 500 mg/dl)
- Glucose:(1+ ~ 50 mg/dl, 2+ ~100 mg/dl, 3+ ~300 mg/dl,4+ ~1000 mg/dl)
- Ketone:(1+ ~5 mg/dl, 2+ ~15 mg/dl, 3+ ~ 50 mg/dl, 4+ ~ 150 mg/dl)

Reference: Pack insert

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*





Dr.TRUPTI SHETTY M. D. (PATH) Pathologist

Page 8 of 13



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

**Reg. Location**: Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected

Reported

: 21-Jun-2023 / 08:04

:21-Jun-2023 / 12:42

## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO BLOOD GROUPING & Rh TYPING

<u>PARAMETER</u> <u>RESULTS</u>

ABO GROUP B

Rh TYPING Positive

NOTE: Test performed by automated Erythrocytes magnetized technology (EMT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

#### Clinical significance:

ABO system is most important of all blood group in transfusion medicine

#### Limitations:

- ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- · Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

#### Refernces:

- 1. Denise M Harmening, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia
- 2. AABB technical manual

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*





Dr.TRUPTI SHETTY M. D. (PATH) Pathologist

Page 9 of 13



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

Reg. Location

: Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

:21-Jun-2023 / 08:04

**Reported** :21-Jun-2023 / 14:10

Collected

### MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO LIPID PROFILE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
CHOLESTEROL, Serum	172.3	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >/=240 mg/dl	CHOD-POD
TRIGLYCERIDES, Serum	167.0	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high:>/=500 mg/dl	Enzymatic colorimetric
HDL CHOLESTEROL, Serum	26.6	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Elimination/ Catalase
NON HDL CHOLESTEROL, Serum	145.7	Desirable: <130 mg/dl Borderline-high:130 - 159 mg/dl High:160 - 189 mg/dl Very high: >/=190 mg/dl	Calculated
LDL CHOLESTEROL, Serum	112.3	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >/= 190 mg/dl	Calculated
VLDL CHOLESTEROL, Serum	33.4	< /= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO, Serum	6.5	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO, Serum	4.2	0-3.5 Ratio	Calculated

<sup>\*</sup>Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*







Page 10 of 13



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. : -

Reg. Location

: Khar West (Main Centre)

Authenticity Check

R

E

Use a QR Code Scanner Application To Scan the Code

**Collected** : 21-Jun-2023 / 08:04

**Reported** :21-Jun-2023 / 14:04

## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO THYROID FUNCTION TESTS

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Free T3, Serum	5.1	3.5-6.5 pmol/L	CLIA
Free T4, Serum	14.9	11.5-22.7 pmol/L	CLIA
sensitiveTSH, Serum	2.798	0.55-4.78 microIU/ml mIU/ml	CLIA



CID : 2317209024

Name : MR.KISHOR BENDORI

: 60 Years / Male Age / Gender

Consulting Dr. Collected : 21-Jun-2023 / 08:04

Reported Reg. Location : Khar West (Main Centre) :21-Jun-2023 / 14:04

#### Interpretation:

A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

#### Clinical Significance:

- 1)TSH Values between high abnormal upto15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological
- can give falsely high TSH.
- 2)TSH values may be trasiently altered becuase of non thyroidal illness like severe infections, liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3 / T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone, Recovery phase of non-thyroidal illness, TSH Resistance.
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radio iodine Rx, post thyroidectomy, Anti thyroid drugs, tyrosine kinase inhibitors & amiodarone, amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hyperthyroidism, Graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum, hydatiform mole)
Low	Normal	Normal	Subclinical Hyperthyroidism, recent Rx for Hyperthyroidism, drugs like steroids & dopamine), Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal Illness, Recent Rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, Drug interference: Amiodarone, Heparin, Beta Blockers, steroids & anti epileptics.

Diurnal Variation: TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am, and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation:19.7%(with in subject variation)

Reflex Tests: Anti thyroid Antibodies, USG Thyroid , TSH receptor Antibody. Thyroglobulin, Calcitonin

#### Limitations:

- 1. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.
- 2. Patient samples may contain heterophilic antibodies that could react in immunoassays to give falsely elevated or depressed results. this assay is designed to minimize interference from heterophilic antibodies.

- 1.O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
- Interpretation of the thyroid function tests, Dayan et al. THE LANCET. Vol 357
- 3. Tietz , Text Book of Clinical Chemistry and Molecular Biology -5th Edition
- 4. Biological Variation: From principles to Practice-Callum G Fraser (AACC Press)

\*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab \*\*\* End Of Report \*\*\*





Dr.ANUPA DIXIT M.D.(PATH) Consultant Pathologist & Lab Director

Anto

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code

Page 12 of 13



Name : MR.KISHOR BENDORI

Age / Gender : 60 Years / Male

Consulting Dr. :-

**Reg. Location**: Khar West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected

Reported

: 21-Jun-2023 / 08:04

:21-Jun-2023 / 14:10


## MEDIWHEEL FULL BODY HEALTH CHECKUP MALE ABOVE 40/2D ECHO LIVER FUNCTION TESTS

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
BILIRUBIN (TOTAL), Serum	0.51	0.2-1.1 mg/dl	Vanadate oxidation
BILIRUBIN (DIRECT), Serum	0.17	0-0.3 mg/dl	Vanadate oxidation
BILIRUBIN (INDIRECT), Serum	0.34	<1.2 mg/dl	Calculated
TOTAL PROTEINS, Serum	6.7	5.7-8.2 g/dL	Biuret
ALBUMIN, Serum	4.2	3.2-4.8 g/dL	BCG
GLOBULIN, Serum	2.5	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	1.7	1 - 2	Calculated
SGOT (AST), Serum	15.2	<34 U/L	Modified IFCC
SGPT (ALT), Serum	18.2	10-49 U/L	Modified IFCC
GAMMA GT, Serum	27.2	<73 U/L	Modified IFCC
ALKALINE PHOSPHATASE, Serum	47.0	46-116 U/L	Modified IFCC

<sup>\*</sup>Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
\*\*\* End Of Report \*\*\*







Page 13 of 13