

Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : -

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

: 22-Mar-2023 / 09:56 : 22-Mar-2023 / 16:06 E

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

Collected

Reported

CBC (Complete Blood Count), Blood			
<u>PARAMETER</u>	RESULTS	BIOLOGICAL REF RANGE	<u>METHOD</u>
RBC PARAMETERS			
Haemoglobin	13.5	13.0-17.0 g/dL	Spectrophotometric
RBC	4.09	4.5-5.5 mil/cmm	Elect. Impedance
PCV	44.6	40-50 %	Measured
MCV	109	80-100 fl	Calculated
MCH	33.1	27-32 pg	Calculated
MCHC	30.4	31.5-34.5 g/dL	Calculated
RDW	14.9	11.6-14.0 %	Calculated
WBC PARAMETERS			
WBC Total Count	5450	4000-10000 /cmm	Elect. Impedance
WBC DIFFERENTIAL AND ABSO	WBC DIFFERENTIAL AND ABSOLUTE COUNTS		
Lymphocytes	26.6	20-40 %	
Absolute Lymphocytes	1449.7	1000-3000 /cmm	Calculated
Monocytes	6.5	2-10 %	
Absolute Monocytes	354.3	200-1000 /cmm	Calculated
Neutrophils	62.9	40-80 %	
Absolute Neutrophils	3428.1	2000-7000 /cmm	Calculated
Eosinophils	3.7	1-6 %	
Absolute Eosinophils	201.7	20-500 /cmm	Calculated
Basophils	0.3	0.1-2 %	

WBC Differential Count by Absorbance & Impedance method/Microscopy.

16.4

PLATELET PARAMETERS

Platelet Count	140000	150000-400000 /cmm	Elect. Impedance
MPV	13.6	6-11 fl	Calculated
PDW	33.2	11-18 %	Calculated

20-100 /cmm

RBC MORPHOLOGY

Absolute Basophils

Immature Leukocytes

Calculated



CID :2308109508

Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Collected Consulting Dr. :22-Mar-2023 / 09:56 Reported :22-Mar-2023 / 12:16 : Borivali West (Main Centre) Reg. Location

Hypochromia

Microcytosis

Macrocytosis

Anisocytosis

Poikilocytosis

Polychromasia

Target Cells

Basophilic Stippling

Others

WBC MORPHOLOGY

PLATELET MORPHOLOGY Megaplatelets seen on smear

COMMENT

Advice:1)Serum vitamin B12 and folic acid estemation.

2)Reticulocyte count.

Normoblasts

Specimen: EDTA Whole Blood

ESR, EDTA WB-ESR 2-15 mm at 1 hr. Sedimentation

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West *** End Of Report ***







Dr.KETAKI MHASKAR M.D. (PATH) **Pathologist**

BMhaskar

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Page 2 of 11



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : -

Reg. Location

: Borivali West (Main Centre)

Authenticity Check

E

Use a QR Code Scanner Application To Scan the Code

: 22-Mar-2023 / 09:56

Reported :22-Mar-2023 / 13:00

Collected

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
GLUCOSE (SUGAR) FASTING, Fluoride Plasma	99.5	Non-Diabetic: < 100 mg/dl Impaired Fasting Glucose: 100-125 mg/dl Diabetic: >/= 126 mg/dl	Hexokinase
GLUCOSE (SUGAR) PP, Fluoride Plasma PP/R	85.4	Non-Diabetic: < 140 mg/dl Impaired Glucose Tolerance: 140-199 mg/dl Diabetic: >/= 200 mg/dl	Hexokinase
BILIRUBIN (TOTAL), Serum	0.88	0.1-1.2 mg/dl	Colorimetric
BILIRUBIN (DIRECT), Serum	0.34	0-0.3 mg/dl	Diazo
BILIRUBIN (INDIRECT), Serum	0.54	0.1-1.0 mg/dl	Calculated
TOTAL PROTEINS, Serum	6.8	6.4-8.3 g/dL	Biuret
ALBUMIN, Serum	4.5	3.5-5.2 g/dL	BCG
GLOBULIN, Serum	2.3	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	2.0	1 - 2	Calculated
SGOT (AST), Serum	30.4	5-40 U/L	NADH (w/o P-5-P)
SGPT (ALT), Serum	42.2	5-45 U/L	NADH (w/o P-5-P)
GAMMA GT, Serum	18.8	3-60 U/L	Enzymatic
ALKALINE PHOSPHATASE, Serum	90.0	40-130 U/L	Colorimetric
BLOOD UREA, Serum	19.8	12.8-42.8 mg/dl	Kinetic
BUN, Serum	9.2	6-20 mg/dl	Calculated
CREATININE, Serum	0.72	0.67-1.17 mg/dl	Enzymatic



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. :

Reg. Location : Borivali West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Collected : 22-Mar-2023 / 12:49

Reported :22-Mar-2023 / 18:27

eGFR, Serum 129 >60 ml/min/1.73sqm Calculated

URIC ACID, Serum 8.5 3.5-7.2 mg/dl Enzymatic

Urine Sugar (Fasting) Absent Absent
Urine Ketones (Fasting) Absent Absent

Urine Sugar (PP) Absent Absent Urine Ketones (PP) Absent Absent

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West
*** End Of Report ***







Dr.KETAKI MHASKAR M.D. (PATH) Pathologist

Page 4 of 11



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : -

Reg. Location: Borivali West (Main Centre)



Authenticity Check

Use a QR Code Scanner Application To Scan the Code

: 22-Mar-2023 / 09:56

:22-Mar-2023 / 13:00

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE GLYCOSYLATED HEMOGLOBIN (HbA1c)

Collected

Reported

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Glycosylated Hemoglobin (HbA1c), EDTA WB - CC	5.0	Non-Diabetic Level: < 5.7 % Prediabetic Level: 5.7-6.4 % Diabetic Level: >/= 6.5 %	HPLC
Estimated Average Glucose (eAG), EDTA WB - CC	96.8	mg/dl	Calculated

Intended use:

- In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year
- · In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly
- For microvascular disease prevention, the HbA1C goal for non pregnant adults in general is Less than 7%.

Clinical Significance:

- HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.
- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

Factors affecting HbA1c results:

Increased in: High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, Splenectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.

Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies

Reflex tests: Blood glucose levels, CGM (Continuous Glucose monitoring)

References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West
*** End Of Report ***



Dr.KETAKI MHASKAR M.D. (PATH) Pathologist

Page 5 of 11



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : - Collected : 22-Mar-2023 / 09:56
Reg. Location : Borivali West (Main Centre) Reported : 22-Mar-2023 / 16:53



Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Application To Scan the Code
• 22-Mar-2023 / 00-5

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE URINE EXAMINATION REPORT

<u>PARAMETER</u>	RESULTS	BIOLOGICAL REF RANGE	<u>METHOD</u>	
PHYSICAL EXAMINATION				
Color	Pale yellow	Pale Yellow	-	
Reaction (pH)	5.0	4.5 - 8.0	Chemical Indicator	
Specific Gravity	1.010	1.001-1.030	Chemical Indicator	
Transparency	Clear	Clear	-	
Volume (ml)	50	-	-	
CHEMICAL EXAMINATION				
Proteins	Absent	Absent	pH Indicator	
Glucose	Absent	Absent	GOD-POD	
Ketones	Absent	Absent	Legals Test	
Blood	Absent	Absent	Peroxidase	
Bilirubin	Absent	Absent	Diazonium Salt	
Urobilinogen	Normal	Normal	Diazonium Salt	
Nitrite	Absent	Absent	Griess Test	
MICROSCOPIC EXAMINATION				
Leukocytes(Pus cells)/hpf	1-2	0-5/hpf		
Red Blood Cells / hpf	Absent	0-2/hpf		
Epithelial Cells / hpf	0-1			
Casts	Absent	Absent		
Crystals	Absent	Absent		
Amorphous debris	Absent	Absent		
Bacteria / hpf	3-4	Less than 20/hpf		

Interpretation: The concentration values of Chemical analytes corresponding to the grading given in the report are as follows:

- Protein: (1+ ~25 mg/dl, 2+ ~75 mg/dl, 3+ ~ 150 mg/dl, 4+ ~ 500 mg/dl)
- Glucose:(1+ ~ 50 mg/dl, 2+ ~100 mg/dl, 3+ ~300 mg/dl,4+ ~1000 mg/dl)
- Ketone:(1+ ~5 mg/dl, 2+ ~15 mg/dl, 3+ ~ 50 mg/dl, 4+ ~ 150 mg/dl)

Reference: Pack insert

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West





Others



Dr.KETAKI MHASKAR M.D. (PATH) Pathologist

Page 6 of 11



CID : 2308109508

Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. **Collected** Reg. Location : Borivali West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Reported

*** End Of Report ***



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : Reg. Location : Borivali West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

: 22-Mar-2023 / 09:56 : 22-Mar-2023 / 14:42

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE BLOOD GROUPING & Rh TYPING

Collected

Reported

<u>PARAMETER</u> <u>RESULTS</u>

ABO GROUP B

Rh TYPING Positive

NOTE: Test performed by automated Erythrocytes magnetized technology (EMT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

Clinical significance:

ABO system is most important of all blood group in transfusion medicine

Limitations:

- ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- · Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

Refernces:

- 1. Denise M Harmening, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia
- 2. AABB technical manual

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD SDRL, Vidyavihar Lab
*** End Of Report ***







Dr.TRUPTI SHETTY M. D. (PATH) Pathologist

Page 8 of 11



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. :

Reg. Location : Borivali West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

: 22-Mar-2023 / 09:56 : 22-Mar-2023 / 13:00

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE LIPID PROFILE

Collected

Reported

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
CHOLESTEROL, Serum	159.4	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >/=240 mg/dl	CHOD-POD
TRIGLYCERIDES, Serum	163.4	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high:>/=500 mg/dl	GPO-POD
HDL CHOLESTEROL, Serum	30.3	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Homogeneous enzymatic colorimetric assay
NON HDL CHOLESTEROL, Serum	129.1	Desirable: <130 mg/dl Borderline-high:130 - 159 mg/d High:160 - 189 mg/dl Very high: >/=190 mg/dl	Calculated l
LDL CHOLESTEROL, Serum	96.0	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >/= 190 mg/dl	Calculated
VLDL CHOLESTEROL, Serum	33.1	< /= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO, Serum	5.3	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO, Serum	3.2	0-3.5 Ratio	Calculated

^{*}Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West
*** End Of Report ***







Dr.KETAKI MHASKAR M.D. (PATH) Pathologist

Page 9 of 11



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : -

Reg. Location : Borivali West (Main Centre)



R

E

Use a QR Code Scanner Application To Scan the Code

Collected : 22-Mar-2023 / 09:56

Reported :22-Mar-2023 / 14:23

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE THYROID FUNCTION TESTS

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
Free T3, Serum	4.7	3.5-6.5 pmol/L	ECLIA
Free T4, Serum	12.1	11.5-22.7 pmol/L	ECLIA
sensitiveTSH, Serum	14.22	0.35-5.5 microIU/ml	ECLIA



Name : MR.PUNEET KUMAR VERMA

Age / Gender : 40 Years / Male

Consulting Dr. : - Collected :22-Mar-2023 / 09:56

Reg. Location : Borivali West (Main Centre) Reported :22-Mar-2023 / 14:23

Interpretation

A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

Clinical Significance:

- 1)TSH Values between high abnormal upto15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological factors
 - can give falsely high TSH.
- 2)TSH values may be trasiently altered becuase of non thyroidal illness like severe infections, liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3 / T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone, Recovery phase of non-thyroidal illness, TSH Resistance.
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radio iodine Rx, post thyroidectomy, Anti thyroid drugs, tyrosine kinase inhibitors & amiodarone, amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hyperthyroidism, Graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum, hydatiform mole)
Low	Normal	Normal	Subclinical Hyperthyroidism, recent Rx for Hyperthyroidism, drugs like steroids & dopamine), Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal Illness, Recent Rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, Drug interference: Amiodarone, Heparin, Beta Blockers, steroids & anti epileptics.

Diurnal Variation:TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am, and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation:19.7%(with in subject variation)

Reflex Tests: Anti thyroid Antibodies, USG Thyroid , TSH receptor Antibody. Thyroglobulin, Calcitonin

Limitations:

- 1. Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.
- 2. Patient samples may contain heterophilic antibodies that could react in immunoassays to give falsely elevated or depressed results. this assay is designed to minimize interference from heterophilic antibodies.

Reference:

- 1.O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
- 2.Interpretation of the thyroid function tests, Dayan et al. THE LANCET . Vol 357
- 3. Tietz , Text Book of Clinical Chemistry and Molecular Biology -5th Edition
- 4.Biological Variation:From principles to Practice-Callum G Fraser (AACC Press)

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD Borivali Lab, Borivali West
*** End Of Report ***







Dr.KETAKI MHASKAR M.D. (PATH) Pathologist

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code

Page 11 of 11