

# **CERTIFICATE OF MEDICAL FITNESS**

NAME: M. R. praveen	Koney
AGE/GENDER: 32ys m	
HEIGHT: 176 cm	WEIGHT: 102/cgs
IDENTIFICATION MARK:	
BLOOD PRESSURE: 170 100 M	utter
PULSE: 170 100 m	
CVS: } RS:P ] Mornal.	
ANY OTHER DISEASE DIAGNOSED IN THE PAST:	Nicl
ALLERGIES, IF ANY:	Mil.
LIST OF PRESCRIBED MEDICINES:	Ni 4
ANY OTHER REMARKS:	
of Ms_Something who has signed disease and is fit for employment	R. pravemkuma\$on/daughter ed in my presence. He/ she has no physical
Signature of candidate	Dr. BINDURAJ. R HABBS, MD Internal Vindicine Reg. N St. 206 Signature of Medical Officer
Place: Spentrom Diagnostice & 1 Date: 30/10/24	
- · · · · · · · · · · · · · · · · · · ·	

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





Dr. Ashok S Bsc., MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827

DATE: 3010.24.

		EYE	EXAMINATION	
NAME: MS.	R.	Praven	Know AGE: 32 yr	GENDER: F/M

RIGHT EYE 616'm Vision Vision With glass Color Vision Normal Normal Anterior segment examination Normal Normal **Fundus Examination** Normal Normal Any other abnormality Nill Nill Diagnosis/ impression Normal Normal Dr. ASHOK SARODHE B.Sc., M.B.B.S., D.O.M.S.

Eye Consultant & Surgeon MC 31827

Consultant (Opthalmologist)







NAME	AGE	GENDER
1. Broven kemmer	3242	tolen

# **DENTAL EXAMINATION REPORT:**

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY & None.

M: MISSING ~ None.

O: OTHERS

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

**REMARKS:** 


SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA) Reg. No: 2247/A





0.15-35H	a VF	aVI	avp.			ID: 0008  PRAVEEN KIMAR  Male 37Vers
z AC50 25mm	2 ;		}	7		
0.15~35Hz AC50 25mm/s 10mm/mV 2*5.0s \(\psi_67\)	>		\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \		RS F.QTc QRS/T V5/SV1	30-10-2024 HR
				<u>}</u>	744	4 10:02:42 For 67 bpm 112 ms
V2.2 SEMIP V1.81	- V6	NV5		} ~33 }	Prolong Low T Report C	BP
				}	Prolonged P-wave Low T Wave(V6)  port Confirmed by:	Diagnosis Information: Sinus Rhythm
SPECTRUM DIAGNOSTICS & HEALTH CARE					A COUNTRIES OF THE PARTY OF THE	





Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC **Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM **Result Date** : 30-Oct-2024 03:47 PM

Report Status : Final

**Test Name** Result Unit Reference Value Method

3010240008

UHID

: 3010240008

### CHEST PA VIEW

- · Visualised lungs are clear.
- Bilateral hila appears normal.
- · Cardia is normal in size.
- · No pleural effusion.

**IMPRESSION**: No significant abnormality.



Printed By : spectrum

Printed On : 30 Oct, 2024 04:00 pm

DR PRAVEEN B, MBBS, DMRD, DNB Consultant Radiologist

Page 1 of 1






## **SPECTRUM DIAGNOSTICS**

Bangalore

Patient ID: 0739

Name: R PRAVEEN KUMAR CR Number: 20241030110356 Registration Date: 30-Oct-2024 Age: 32

Gender : Male Operator : spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	50	50	45	45	45	40	35	35	35	25	10
O - Air Right	55	55	55	50	50	45	40	30	30	30	20
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	37.73 dB	26.25 dB	40.00 dB	47.50 dB
AIR Right	41.82 dB	27.50 dB	45.00 dB	53.75 dB

#### Clinical Notes:

**Not Found** 





NAME AND LAB NO	MR PRAVEEN KUMAR	REG-008
AGE & SEX	32 YRS	MALE
DATE AND AREA OF INTEREST	30.10.2024	
REF BY	C/O APOLO CLINIC	

#### **USG ABDOMEN AND PELVIS**

LIVER:

Normal in size with increased echogenicity

No e/o IHBR dilatation. No evidence of focal lesion. Portal vein appears normal. CBD appears normal.

**GALL BLADDER:** 

Partially distended .No obvious calculus in the visualised luminal portion.

SPLEEN:

Normal in size and echotexture. No e/o focal lesion.

PANCREAS:

Head and body appears normal . Tail obscured by bowel gas shadows .

RETROPERITONEUM:

Suboptimal visualised due to bowel gas

RIGHT KIDNEY:

Right kidney is normal in size & echotexture.

Non obstructive calculus measuring 5.6 mm in the lower pole.

No evidence of hydronephrosis.

No solid lesions.

LEFT KIDNEY:

Left kidney is normal in size & echotexture. No evidence of calculus/ hydronephrosis.

No solid lesions.

**URINARY BLADDER:** 

Well distended. Mild diffuse bladder wall thickening measuring 4.9 mm.

PROSTATE:

Normal in size and echotexture.

No evidence of ascites.

#### IMPRESSION:

- Grade I fatty liver .
- Right non obstructive renal calculus .No hydronephrosis .
- Mild diffuse urinary bladder wall thickening suggested urine analysis correlation to rule out cystitis.

Suggested clinical correlation

DR PRAVEEN B , DMRD , DNB
CONSULTANT RADIOLOGIST





Name

: MR. R PRAVEEN KUMAR

Age / Gender

: 32 years / Male

Ref. By Dr. Reg. No.

: C/O APOLO CLINIC

C/o

: 3010240008

: APOLLO CLINIC

: 3010240008

3010240008

**Bill Date** 

: 30-Oct-2024 08:20 AM Sample Col. Date: 30-Oct-2024 08:20 AM

**Result Date** 

: 30-Oct-2024 01:16 PM

Report Status

: Final

**Test Name** 

Result

Unit

**UHID** 

Reference Value

Method

#### 2D ECHO

## 2D ECHO CARDIOGRAHIC STUDY M-MODE

Cardiograhic Study		Size
Aorta	30	mm
Left Atrium	38	mm
Right Ventricle	27	mm
Left ventricle (Diastole)	48	mm
Left ventricle(Systole)	29	mm
Ventricular Septum (Diastole)	15	mm
Ventricular septum (Systole)	14	mm
Posterior Wall (Diastole)	14	mm
Posterior Wall (Systole)	15	mm
Fractional Shortening	30	%
Ejection fraction	60	%

## DOPPLER /COLOUR FLOW

Mitral Valve Velocity	MVE- 0.68m/s	MVA - 0.5	4m/s	E/A-1.27
Tissue Doppler e' (Septal) 9cm/s		E/e'(Septal	) -7	
Velocity/ Gradient acro valve	ss the Pulmonic	0.83m/s	3mm	Hg
Max. Velocity / Gradiental valve	nt across the Aortic	1.05m/s	4mm	Hg
Velocity / Gradient acro	ss the Tricuspid valve	2.45 m/s	26mi	nHg

Page 1 of 2



Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

**Test Name** 

C/o : APOLLO CLINIC **Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM

Method

**Result Date** : 30-Oct-2024 01:16 PM

Report Status : Final

## **2DECHO Cardiographic Study**

SITUS SOLITUS, LEVOCARDIA

• SYSTEMIC VEINS: Normal drainage. IVC-2.0<50% collapse with inspiration.

UHID

Unit

: 3010240008

Reference Value

3010240008

PULMONARY VEINS: Normal drainage.

• RIGHT ATRIUM: Normal size, LEFT ATRIUM: Normal size.

• RIGHT VENTRICLE: Normal size & Adequate function.

Result

• LEFT VENTRICLE: Con. LVH; No RWMA; LV Systolic function adequate.

• IAS: INTACT; IVS: INTACT.

• MITRAL VALVE: No stenosis; No regurgitation

• TRICUSPID VALVE: No stenosis; Trivial regurgitation

• AORTIC VALVE: No stenosis; No regurgitation

• PULMONIC VALVE: No stenosis; No regurgitation

GREAT ARTERIES: Normally related.

· AORTA: Left aortic arch. No aortic dissection

• PULMONARY ARTERY: Confluent branch pulmonary arteries

· NO PDA.

· No pericardial effusion.

#### IMPRESSION:

- ADEQUATE LEFT VENTRICLE SYSTOLIC FUNCTION
- NO REGIONAL WALL MOTION ABNORMALITY
- ADEQUATE RIGHT VENTRICLE SYSTOLIC FUNCTION

NO PAH



Printed By

: Durga


Printed On


: 30 Oct, 2024 01:16 pm

Ms.Durga V., ECHO Technician

Page 2 of 2











Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC **Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	110.00	mg/dL	0.0-200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	88.00	mg/dL	0.0-150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	30.00	mg/dL	40.0-60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	80	mg/dL	0.0130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	62	mg/dL	0.0-100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	18	mg/dL	0.0-40	Calculated
Cholesterol/HDL Ratio-Serum	3.67	Ratio	0.0-5.0	Calculated

: 3010240008

3010240008

UHID

#### Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.

Fasting Blood Sugar (FBS)-Plasma

96

mg/dL

60.0-110.0

Hexo Kinase

Page 1 of 15













Age / Gender : 32 years / Male Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:15 AM

Report Status : Final

**Test Name** Result Unit Reference Value Method

UHID

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

3010240008

: 3010240008

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By


: spectrum

Printed On : 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 2 of 15









Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC Reg. No. : 3010240008

C/o : APOLLO CLINIC UHID : 3010240008

3010240008

**Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase 18.0 (GGT)-Serum	18.00	U/L	Male: 15.0-85.0	Other g-Glut-
			Female: 5.0-55.0	3-carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.




Printed By : spectrum

Printed On : 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 3 of 15









Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008 C/o : APOLLO CLINIC Bill Date : 30-Oct-2024 08:20 AM : 3010240008 Sample Col. Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TF7 Serum	Γ)-			
Tri-Iodo Thyronine (T3)-Se	erum 1.39	ng/mL	0.60-1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.9	μg/dL	5.50-12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormo (TSH)-Serum	one 3.74	μIU/mL	0.35-5.50	Chemiluminescence Immunoassay (CLIA)

UHID

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15 Years: 0.82-2.13,Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years: 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.

Printed By : spectrum

Printed On : 30 Oct, 2024 04:01 pm

Milliam.

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 4 of 15



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010











Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC

Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Bl	ood EDTA			
Haemoglobin (HB)	18.50	g/dL	Male: 14.0 - 17.0	Spectrophotmeter
Red Blood Cell (RBC)	5.80		m3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	53.50	%	Male: 42.0 - 51.0	Electronic Pulse
Mean corpuscular volume (MCV)	92.10	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	31.90	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	34.60	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	43.40	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	15.00	%	Male: 11.80 - 14.50	Volumetric Impedance
Mean Platelet Volume (MPV)	10.90	fL	8.0-15.0	Volumetric Impedance
Platelet	1.64	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	11.80	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	5470.00	cells/cumm	Male: 4000.0 - 11000.0	Volumetric Impedance
Neutrophils	54.90	%	40.0-75.0	Light scattering/Manual
Lymphocytes	35.40	%	20.0-45.0	Light scattering/Manual
Eosinophils	2.80	%	0.0-8.0	Light scattering/Manual
Monocytes	6.90	%	0.0-10.0	Light scattering/Manual
Basophils	0.00	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	3.01	10^3/uL	2.0-7.0	Calculated

UHID

: 3010240008

3010240008

Page 5 of 15







Age / Gender : 32 years / Male Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC

Bill Date : 30-Oct-2024 08:20 AM Sample Col. Date: 30-Oct-2024 08:20 AM

**Result Date** : 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Absolute Lymphocyte Count	1.94	10^3/uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.37	10^3/uL	0.20-1.00	Calculated
Absolute Eosinophil Count	150.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10^3/uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	16	mm/hr	Male: 0.0 - 10.0	Westergren

3010240008

: 3010240008

UHID

## Peripheral Smear Examination-Whole Blood EDTA

Method : (Microscopy-Manual)

RBC'S : Normocytic Normochromic.

WBC'S : Are normal in total number, morphology and distribution.

Platelets : Adequate in number and normal in morphology.

No abnormal cells or hemoparasites are present.

Normocytic Normochromic Blood Picture. Impression:



Printed By : spectrum

Printed On : 30 Oct, 2024 04:01 pm

Page 6 of 15













Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM **Result Date** : 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serur	n			
Bilirubin Total-Serum	0.62	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.16	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.46	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	23.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	25.00	U/L	Male:16.0-63.0 Female:14.0-59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)- Serum	129.00	U/L	Adult: 45.0-117.0 Children: 48.0-445.0 Infants: 81.90-350.30	PNPP,AMP- Buffer
Protein, Total-Serum	7.69	g/dL	6.40-8.20	Biuret/Endpoint- With Blank
Albumin-Serum	4.60	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	3.09	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serun	n 1.49	Ratio	0.80-2.0	Calculated

UHID

: 3010240008

3010240008



Printed By

: spectrum

Printed On

: 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 7 of 15











Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC Reg. No.

: 3010240008

C/o : APOLLO CLINIC Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date

: 30-Oct-2024 11:15 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method	
Glycosylated Haemoglobin					
(HbA1c)-Whole Blood EDTA					
Glycosylated Haemoglobin	5.40	%	Non diabetic adults :<5.7	HPLC	
(HbA1c)			At risk (Prediabetes): 5.7 - 6.4		
(			Diagnosing Diabetes :>= 6.5		
			Diabetes		
			Excellent Control: 6-7		
			Fair to good Control: 7-8		
			Unsatisfactory Control:8-10		
			Poor Control :>10		
Estimated Average Glucose(eAG)	108.28	mg/dL		Calculated	

3010240008

: 3010240008

UHID

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

Printed On

+91 77604 97644 | 080 2337 1555

: 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 8 of 15









: MR. R PRAVEEN KUMAR Name

Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:24 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Kidney Function Test (KFT)-B	UN,CREA,Uri	ic Acid,Na,K,C	I-Serum	
Kidney Function Test (KFT)- Serum			. V	
Blood Urea Nitrogen (BUN)	10.5	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	1.19	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	5.75	mg/dL	Male: 3.50-7.20 Female: 2.60-6.0	
Electrolytes				
Sodium (Na+)-Serum	138.1	mmol/L	135.0-145.0	ISE-Direct
Potassium (K+)-Serum	3.95	mmol/L	3.50-5.50	ISE-Direct
Chloride (Cl-)-Serum	99.70	mmol/L	96.0-108.0	ISE-Direct

: 3010240008

3010240008

UHID

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By : spectrum

Printed On : 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 9 of 15













Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

**Reg. No.** : 3010240008

C/o : APOLLO CLINIC

Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:24 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)

3010240008

: 3010240008

UHID



Printed By : spectrum

Printed On : 30 Oct, 2024 04:01 pm

Milliam.

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 10 of 15







Age / Gender : 32 years / Male Ref. By Dr.

: C/O APOLO CLINIC Reg. No. : 3010240008

C/o

: APOLLO CLINIC

**Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 11:44 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination	on-Urine			
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	6.5		5.0-7.5	Dipstick
Specific Gravity	1.020		1.000-1.030	Dipstick
<b>Biochemical Examinatio</b>	n			Diponen
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination	n		•	
Pus Cells	3-4	hpf	0.0-5.0	Microscopy
Epithelial Cells		hpf	0.0-10.0	Microscopy
RBCs		hpf	Absent	Microscopy
Casts	Absent	•	Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

UHID

: 3010240008

3010240008

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum

Printed On : 30 Oct, 2024 04:01 pm Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 11 of 15







Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008 C/o

: APOLLO CLINIC

**Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 01:41 PM

Report Status : Final

**Test Name** Result Unit Reference Value Method Blood Group & Rh Typing-Whole Blood EDTA **Blood Group** Slide/Tube agglutination Rh Type Positive Slide/Tube agglutination

: 3010240008

UHID

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum

: 30 Oct, 2024 04:01 pm Printed On

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 12 of 15



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 info@spectrumdiagnostics.org www.spectrumdiagnostics.org +91 77604 97644 | 080 2337 1555





Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC **Bill Date** 

: 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date

Report Status

: 30-Oct-2024 01:41 PM : Final

Test Name	Result	Unit	Reference Value	Method
Post Prandial Urine Sugar	Negative		Negative	Dipstick/Benedicts(Manual

3010240008

: 3010240008

UHID



Printed By

: spectrum

Printed On

: 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 13 of 15



SCAN FOR LOCATION





Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC Bill Date : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM Result Date : 30-Oct-2024 01:41 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	102	mg/dL	70-140	Hexo Kinase

3010240008

: 3010240008

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you cat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

UHID

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By

: spectrum

Printed On

: 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 14 of 15







Age / Gender : 32 years / Male

Ref. By Dr. : C/O APOLO CLINIC

Reg. No. : 3010240008

C/o : APOLLO CLINIC **Bill Date** : 30-Oct-2024 08:20 AM

Sample Col. Date: 30-Oct-2024 08:20 AM **Result Date** : 30-Oct-2024 03:35 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Calcium, Total- Serum	9.00	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein complexone)

3010240008

: 3010240008

UHID



Printed By

: spectrum

Printed On

: 30 Oct, 2024 04:01 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 15 of 15



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 9 +91 77604 97644 | 080 2337 1555



info@spectrumdiagnostics.org

