Ref. Dr : MediWheel Type : OP

<u>Investigation</u>	Observed Value	<u>Unit</u>	Biological Reference Interval
<b>IMMUNOHAEMATOLOGY</b>			
<b>BLOOD GROUPING AND Rh TYPING</b> (Blood /Agglutination)	'O' 'Positive'		
<b>HAEMATOLOGY</b>			
Complete Blood Count With - ESR			
Haemoglobin (Blood/Spectrophotometry)	09.66	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (Blood/Derived from Impedance)	31.37	%	37 - 47
RBC Count (Blood/Impedance Variation)	04.26	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (Blood/ Derived from Impedance)	73.69	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (Blood/Derived from Impedance)	22.68	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (Blood/Derived from Impedance)	30.78	g/dL	32 - 36
RDW-CV(Derived from Impedance)	15.5	%	11.5 - 16.0
RDW-SD(Derived from Impedance)	39.98	fL	39 - 46
Total Leukocyte Count (TC) (Blood/ Impedance Variation)	8130	cells/cu.mm	4000 - 11000
<b>Neutrophils</b> (Blood/Impedance Variation & Flow Cytometry)	57.30	%	40 - 75
<b>Lymphocytes</b> (Blood/Impedance Variation & Flow Cytometry)	35.50	%	20 - 45
<b>Eosinophils</b> (Blood/Impedance Variation & Flow Cytometry)	03.20	%	01 - 06
<b>Monocytes</b> (Blood/Impedance Variation & Flow Cytometry)	03.70	%	01 - 10
<b>Basophils</b> (Blood/Impedance Variation & Flow Cytometry)	00.30	%	00 - 02
<b>INTERPRETATION:</b> Tests done on Automated microscopically.	Five Part cell counter. A	ll abnormal resu	ults are reviewed and confirmed
Absolute Neutrophil count (Blood/ Impedance Variation & Flow Cytometry)	4.66	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (Blood/ Impedance Variation & Flow Cytometry)	2.89	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (Blood/ Impedance Variation & Flow Cytometry)	0.26	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (Blood/ Impedance Variation & Flow Cytometry)	0.30	10^3 / µl	< 1.0







Ref. Dr : MediWheel Type : OP

<u>Investigation</u>	Observed Value	<u>Unit</u>	<b>Biological Reference Interval</b>
<b>Absolute Basophil count</b> (Blood/Impedance Variation & Flow Cytometry)	0.02	10^3 / µl	< 0.2
Platelet Count (Blood/Impedance Variation)	134	10^3 / µl	150 - 450
MPV (Blood/Derived from Impedance)	09.09	fL	8.0 - 13.3
PCT(Automated Blood cell Counter)	0.12	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated ESR analyser)	56	mm/hr	< 20
<b>BIOCHEMISTRY</b>			
BUN / Creatinine Ratio	15.3		
<b>Glucose Fasting (FBS)</b> (Plasma - F/GOD-PAP)	81.8	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/	83.2	mg/dL	70 - 140

#### INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Negative		Negative
9.5	mg/dL	7.0 - 21
0.62	mg/dL	0.6 - 1.1
3.0	mg/dL	2.6 - 6.0
0.46	mg/dL	0.1 - 1.2
0.12	mg/dL	0.0 - 0.3
0.34	mg/dL	0.1 - 1.0
21.5	U/L	5 - 40
20.6	U/L	5 - 41
15.9	U/L	< 38
	9.5 0.62 3.0 0.46 0.12 0.34 21.5 20.6	9.5 mg/dL  0.62 mg/dL  3.0 mg/dL  0.46 mg/dL  0.12 mg/dL  0.34 mg/dL  21.5 U/L







Ref. Dr : MediWheel Type : OP

<u>Investigation</u>	Observed Value	<u>Unit</u>	Biological Reference Interval
Alkaline Phosphatase (SAP) (Serum/ Modified IFCC)	73.2	U/L	42 - 98
Total Protein (Serum/Biuret)	7.15	gm/dL	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.85	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	3.30	gm/dL	2.3 - 3.6
A: GRATIO (Serum/Derived)	1.17		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	173.2	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	80.8	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the husual+dicirculating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	40.1	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	116.9	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	16.2	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	133.1	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.







Ref. Dr : MediWheel Type : OP

Investigation Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	Observed Value 4.3	<u>Unit</u>	Biological Reference Interval Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	2		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/ Calculated)	2.9		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)			
<b>HbA1C</b> (Whole Blood/Ion exchange HPLC by D10)	5.3	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose (Whole Blood) 105.41 mg/dL

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia,hyperbilirubinemia,Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

### **IMMUNOASSAY**

#### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total (Serum/ 1.24 ng/mL 0.7 - 2.04

Chemiluminescent Immunometric Assay (CLIA))

#### INTERPRETATION:

#### Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

**T4 (Tyroxine) - Total** (Serum/ 10.81 μg/dL 4.2 - 12.0

Chemiluminescent Immunometric Assay

(CLIA))

### INTERPRETATION:

#### Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.







Ref. Dr : MediWheel Type : OP

<u>Investigation</u> <u>Observed Value</u> <u>Unit</u> <u>Biological Reference Interval</u>

TSH (Thyroid Stimulating Hormone) (Serum 1.57 µIU/mL 0.35 - 5.50

/Chemiluminescent Immunometric Assay

(CLIA))

#### **INTERPRETATION:**

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and

BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations. 3. Values&amplt;0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

### **CLINICAL PATHOLOGY**

#### **Urine Analysis - Routine**

Colour (Urine) Appearance (Urine)	Pale Yellow Clear		Yellow to Amber Clear
Protein (Urine)	Negative		Negative
Glucose (Urine)	Negative		Negative
Pus Cells (Urine)	1-2	/hpf	NIL
Epithelial Cells (Urine)	3-4	/hpf	NIL
RBCs (Urine)	Nil	/hpf	NIL

-- End of Report --







Name	MRS.SILUVAI ANTENEETA SONIYA D	ID	MED121758925
Age & Gender	37Y/FEMALE	Visit Date	22 Mar 2023
Ref Doctor Name	MediWheel		

### Thanks for your reference

# ECHOCARDIOGRAM WITH COLOUR DOPPLER:

LVID d	5.0 cm
LVID s	2.9 cm
EF	73 %
IVS d	1.1 cm
IVS s	1.3 cm
LVPW d	0.8 cm
LVPW s	1.1 cm
LA	3.0 cm
AO	3.2 cm
TAPSE	20mm
IVC	1.2cm

Left ventricle, Left atrium normal.

Right ventricle, Right atrium normal.

No regional wall motion abnormality present.

Mitral valve, Aortic valve, Tricuspid valve & Pulmonary valve normal.

Aorta normal.

Inter atrial septum intact.

Inter ventricular septum intact.

No pericardial effusion.

# Doppler:

Name	MRS.SILUVAI ANTENEETA SONIYA D	ID	MED121758925
Age & Gender	37Y/FEMALE	Visit Date	22 Mar 2023
Ref Doctor Name	MediWheel		

Mitral valve: E: 0.62 m/s A:1.10 m/s

E/A Ratio: 0.57 E/E:8.23

Aortic valve: AV Jet velocity:1.71 m/s

Tricuspid valve: TV Jet velocity: 2.35 m/s TRPG:22.16mmHg.

Pulmonary valve: PV Jet velocity: 1.61 m/s

# **IMPRESSION:**

1. Normal chambers & Valves.

- 2. No regional wall motion abnormality present.
- 3. Normal LV systolic function.
- 4. Pericardial effusion Nil.

5. No pulmonary artery hypertension.

Dr. S. MANIKANDAN. MD.DM.(Cardio) Cardiologist

Name	MRS.SILUVAI ANTENEETA SONIYA D	ID	MED121758925
Age & Gender	37Y/FEMALE	Visit Date	22 Mar 2023
Ref Doctor Name	MediWheel		

Name	MRS.SILUVAI ANTENEETA SONIYA D	ID	MED121758925
Age & Gender	37Y/FEMALE	Visit Date	22 Mar 2023
Ref Doctor Name	MediWheel		

## 

Poor penetration of USG due to thick abdominal wall.

Liver: The liver is normal in size and shows uniform echotexture with

no focal abnormality. There is no intra or extra hepatic biliary

ductal dilatation.

Gallbladder: The gall bladder is contracted. A linear echogenic focus measuring

about 28 mm with posterior acoustic shadowing, noted within the

lumen.

Pancreas: The pancreas shows a normal configuration and echotexture.

The pancreatic duct is normal.

Spleen: The spleen is normal.

Kidneys: The right kidney measures 10.3 x 4.5 cm. Normal architecture.

The collecting system is not dilated.

The left kidney measures 11.1 x 5.5 cm. Normal architecture.

The collecting system is not dilated.

Urinary

bladder: The urinary bladder is smooth walled and uniformly transonic.

There is no intravesical mass or calculus.

Uterus: The uterus is anteverted, and measures 8.1 x 4.4 cm.

Myometrial echoes are homogeneous.

Name	MRS.SILUVAI ANTENEETA SONIYA D	ID	MED121758925
Age & Gender	37Y/FEMALE	Visit Date	22 Mar 2023
Ref Doctor Name	MediWheel		

The endometrium is central and normal measures 9.8 mm in thickness.

Ovaries: The right ovary measure 2.6 x 2.6 cm.

The left ovary measures 3.2 x 2.8 cm. Few small follicles noted in both ovaries.

No significant mass or cyst is seen in the ovaries.

Parametria are free.

RIF: Iliac fossae are normal.

No mass or fluid collection is seen in the right iliac fossa.

The appendix is not visualized.

There is no free or loculated peritoneal fluid. No para aortic lymphadenopathy is seen.

## **IMPRESSION**:

Suggestive of Cholelithiasis.-To be reviewed with 12 hrs fasting if necessary.

DR.T.ANNIE STALIN MBBS.,F.USG., SONOLOGIST.

Name	SILUVAI ANTENEETA SONIYA D	Customer ID	MED121758925
Age & Gender	37Y/F	Visit Date	Mar 22 2023 8:37AM
Ref Doctor	MediWheel		

## Thanks for your reference

## **DIGITAL X- RAY CHEST PA VIEW**

Trachea appears normal.

Cardiothoracic ratio is within normal limits.

Bilateral lung fields appear normal.

Both costophrenic angles appear normal.

Visualised bony structures appear normal.

Extra thoracic soft tissues shadow grossly appears normal.

## **IMPRESSION:**

• NO SIGNIFICANT ABNORMALITY DEMONSTRATED.

DR. DANIEL STANLEY PETER, M.D.R.D. Consultant Radiologist

Reg. No: 82342