

CERTIFICATE OF MEDICAL FITNESS

NAME: M. Amresh

AGE/ GENDER: 59 y / male

HEIGHT: 168 cm

WEIGHT: 69.4 kg

IDENTIFICATION MARK: _____

BLOOD PRESSURE: 100/70 mmHg

PULSE: 68 bpm

CVS: Normal

RS:P

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Diabetic - glycimip m,
2ni Ryzoledg 8 units every

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: nil

ANY OTHER REMARKS: no

I Certify that I have carefully examined Mr/Mrs. Amresh son/daughter of Mrs M. Sadanandiah, who has signed in my presence. He/ she has no physical disease and is fit for employment.


Signature of candidate


Dr. BINDURAJ. R
MBBS, MD
Internal Medicine
Reg. No. 62856
Signature of Medical Officer

Place: Spectrum Diagnostics & Health Care

Date: 30/07/24

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined



Dr. Ashok S
Bsc., MBBS., D.O.M.S
Consultant Ophthalmologist
KMC No: 31827

DATE: 30/07/24

EYE EXAMINATION

NAME: ms. m. Anandesh AGE: 59 yrs. GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	<u>6/6p: 10</u>	<u>6/12: 10</u>
Vision With glass	-----	-----
Color Vision	<u>Normal</u>	<u>Normal</u>
Anterior segment examination	<u>Normal</u>	<u>Normal</u>
Fundus Examination	<u>Normal</u>	<u>Normal</u>
Any other abnormality	<u>Nil</u>	<u>Nil</u>
Diagnosis/ impression	<u>Normal</u>	<u>Normal</u>

Dr. ASHOK SARODHE
B.Sc., M.B.B.S., D.O.M.S.
E. J. Consultant & Surgeon
KMC 31827
Consultant (Ophthalmologist)




0037
Mr Anarash
M 59y 000Kg
09:42 AM
30/07/2024

25 mm/s
10mm/mV
1.0 - 35Hz
50Hz ReJ-Y
BLC-Y

P = 92 ms
QRS = 100 ms
QT = 165 ms
QTc = 347 ms
QTc = 396 ms

QT/QTc = 87%
QT/RR = 45%
QRS axis = -36°
P axis = -15°
T axis = 00°
HR = 78bpm



SPECTRUM DIAGNOSTICS

Dr

Ref By Dr

ASSW CE

Name	: MR. M AMARESH	UHIP	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 01:07 PM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

XRAY CHEST GENERAL


CHEST PA VIEW

- Visualised lungs are clear.
- Bilateral hila appears normal.
- Cardia is normal in size.
- No pleural effusion.

IMPRESSION: No significant abnormality.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:44 pm



DR PRAVEEN B, MBBS, DMRD, DNB Consultant
Radiologist

Page 1 of 1

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010
+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

Other Branch: #486/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

Name : MR. M AMARESH	UHIP : 3007240007	Bill Date : 30-Jul-2024 08:09 AM
Age / Gender : 59 Years / Male		Sample Col. Date : 30-Jul-2024 08:09 AM
Ref. By Dr. : Dr. APOLO CLINIC	3007240007	Result Date : 30-Jul-2024 11:13 AM
Reg. No. : 3007240007		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blood EDTA				
Haemoglobin (HB)	14.90	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotometer
Red Blood Cell (RBC)	4.87	million/cumm	3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	44.10	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	90.50	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	30.60	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	33.80	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	40.60	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	14.50	%	Male: 11.80-14.50 Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	10.90	fL	8.0-15.0	Volumetric Impedance
Platelet	2.39	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	13.80	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	7820.00	cells/cumm	Male: 4000-11000 Female: 4000-11000 Children: 6000-17500 Infants : 9000-30000	Volumetric Impedance
Neutrophils	50.80	%	40.0-75.0	Light scattering/Manual
Lymphocytes	42.00	%	20.0-40.0	Light scattering/Manual
Eosinophils	2.70	%	0.0-8.0	Light scattering/Manual

Name	: MR. M AMARESH	UHIP	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:13 AM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Monocytes	4.50	%	0.0-10.0	Light scattering/Manual
Basophils	0.00	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	3.98	10 ³ /uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	3.28	10 ³ /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.35	10 ³ /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	210.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10 ³ /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	02	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

- RBC'S : Normocytic Normochromic.
- WBC'S : Are normal in total number, morphology and distribution.
- Platelets : Adequate in number and normal in morphology.
No abnormal cells or hemoparasites are present.
- Impression : Normocytic Normochromic Blood picture.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Name	: MR. M AMARESH	UHID	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 12:21 PM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	190.00	mg/dL	0.0-200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	111.00	mg/dL	0.0-150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	44.00	mg/dL	40.0-60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	146	mg/dL	0.0-130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	156.0	mg/dL	0.0-100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	22	mg/dL	0.0-40	Calculated
Cholesterol/HDL Ratio-Serum	4.32	Ratio	0.0-5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name : MR. M AMARESH	UHID : 3007240007	Bill Date : 30-Jul-2024 08:09 AM
Age / Gender : 59 Years / Male	 3007240007	Sample Col. Date : 30-Jul-2024 08:09 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 30-Jul-2024 12:21 PM
Reg. No. : 3007240007		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serum				
Bilirubin Total-Serum	0.72	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.14	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.58	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	22.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	26.00	U/L	Male:16.0-63.0 Female:14.0-59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	63.00	U/L	Adult: 45.0-117.0 Children: 48.0-445.0 Infants: 81.90-350.30	PNPP,AMP-Buffer
Protein, Total-Serum	6.62	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	4.01	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	2.61	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.54	Ratio	0.80-2.0	Calculated



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MR. M AMARESH	UHID	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 12:21 PM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test) :				
Blood Urea Nitrogen (BUN)-Serum	13.20	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	1.15	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	6.65	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	138.8	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	3.98	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	103.70	mmol/L	96.0-108.0	Ion-Selective Electrodes (ISE)

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MR. M AMARESH	UHID	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 12:21 PM
Reg. No.	: 3007240007		3007240007	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)- Serum				
Tri-Iodo Thyronine (T3)-Serum	1.19	ng/mL	0.60-1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.30	µg/dL	5.50-12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	3.53	µIU/mL	0.35-5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).

Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy

Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II- trimester: 0.2-3.0; III- trimester: 0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithin Reddy C, MD, Consultant Pathologist



Name : MR. M AMARESH	UHID : 3007240007	Bill Date : 30-Jul-2024 08:09 AM
Age / Gender : 59 Years / Male		Sample Col. Date : 30-Jul-2024 08:09 AM
Ref. By Dr. : Dr. APOLO CLINIC	3007240007	Result Date : 30-Jul-2024 12:21 PM
Reg. No. : 3007240007		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)- Plasma	135	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA

Glycosylated Haemoglobin (HbA1c)	9.10	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
---	------	---	---	------

Estimated Average Glucose(eAG)	214.47	mg/dL		Calculated
---------------------------------------	--------	-------	--	------------

Name	: MR. M AMARESH	UHIP	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 12:21 PM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 8 of 11

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org

SCAN FOR LOCATION



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 087 | 080-2891 6944 | 080-49511985

Name	: MR. M AMARESH	UHIP	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 12:21 PM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-Urine				
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.025		1.000-1.030	Dipstick
Biochemical Examination				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Positive (++++)		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				
Pus Cells	2-3	hpf	0.0-5.0	Microscopy
Epithelial Cells	1-2	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cell, may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithun Roddy C,MD,Consultant Pathologist

Page 9 of 11

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010

+91 77604 97644 | 080 2337 1555 Info@spectrumdiagnostics.org www.spectrumdiagnostics.org

SCAN FOR LOCATION



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6044 | 080-4981985

Name	: MR. M AMARESH	UHIP	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 12:46 PM
Reg. No.	: 3007240007			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	22.00	U/L	Male: 15.0-85.0 Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm

N. Nithun

Dr. Nithun Reddy C, MD, Consultant Pathologist



Name	: MR. M AMARESH	UHID	: 3007240007	Bill Date	: 30-Jul-2024 08:09 AM
Age / Gender	: 59 Years / Male			Sample Col. Date	: 30-Jul-2024 08:09 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 01:09 PM
Reg. No.	: 3007240007		3007240007	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	228	mg/dL	70-140	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:45 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



CERTIFICATE OF MEDICAL FITNESS

NAME: M. Sumathi

AGE/ GENDER: 53y / female

HEIGHT: 151 cm

WEIGHT: 71.3 kg

IDENTIFICATION MARK: -

BLOOD PRESSURE: 130/80 mmHg

PULSE: 84 bpm

CVS: normal
RS:P

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Diabetic - udapa 100/1000mg

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: Nil

ANY OTHER REMARKS: NO

I Certify that I have carefully examined Mr/Mrs. M. Sumathi son/daughter of Mr Seetharamiah who has signed in my presence. He/ she has no physical disease and is fit for employment.



Signature of candidate

Dr. BINDURAJ. R
MBBS, MD
Internal Medicine
Reg. No. 62306

Signature of Medical Officer

Place: Spectrum Diagnostics & Health Care

Date: 30/07/24

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined

SCAN FOR LOCATION



EYE EXAMINATION

NAME: *Ms. Smalvi* AGE: *53 yrs* GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	<i>6/12 = 2/10</i>	<i>6/12 = 2/10</i>
Vision With glass	<i>6/6 = 2/6</i>	<i>6/6 = 2/6</i>
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal

to wear spectacles
Dr. ASHOK SARODHE
 B.Sc., M.B.B.S., D.O.M.S.
 Eye Consultant & Surgeon
 KMC 31827




0037
Mrs sumathi
F 53y 000kg
09:47 AM
30/07/2024

AUTO SEQ
BLC-V

25 mm/s
10mm/mV
1.0 - 35Hz
50Hz Rej-Y

P	=	95 ms	QT/QTc	=	93x
QRS	=	87 ms	QT/RR	=	40x
PQ	=	157 ms	QRS ax15	=	35°
QT	=	360 ms	P ax15	=	-07°
QTc	=	383 ms	T ax15	=	-23°

To be clinically correlated: HR = 68bpm



SPECTRUM DIAGNOSTICS

Dr

Ref By Dr

ANW CE





SPECTRUM DIAGNOSTICS

Bangalore

Patient ID : 0533
Name : M SUMATHI
CR Number : 20240730115437
Registration Date : 30-Jul-2024

Age : 53
Gender : Female
Operator : spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	20	25	25	20	20	20	25	20	15	20	15
O - Air Right	25	30	35	35	30	30	25	25	20	25	20
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	20.45 dB	17.50 dB	21.67 dB	22.50 dB
AIR Right	27.27 dB	22.50 dB	28.33 dB	31.25 dB

Clinical Notes :

Not Found



NAME AND LAB NO	MRS SUMATHI M	REG -008
AGE & SEX	53 YRS	FEMALE
DATE AND AREA OF INTEREST	30.07.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

- LIVER:** Normal in size and shows diffuse increased echogenicity. No e/o IHBR dilatation. No evidence of focal lesion. Portal vein appears normal. CBD appears normal.
- GALL BLADDER:** Partially distended. No obvious calculus in the visualised luminal portion.
- SPLEEN:** Normal in size and echotexture. No focal lesion
- PANCREAS:** Head and body appears normal. Tail obscured by bowel gas shadows
- RETROPERITONEUM:** Suboptimal visualised due to bowel gas.
- RIGHT KIDNEY:** Right kidney is normal in size & echotexture. No evidence of calculus/ hydronephrosis.
- LEFT KIDNEY:** Left kidney is normal in size & echotexture. No evidence of calculus/ hydronephrosis.
- URINARY BLADDER:** Well distended. No wall thickening/ calculi.
- UTERUS: OVARIES:** Post menopausal status. Endometrium is normal. ET – 2.4 mm. No obvious adnexal mass lesions

- No evidence of ascites.

IMPRESSION:

- *Grade I fatty liver.*


DR PRAVEEN B, DMRD, DNB
CONSULTANT RADIOLOGIST

SCAN FOR LOCATION



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:14 AM
Reg. No.	: 3007240008		3007240008	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blood EDTA				
Haemoglobin (HB)	14.40	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotometer
Red Blood Cell (RBC)	4.73	million/cumm	3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	42.60	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	90.10	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	30.30	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	33.70	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	42.00	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	15.10	%	Male: 11.80-14.50 Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	9.30	fL	8.0-15.0	Volumetric Impedance
Platelet	3.17	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	9.30	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	5840.00	cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants : 9000-30000	Volumetric Impedance
Neutrophils	54.10	%	40.0-75.0	Light scattering/Manual
Lymphocytes	40.00	%	20.0-40.0	Light scattering/Manual
Eosinophils	2.80	%	0.0-8.0	Light scattering/Manual

SCAN FOR LOCATION



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:14 AM
Reg. No.	: 3007240008			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.00	%	0.0-10.0	Light scattering/Manual
Basophils	0.10	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	3.15	10 ³ /uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	2.33	10 ³ /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.18	10 ³ /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	170.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.01	10 ³ /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	18	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.
WBC'S : Are normal in total number, morphology and distribution.
Platelets : Adequate in number and normal in morphology.
No abnormal cells or hemoparasites are present.
Impression : Normocytic Normochromic Blood picture.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:28 AM
Reg. No.	: 3007240008			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test) :				
Blood Urea Nitrogen (BUN)-Serum	12.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.64	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	4.44	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	138.8	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.03	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	102.80	mmol/L	96.0-108.0	Ion-Selective Electrodes (ISE)

Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name : MRS. M SUMATHI	UHID : 3007240008	Bill Date : 30-Jul-2024 08:14 AM
Age / Gender : 53 years / Female	 3007240008	Sample Col. Date : 30-Jul-2024 08:14 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 30-Jul-2024 11:28 AM
Reg. No. : 3007240008		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	191.00	mg/dL	0.0-200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	98.00	mg/dL	0.0-150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	49.00	mg/dL	40.0-60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	142	mg/dL	0.0-130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	136.00	mg/dL	0.0-100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	20	mg/dL	0.0-40	Calculated
Cholesterol/HDL Ratio-Serum	3.90	Ratio	0.0-5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:28 AM
Reg. No.	: 3007240008			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)- Plasma	144	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol, Dietary - Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA

Glycosylated Haemoglobin (HbA1c)	7.70	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
Estimated Average Glucose (eAG)	174.28	mg/dL		Calculated



Name : MRS. M SUMATHI	UHID : 3007240008	Bill Date : 30-Jul-2024 08:14 AM
Age / Gender : 53 years / Female	 3007240008	Sample Col. Date : 30-Jul-2024 08:14 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 30-Jul-2024 11:28 AM
Reg. No. : 3007240008		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:28 AM
Reg. No.	: 3007240008			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serum				
Bilirubin Total-Serum	0.66	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.12	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.54	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	22.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	18.00	U/L	Male:16.0-63.0 Female:14.0-59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	77.00	U/L	Adult: 45.0-117.0 Children: 48.0-445.0 Infants: 81.90-350.30	PNPP,AMP-Buffer
Protein, Total-Serum	7.02	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	4.15	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	2.87	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.45	Ratio	0.80-2.0	Calculated



Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:28 AM
Reg. No.	: 3007240008			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	26	U/L	Male: 15.0-85.0 Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.

Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



Name	: MRS. M SUMATHI	UHID	: 3007240008	Bill Date	: 30-Jul-2024 08:14 AM
Age / Gender	: 53 years / Female			Sample Col. Date	: 30-Jul-2024 08:14 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 30-Jul-2024 11:29 AM
Reg. No.	: 3007240008			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)-Serum				
Tri-Iodo Thyronine (T3)-Serum	1.13	ng/mL	0.60-1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	11.90	µg/dL	5.50-12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	5.02	µIU/mL	0.35-5.50	Chemiluminescence Immunoassay (CLIA)

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10
 Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,
 Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).
Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.9-16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3
 1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.
Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, calorie intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0
 Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.
Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.



Printed By : spectrum
 Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



Name : MRS. M SUMATHI	UHID : 3007240008	Bill Date : 30-Jul-2024 08:14 AM
Age / Gender : 53 years / Female	 3007240008	Sample Col. Date : 30-Jul-2024 08:14 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 30-Jul-2024 12:22 PM
Reg. No. : 3007240008		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-Urine				
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Slightly Turbid		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.025		1.000-1.030	Dipstick
Biochemical Examination				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Positive (+++)		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				
Pus Cells	10-12	hpf	0.0-5.0	Microscopy
Epithelial Cells	7-9	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Bacteria Present		Absent	Microscopy

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.

Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



Name : MRS. M SUMATHI	Uhid : 3007240008	Bill Date : 30-Jul-2024 08:14 AM
Age / Gender : 53 years / Female	 3007240008	Sample Col. Date : 30-Jul-2024 08:14 AM
Ref. By Dr. : Dr. APOLO CLINIC		Result Date : 30-Jul-2024 01:36 PM
Reg. No. : 3007240008		Report Status : Final
C/o : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Blood Group & Rh Typing-Whole Blood EDTA				
Blood Group	A			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum
Printed On : 30 Jul, 2024 02:41 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

