

CERTIFICATE OF MEDICAL FITNESS

NAME: Robout Nayaka-M
AGE/GENDER: 464 / Male
HEIGHT: 171 Cm WEIGHT: 93 Cg
IDENTIFICATION MARK: Bluele mole on the chest
BLOOD PRESSURE: 140 /80 mmtg
PULSE: 68 b/m
CVS: RS:P Normal
ANY OTHER DISEASE DIAGNOSED IN THE PAST: The you'd (Theyenorm - 62 mg
ALLERGIES, IF ANY:
LIST OF PRESCRIBED MEDICINES: N°C
ANY OTHER REMARKS:
I Certify that I have carefully examined Mr/Mrs. Roll Mayalca son/daughter of Mr. Muralidha who has signed in my presence. He/ she has no physical disease and is fit for employment.
Dr. BINDURAJ. R
Signature of candidate Signature of Medical Officer
Place: Spectner Diagnostics & health call
Date: 24 2 24
The state of the s

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





Dr. Ashok S Bsc., MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827

DATE: 24-02-24

EYE EXAMINATION

NAME: MS.	Kohit	Wagaka WAGE:	GENDE	R:F/M
		DIGHT EVE	I FFT FY	'F

Vision Vision With glass Normal Normal **Color Vision** Normal Normal Anterior segment examination Normal Normal **Fundus Examination** Nill Nill Any other abnormality Normal Normal Diagnosis/ impression

> Consultant (Opthalmologist)M.S. sultant & Surgeon KMC 31827



0.15-35Hz AC50 25mm/s			\[\frac{1}{2} \]	ID: 2240040 MR ROHIT NAYAKA M Male 46Years
10mm/mV 2*5.0s •78			\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	24-02-2024 10:27:31 For BPL HR : 78 bpm P : 103 ms PR : 140 ms QRS : 94 ms QT/QTc : 362/414 ms P/QRS/T : 12/46/16 ° RV5/SV1 : 1.380/0.797 mV
V2.2 SEMIP VI.81 SPECTRUM DIAGNOSTICS & HEALTH CARE				Diagnosis Information: Sinus Rhythm ***Normal ECG*** Report Confirmed by:



NAME	: MR.ROHIT NAYAKA M	DATE : 24/02/2024
AGE/SEX	: 46YEARS/MALE	REG NO: 2402240040
REF BY	: APOLLO CLINIC	

CHEST PA VIEW

- Visualised lungs are clear.
- Bilateral hila appears normal.
- Cardia is normal in size
- No pleural effusion

IMPRESSION: No Significant Abnormality Detected

Transcorp

DR PRAVEEN B, DMRD, DNB **Consultant Radiologist**







: 46 years / Male Age / Gender

Ref. By Dr. : Dr. APOLO CLINIC

: 2402240040 Reg. No.

C/o : Apollo Clinic

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM : 24-Feb-2024 10:59 AM **Result Date**

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood ED			clinical symptoms.	
n hamen traiten	6.80	%	Non diabetic adults :<5.7	HPLC
Glycosylated Haemoglobin (HbA1c)			At risk (Prediabetes): 5.7 - 6.4	
			Diagnosing Diabetes :>= 6.5	
			Diabetes	
			Excellent Control: 6-7	
			Fair to good Control: 7-8	
			Chambras of Commercial	
			Poor Control :>10	
Estimated Average Glucose(eAG)	148.46	mg/dL		Calculated

: 2402240040

2402240040

UHID

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

: 24 Feb, 2024 06:17 pm Printed On

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page I of II















: 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240040

Age / Gender

: Apollo Clinic C/o

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM **Result Date** : 24-Feb-2024 11:53 AM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examinati	on-Urine		h clinical symptoms.	
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.015		1.000-1.030	Dipstick
Biochemical Examination	on 180 smos ni nzis h			
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				
Pus Cells	1-2	hpf	0.0-5.0	Microscopy
Epithelial Cells	1-2	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent	-	Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

: 2402240040

2402240040

UHID

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By

: spectrum

Printed On


: 24 Feb, 2024 06:17 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 2 of 11















Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240040

C/o : Apollo Clinic

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM **Result Date** : 24-Feb-2024 12:04 PM

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blo	ood EDTA		clinical symptoms.	conjunction with
Haemoglobin (HB)	13.40	g/dL	Male: 14.0-17.0 Female:12.0-15.0 Newborn:16.50 - 19.50	Spectrophotmeter
Red Blood Cell (RBC)	4.74	million/cumr	n3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	38.40	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	81.00	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	28.30	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	34.90	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	40.50	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	15.90	%	Male: 11.80-14.50 Female:12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	9.30	fL	8.0-15.0	Volumetric Impedance
Platelet	2.06	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	13.10	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	8090.00	cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants: 9000-30000	Volumetric Impedance
Neutrophils	59.60	%	40.0-75.0	Light scattering/Manual
Lymphocytes	30.20	%	20.0-40.0	Light scattering/Manual
Eosinophils	6.50	%	0.0-8.0	Light scattering/Manual

UHID

: 2402240040

2402240040









Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC : 2402240040 Reg. No.

: Apollo Clinic C/o

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM : 24-Feb-2024 12:04 PM **Result Date**

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.60	%	0.0-10.0	Light scattering/Manual
Basophils	0.10	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	4.82	10^3/uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	2.44	10^3/uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.29	10^3/uL	0.20-1.00	Calculated
Absolute Eosinophil Count	530.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.01	10^3/uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	25	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren

: 2402240040

2402240040

UHID

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

: Normocytic Normochromic. RBC'S

: Are normal in total number, morphology and distribution. WBC'S

: Adequate in number and normal in morphology. **Platelets**

No abnormal cells or hemoparasites are present.

Impression: Normocytic Normochromic Blood picture.



Printed By

: spectrum

Printed On : 24 Feb, 2024 06:17 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist













Name : MR. ROHIT NAYAKA M

Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240040 C/o

: Apollo Clinic

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM **Result Date** : 24-Feb-2024 12:55 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serun	1		al symptoms.	conjunction with clinic
Bilirubin Total-Serum	0.56	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.12	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.44	mg/dL	Male: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	22.00	U/L	Male: 15.0 - 37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	27.00	U/L	Male: 16.0 - 63.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)- Serum	101.00	U/L	Male: 45.0 - 117.0	PNPP,AMP- Buffer
Protein, Total-Serum	6.83	g/dL	6.40-8.20	Biuret/Endpoint- With Blank
Albumin-Serum	3.92	g/dL	Male: 3.40 - 5.50	Bromocresol Purple
Globulin-Serum	2.91	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serus	m 1.35	Ratio	0.80-2.0	Calculated

UHID

: 2402240040

2402240040



Printed By

: spectrum

Printed On : 24 Feb, 2024 06:17 pm


Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 5 of 11



+91 77604 97644 | 080 2337 1555









Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240040 C/o

: Apollo Clinic

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM : 24-Feb-2024 12:55 PM

Report Status : Final

Result Date

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	171.00	mg/dL	Male: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	89.00	mg/dL	Male: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	38.00	mg/dL	Male: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	133	mg/dL	Male: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	115.00	mg/dL	Male: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	18	mg/dL	Male: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	4.50	Ratio	Male: 0.0 - 5.0	Calculated

: 2402240040

2402240040

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum

Printed On : 24 Feb, 2024 06:17 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 6 of 11













Age / Gender : 46 years / Male

: Dr. APOLO CLINIC Ref. By Dr.

Reg. No. : 2402240040

: Apollo Clinic C/o

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM **Result Date** : 24-Feb-2024 12:55 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test)				
Blood Urea Nitrogen (BUN)- Serum	8.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	1.04	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	7.07	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	140.7	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.22	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	99.30	mmol/L	96.0-108.0	Ion-Selective Electrodes (ISE)
Calcium, Total- Serum	8.90	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein complexone)

: 2402240040

2402240040

UHID



Printed By

: spectrum

: 24 Feb, 2024 06:17 pm Printed On

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 7 of 11
Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010











Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

: 2402240040 Reg. No. C/o

: Apollo Clinic

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM **Result Date** : 24-Feb-2024 12:55 PM

: Final Report Status

Test Name	Result	Unit	Reference Value	Method
Gamma-Glutamyl Transferase (GGT)-Serum	28.00 U	U/L	Male: 15.0-85.0	Other g-Glut- 3-carboxy-4
			Female: 5.0-55.0	nitro

2402240040

: 2402240040

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.

Fasting Blood Sugar (FBS)-Plasma

96.00

mg/dL

60.0-110.0

Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By


: spectrum

Printed On

: 24 Feb, 2024 06:17 pm


Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 8 of 11





+91 77604 97644 | 080 2337 1555







Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

: 2402240040 Reg. No.

: Apollo Clinic C/o

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM

: 24-Feb-2024 12:55 PM

Report Status : Final

Result Date

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT) Serum	_			
Tri-Iodo Thyronine (T3)-Ser	um 1.01	ng/mL	Male: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	8.5	μg/dL	Male: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormon (TSH)-Serum	ne 4.91	μIU/mL	Male: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

: 2402240040

2402240040

UHID

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15

Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90,Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-

16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.


els: Graves disease, Autonomous thyroid hormone secretion, TSH defic


Printed By : spectrum


Printed On : 24 Feb, 2024 06:17 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru













Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

: 2402240040 Reg. No.

: Apollo Clinic C/o

: 24-Feb-2024 09:09 AM **Bill Date**

Sample Col. Date: 24-Feb-2024 09:09 AM

: 24-Feb-2024 01:00 PM

: Final

Result Date

Report Status

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	173	mg/dL	70-140	Hexo Kinase

2402240040

: 2402240040

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

UHID

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By


: spectrum


: 24 Feb, 2024 06:17 pm Printed On

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 10 of 11













Age / Gender : 46 years / Male

Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 2402240040

C/o : Apollo Clinic **Bill Date** : 24-Feb-2024 09:09 AM

Sample Col. Date: 24-Feb-2024 09:09 AM

: 24-Feb-2024 03:55 PM

Report Status : Final

Result Date

Unit Reference Value Method **Test Name** Result

UHID

Blood Group & Rh Typing-Whole Blood EDTA

Slide/Tube **Blood Group**

agglutination Negative Slide/Tube Rh Type agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.

: 2402240040

2402240040



Printed By : spectrum

: 24 Feb, 2024 06:17 pm Printed On

Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 11 of 11

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 www.spectrumdiagnostics.org info@spectrumdiagnostics.org +91 77604 97644 | 080 2337 1555



