PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 18/06/2022 4:21 PM

Printed On

MEDALI

Type : OP

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>

: 25/06/2022 5:15 PM

	<u>value</u>		<u>Reference interval</u>
<b>HAEMATOLOGY</b>			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	8.93	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	29.3	%	37 - 47
RBC Count (EDTA Blood)	4.73	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	62.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	18.9	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	30.5	g/dL	32 - 36
RDW-CV	20.7	%	11.5 - 16.0
RDW-SD	44.92	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	12680	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	58.81	%	40 - 75
Lymphocytes (Blood)	25.64	%	20 - 45
Eosinophils	10.86	%	01 - 06



(Blood)

**VERIFIED BY** 



PID No. : MED121139091 **Register On** : 18/06/2022 9:30 AM : 522219369 SID No. Collection On : 18/06/2022 11:37 AM Age / Sex : 43 Year(s) / Female Report On : 18/06/2022 4:21 PM

**Printed On** 



Type : OP

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Monocytes (Blood)	4.35	%	01 - 10
Basophils (Blood)	0.35	%	00 - 02
INTERPRETATION: Tests done on Automated Five Pa	art cell counter. All a	bnormal results are re	eviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	7.46	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	3.25	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	1.38	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.55	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.04	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	582.8	10^3 / μl	150 - 450
MPV (Blood)	7.55	fL	8.0 - 13.3
PCT (Automated Blood cell Counter)	0.44	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate)	51	mm/hr	< 20

: 25/06/2022 5:15 PM



(Citrated Blood)

**VERIFIED BY** 



 PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 18/06/2022 4:21 PM

**Printed On** 



Type : OP

Ref. Dr : MediWheel

Investigation BIOCHEMISTRY	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.29	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.13	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.16	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	6.29	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	7.21	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	21.00	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	71.42	U/L	42 - 98
Total Protein (Serum/Biuret)	6.94	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.85	gm/dl	3.5 - 5.2

3.09

1.25

gm/dL

: 25/06/2022 5:15 PM



Globulin

(Serum/Derived)
A: GRATIO

(Serum/Derived)

**VERIFIED BY** 



2.3 - 3.6

1.1 - 2.2

 PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 18/06/2022 4:21 PM



Type : OP

Ref. Dr : MediWheel

**Printed On** : 25/06/2022 5:15 PM

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	218.31	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	152.66	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_circulating level of triglycerides during most part of the day.

part of the day.			
HDL Cholesterol (Serum/Immunoinhibition)	36.78	mg/dL	Optimal(Negative Risk Factor): >= 60  Borderline: 50 - 59  High Risk: < 50
LDL Cholesterol (Serum/Calculated)	151	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	30.5	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	181.5	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



**VERIFIED BY** 



 PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 18/06/2022 4:21 PM



Type : OP

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

**INTERPRETATION:** 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2. It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 25/06/2022 5:15 PM

Total Cholesterol/HDL Cholesterol Ratio 5.9 (Serum/Calculated)

**Printed On** 

erum/Calculated)

Low Risk: 3.4 - 4.4

Average Risk: 4.5 - 7.1

Moderate Risk: 7.2 - 11.0

oderate Risk: 7.2 - 11.0 High Risk: > 11.0

Optimal: < 3.3

Triglyceride/HDL Cholesterol Ratio 4.2 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

(Serum/Calculated) High Risk: > 5.0

LDL/HDL Cholesterol Ratio 4.1 Optimal: 0.5 - 3.0

(Serum/Calculated)
Borderline: 3.1 - 6.0
High Risk: > 6.0



**VERIFIED BY** 



**Printed On** 

**Age / Sex** : 43 Year(s) / Female **Report On** : 18/06/2022 4:21 PM

Type : OP

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC</i> )	6.4	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 25/06/2022 5:15 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 136.98 mg/dL

(Whole Blood)

### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



**VERIFIED BY** 



 PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

**Age / Sex**: 43 Year(s) / Female **Report On**: 18/06/2022 4:21 PM

**Printed On** 

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

: 25/06/2022 5:15 PM

# **IMMUNOASSA**Y

### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.06 ng/ml 0.7 - 2.04

(Serum/ECLIA)

#### INTERPRETATION:

### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 7.58  $\mu$ g/dl 4.2 - 12.0

(Serum/ECLIA)

### INTERPRETATION:

### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 4.95 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

### **Comment:**

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



**VERIFIED BY** 



 PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 18/06/2022 4:21 PM

**Printed On** 



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	Value	Reference Interval

: 25/06/2022 5:15 PM

# **CLINICAL PATHOLOGY**

# PHYSICAL EXAMINATION (URINE COMPLETE)

Colour	Yellow	Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 20

(Urine)

# CHEMICAL EXAMINATION (URINE

**COMPLETE**)

pH 5.0 4.5 - 8.0

(Urine)

Specific Gravity 1.020 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)



**VERIFIED BY** 



 PID No.
 : MED121139091
 Register On
 : 18/06/2022 9:30 AM

 SID No.
 : 522219369
 Collection On
 : 18/06/2022 11:37 AM

 Age / Sex
 : 43 Year(s) / Female
 Report On
 : 18/06/2022 4:21 PM

**Printed On** 



Type : OP

Ref. Dr : MediWheel

: 25/06/2022 5:15 PM

InvestigationObserved<br/>ValueUnit<br/>Neference IntervalBiological<br/>Reference Interval

Glucose Negative Negative

(Urine/GOD - POD)

Leukocytes(CP) Negative

(Urine)

MICROSCOPIC EXAMINATION (URINE COMPLETE)

Pus Cells 0-1 /hpf NIL

(Urine)

Epithelial Cells (Urine) O-1 /hpf NIL

RBCs Nil /hpf NIL

(Urine)

Others

(Urine)

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.



**VERIFIED BY** 



**Printed On** 

 $\textbf{Age / Sex} \quad \textbf{:} \quad 43 \; \text{Year(s) / Female} \qquad \qquad \textbf{Report On} \qquad \textbf{:} \quad 18/06/2022 \; 4:21 \; \text{PM}$ 

Type : OP

Ref. Dr : MediWheel



InvestigationObservedUnitBiologicalValueReference Interval

: 25/06/2022 5:15 PM

# **IMMUNOHAEMATOLOGY**

BLOOD GROUPING AND Rh TYPING 'B' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.



**VERIFIED BY** 



PID No. : MED121139091 Register On : 18/06/2022 9:30 AM : 522219369 SID No.

Collection On : 18/06/2022 11:37 AM

Age / Sex : 43 Year(s) / Female

Report On 18/06/2022 4:21 PM



**Type** : OP

Ref. Dr

: MediWheel

**Printed On** : 25/06/2022 5:15 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<b>BIOCHEMISTRY</b>			
BUN / Creatinine Ratio	14.0		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	110.61	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	111.66	mg/dL	70 - 140

# **INTERPRETATION:**

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	8.2	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.57	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists, N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 4.55 2.6 - 6.0mg/dL (Serum/Enzymatic)



**VERIFIED BY** 



**APPROVED BY** 

-- End of Report --



Name	Ms.MUBUSHEERA	ID	MED121139091
Age & Gender	43/FEMALE	Visit Date	18/06/2022
Ref Doctor Name	MediWheel		

X-ray mammogram (mediolateral oblique & craniocaudal views) followed by Sonomammography.

### **BILATERAL MAMMOGRAPHY**

Breast composition Type B (These are scattered areas of fibroglandular density).

No evidence of focal soft tissue lesion.

No evidence of cluster microcalcification.

Subcutaneous fat deposition is within normal limits.

Bilateral axillary lymphnodes are seen.

### **BILATERAL SONOMAMMOGRAPHY**

Right breast shows well defined hypoechoic wider than tall lesion at 10 o' clock position, measuring about 8.5 x 3.4 mm.

Cysts are seen in both the breasts, measuring about 2.8 x 2.6 mm at 7 o' clock position in right breast and 4.0 x 3.6 mm at 12 o' clock position in left breast.

Right breasts shows dilated duct measuring 3.2 mm in maximum diameter seen at the 12 o' clock position.

Bilateral prominent ducts are seen, predominantly in the retroareolar region.

No evidence of axillary lymphadenopathy on both sides.

### **IMPRESSION:**

- Hypoechoic lesion in right breast likely fibroadenoma.
- Cysts in both the breast.
- Dilated duct in right breast.

**ASSESSMENT: BI-RADS CATEGORY - 3** 

# **BI-RADS CLASSIFICATION**

### REPORT DISCLAIMER

- 1.This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- 10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Ms.MUBUSHEERA	ID	MED121139091
Age & Gender	43/FEMALE	Visit Date	18/06/2022
Ref Doctor Name	MediWheel		

### **CATEGORY RESULT**

3 Probably benign finding. Short interval follow-up suggested.

DR. HEMANANDINI V.N
CONSULTANT RADIOLOGIST
Hn/Ss

### REPORT DISCLAIMER

<sup>1.</sup>This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiological reports should be interpreted in correlation with clinical and pathological findings.

<sup>2.</sup> The results reported here in are subject to interpretation by qualified medical professionals only.

<sup>3.</sup> Customer identities are accepted provided by the customer or their representative.

<sup>4.</sup>information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

<sup>5.</sup>If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

<sup>6.</sup>Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

<sup>7.</sup>Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

<sup>8.</sup>If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

<sup>9.</sup>Liability is limited to the extend of amount billed.

<sup>10.</sup>Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

<sup>11.</sup>Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.