





CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School , Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

PATIENT NAME : NIDHI ROSHAN (10000292)	PATIENT ID : NIDHF141287290
ACCESSION NO : 0290WB00605 AGE : 35 Years SEX : Female	ABHA NO :
DRAWN : RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCTOR : DR. ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:

Test Report Status	Eine al	Results	Biological Reference Interval Units
Test Report Status	<u>Final</u>	Results	Biological Reference Interval Onits

#### MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

BLOOD COUNTS,EDTA WHOLE BLOOD				
HEMOGLOBIN (HB)	11.9	Low	12.0 - 15.0	g/dL
METHOD : SPECTROPHOTOMETRY				
RED BLOOD CELL (RBC) COUNT	4.61		3.8 - 4.8	mil/µL
METHOD : ELECTRICAL IMPEDANCE				
WHITE BLOOD CELL (WBC) COUNT	5.90		4.0 - 10.0	thou/µL
METHOD : ELECTRICAL IMPEDANCE				
PLATELET COUNT	150		150 - 410	thou/µL
METHOD : ELECTRICAL IMPEDANCE				
RBC AND PLATELET INDICES				
HEMATOCRIT (PCV)	35.2	Low	36 - 46	%
METHOD : CALCULATED				
MEAN CORPUSCULAR VOLUME (MCV)	76.0	Low	83 - 101	fL
METHOD : CALCULATED				
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	25.8	Low	27.0 - 32.0	pg
METHOD : CALCULATED				
MEAN CORPUSCULAR HEMOGLOBIN	33.8		31.5 - 34.5	g/dL
CONCENTRATION (MCHC) METHOD : CALCULATED				
RED CELL DISTRIBUTION WIDTH (RDW)	14.3	High	11.6 - 14.0	%
METHOD : CALCULATED				
MENTZER INDEX	16.5			
MEAN PLATELET VOLUME (MPV)	14.4	High	6.8 - 10.9	fL
METHOD : CALCULATED				
WBC DIFFERENTIAL COUNT				
NEUTROPHILS	58		40 - 80	%
METHOD : IMPEDANCE / MICROSCOPY				
LYMPHOCYTES	35		20 - 40	%
METHOD : IMPEDANCE / MICROSCOPY				
MONOCYTES	05		2 - 10	%
METHOD : IMPEDANCE / MICROSCOPY				
EOSINOPHILS	02		1 - 6	%
METHOD : IMPEDANCE / MICROSCOPY				
BASOPHILS	00		0 - 2	%
METHOD : IMPEDANCE / MICROSCOPY				
ABSOLUTE NEUTROPHIL COUNT	3.42		2.0 - 7.0	thou/µL










CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

8800465156	Tel : 0731 2490008		
PATIENT NAME : NIDHI ROSHAN (1000	0292)	PATIENT ID : NI	DHF141287290
ACCESSION NO : 0290WB00605 AGE :	35 Years SEX : Female	ABHA NO :	
DRAWN : RECI	EIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 1	2:27
REFERRING DOCTOR : DR. ACROFEMI HEALT	HCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:	
Test Report Status <u>Final</u>	Results	Biological Reference Inte	erval Units
METHOD : CALCULATED			
ABSOLUTE LYMPHOCYTE COUNT METHOD : CALCULATED	2.06	1.0 - 3.0	thou/µL
ABSOLUTE MONOCYTE COUNT METHOD : CALCULATED	0.30	0.2 - 1.0	thou/µL
ABSOLUTE EOSINOPHIL COUNT METHOD : CALCULATED	0.12	0.02 - 0.50	thou/µL
ERYTHROCYTE SEDIMENTATION RATE ( BLOOD	ESR),WHOLE		
E.S.R METHOD : MODIFIED WESTERGREN	15	0 - 20	mm at 1 hr
GLUCOSE FASTING, FLUORIDE PLASMA			
FBS (FASTING BLOOD SUGAR) METHOD : HEXOKINASE	91	74 - 99	mg/dL
GLYCOSYLATED HEMOGLOBIN(HBA1C), BLOOD	EDTA WHOLE		
HBA1C	5.6	Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 Therapeutic goals: < 7.0 Action suggested : > 8.0 (ADA Guideline 2021)	%
METHOD : HPLC TECHNOLOGY			

ESTIMATED AVERAGE GLUCOSE(EAG)	114.0
GLUCOSE, POST-PRANDIAL, PLASMA	
PPBS(POST PRANDIAL BLOOD SUGAR)	103
METHOD : HEXOKINASE	
LIPID PROFILE, SERUM	
CHOLESTEROL, TOTAL	214
CHOLESTEROL, TOTAL	214
CHOLESTEROL, TOTAL METHOD : OXIDASE, ESTERASE, PEROXIDASE	214
	<b>214</b> 91
METHOD : OXIDASE, ESTERASE, PEROXIDASE	
METHOD : OXIDASE, ESTERASE, PEROXIDASE	

64



Borderline High: 150 - 199

< 116.0

199

**High** Desirable: <200

Desirable: < 150

High: 200 - 499

Normal: < 140,

Diabetic > or = 200

BorderlineHigh : 200-239 High : > or = 240

Impaired Glucose Tolerance:140-

METHOD : DIRECT- NON IMMUNOLOGICAL



HDL CHOLESTEROL

mg/dL

mg/dL

mg/dL

mg/dL







ſ

SRL LTD
Opposite St Raphael's Higher Secondary School , Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

PATIENT NAME: NIDHI ROSHAN (10000292)			PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605 ₁	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	<b>FOR :</b> DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Interv	al Units
CHOLESTEROL LDL	132	High	Adult levels: Optimal < 100 Near optimal/above optimal: 1 129 Borderline high : 130-159 High : 160-189	mg/dL 100-
NON HDL CHOLESTEROL	150	High	Very high : = 190 Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD : CALCULATED VERY LOW DENSITY LIPOPROTEIN METHOD : CALCULATED	18.2			mg/dL
CHOL/HDL RATIO	3.3			
LDL/HDL RATIO	2.1		0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate >6.0 High Risk	Risk









CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School , Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

PATIENT NAME : NIDHI ROSHAN (10000292)			PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	OR: DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:
Test Report Stat	tus <u>Final</u>	Results	Biological Reference Interval Units

#### Interpretation(s)

1) Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.

2) Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.

3)HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL

4) LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.

5)Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction.Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

#### Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

<b>Risk Category</b>			
Extreme risk group	A.CAD with $> 1$ feature of high risk group		
	B. CAD with $> 1$ feature of Very high risk	group or recurrent ACS (within 1 year) despite LDL-C	
	< or $=$ 50 mg/dl or polyvascular disease		
Very High Risk	1. Established ASCVD 2. Diabetes with 2	major risk factors or evidence of end organ damage 3.	
	Familial Homozygous Hypercholesterolemi	ia	
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end		
		DL >190 mg/dl 5. Extreme of a single risk factor. 6.	
	Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid		
	plaque		
Moderate Risk	2 major ASCVD risk factors		
Low Risk	0-1 major ASCVD risk factors		
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors			
1. Age $>$ or $=$ 45 years in males and $>$ or $=$ 55 years in females 3. Current Cigarette smoking or tobacco u			
2. Family history of p	oremature ASCVD	4. High blood pressure	
5. Low HDL			

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
Category A	< OR = 30)	< OR = 60)		









8800465156

CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA

SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

PATIENT NAME: NIDHI ROSHAN (10000292)	PATIENT ID : NIDHF141287290
ACCESSION NO : 0290WB00605 AGE : 35 Years SEX : Female	ABHA NO :
DRAWN : RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
<b>REFERRING DOCTOR :</b> DR. ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID :

Test Report Status         Final         Results         Biological Reference Interval         Unit	s
---	---

Extreme Risk Group Category B	<or 30<="" =="" th=""><th><or 60<="" =="" th=""><th>&gt; 30</th><th>&gt;60</th></or></th></or>	<or 60<="" =="" th=""><th>&gt; 30</th><th>&gt;60</th></or>	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR=100
Moderate Risk	<100	<130	>OR=100	>OR=130
Low Risk	<100	<130	>OR=130*	>OR=160

\*After an adequate non-pharmacological intervention for at least 3 months.

**References:** Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

### LIVER FUNCTION PROFILE, SERUM

,,			
BILIRUBIN, TOTAL	0.46	0.0 - 1.2	mg/dL
METHOD : JENDRASSIK AND GROFF			
BILIRUBIN, DIRECT	0.20	0.0 - 0.2	mg/dL
METHOD : DIAZOTIZATION			
BILIRUBIN, INDIRECT	0.26	0.00 - 1.00	mg/dL
METHOD : CALCULATED			
TOTAL PROTEIN	7.4	6.4 - 8.3	g/dL
METHOD : BIURET			
ALBUMIN	4.5	3.50 - 5.20	g/dL
METHOD : BROMOCRESOL GREEN			
GLOBULIN	2.9	2.0 - 4.1	g/dL
METHOD : CALCULATED			
ALBUMIN/GLOBULIN RATIO	1.6	1.0 - 2.0	RATIO
METHOD : CALCULATED			
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	16	UPTO 32	U/L
METHOD : UV WITH P5P			
ALANINE AMINOTRANSFERASE (ALT/SGPT)	14	UPTO 34	U/L
METHOD : UV WITH P5P			
ALKALINE PHOSPHATASE	65	35 - 104	U/L
METHOD : PNPP			
GAMMA GLUTAMYL TRANSFERASE (GGT)	11	5 - 36	U/L
METHOD : G-GLUTAMYL-CARBOXY-NITROANILIDE			
LACTATE DEHYDROGENASE	213	135 - 214	U/L
METHOD : ENZYMATIC LACTATE - PYRUVATE(IFCC)			
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	11	6 - 20	mg/dL
METHOD : UREASE KINETIC			













CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD	
Opposite St Raphael's Higher Secondary School , (	Old Seshore Road,
Residency Area	
INDORE, 452001	
Madhya Pradesh, India	
Tel : 0731 2490008	

CLIENT PATIENT ID:

PATIENT NAME : NIDHI ROSHAN (10000292)			PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )

Test Report Status	<u>Final</u>	Results		<b>Biological Reference Interva</b>	I Units
CREATININE		0.67		0.50 - 0.90	mg/dL
METHOD : ALKALINE PICRA	TE KINETIC JAFFES				
BUN/CREAT RATIO					
BUN/CREAT RATIO		16.42	High	5.0 - 15.0	
METHOD : CALCULATED					
URIC ACID, SERUM					
URIC ACID		4.0		2.6 - 6.0	mg/dL
METHOD : URICASE/CATALA	ASE UV				
TOTAL PROTEIN, SE	RUM				
TOTAL PROTEIN		7.4		6.4 - 8.3	g/dL
METHOD : BIURET					
ALBUMIN, SERUM					
ALBUMIN		4.5		3.5 - 5.2	g/dL
METHOD : BROMOCRESOL C	GREEN				
GLOBULIN					
GLOBULIN		2.9		2.0 - 4.1	g/dL
ELECTROLYTES (NA/	/K/CL), SERUM				
SODIUM, SERUM		143.2		136.0 - 146.0	mmol/L
METHOD : DIRECT ION SELE	ECTIVE ELECTRODE				
POTASSIUM, SERUM		4.96		3.50 - 5.10	mmol/L
METHOD : DIRECT ION SELE	ECTIVE ELECTRODE				
CHLORIDE, SERUM		105.6		98.0 - 106.0	mmol/L
METHOD : DIRECT ION SELE	ECTIVE ELECTRODE				







<u>Final</u>





# CLIENT CODE : C000138355

CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

**Biological Reference Interval** Units

PATIENT NAME: NIDHI ROSHAN (10000292)			PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN:		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	<b>FOR :</b> DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:

Results

## Interpretation(s)

**Test Report Status** 

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake,prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome,osmotic diuresis (e.g., hyperglycemia),alkalosis, familial periodic paralysis,trauma (transient).Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism,metabolic alkalosis. Drugs: chronic laxative,corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea),diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice,oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, high- dose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences:Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

### **PHYSICAL EXAMINATION, URINE**

COLOR	PALE YELLOW	
APPEARANCE	CLEAR	
CHEMICAL EXAMINATION, URINE		
PH	7.0	4.7 - 7.5
SPECIFIC GRAVITY	1.010	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED
MICROSCOPIC EXAMINATION, URINE		
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED



Page 7 Of 18 лж. 多回 

Scan to View Report £,

Ξ

/HPF







CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School , Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel : 0731 2490008

	Deculto	Dielegies   Deference Interval   Unite
<b>REFERRING DOCTOR :</b> DR.	ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:
DRAWN :	RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
ACCESSION NO : 0290WE	BOO605 AGE : 35 Years SEX : Female	ABHA NO :
PATIENT NAME : NIDHI	ROSHAN (10000292)	PATIENT ID : NIDHF141287290

Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
PUS CELL (WBC'S)	2-3	0-5	/HPF
EPITHELIAL CELLS	3-5	0-5	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	
REMARKS	Please note that all the urinary findings are confirmed manually as well.		nually as well.

### Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions		
Proteins	Inflammation or immune illnesses		
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment		
Glucose	Diabetes or kidney disease		
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst		
Urobilinogen	Liver disease such as hepatitis or cirrhosis		
Blood	Renal or genital disorders/trauma		
Bilirubin	Liver disease		
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases		
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions		
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time		
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein		
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases		
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice		
Uric acid	arthritis		
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.		
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis		

**THYROID PANEL, SERUM** 











ſ

SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

PATIENT NAME	: NIDHI ROSHAN	(10000292)	PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	TOR : DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	<b>Biological Reference Interval</b> Units
ТЗ	107.40	Non-Pregnant Women ng/dL 80.0 - 200.0 Pregnant Women 1st Trimester:105.0 - 230.0 2nd Trimester:129.0 - 262.0
METHOD : CHEMILUMINESCENCE TECHNOLOGY		3rd Trimester:135.0 - 262.0
T4	8.12	Non-Pregnant Women         µg/dL           5.10 - 14.10         Pregnant Women           1st Trimester: 7.33 - 14.80         2nd Trimester: 7.93 - 16.10           3rd Trimester: 6.95 - 15.70
TSH (ULTRASENSITIVE)	2.910	Non Pregnant Women µIU/mL 0.27 - 4.20 Pregnant Women 1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15
METHOD : CHEMILUMINESCENCE TECHNOLOGY		











**CLIENT'S NAME AND ADDRESS :** ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel : 0731 2490008

Test Report Sta	tus <u>Final</u>	Results	Biological Reference Interval Units
REFERRING DOCT	OR: DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
PATIENT NAME	: NIDHI ROSHAN	(10000292)	PATIENT ID : NIDHF141287290

#### Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

### ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

»»	BOTH THE LUNG FIELDS ARE
XRAY-CHEST	
METHOD : TUBE AGGLUTINATION	
RH TYPE	POSITIVE
METHOD : TUBE AGGLUTINATION	
ABO GROUP	TYPE A

BOTH THE LUNG FIELDS ARE CLEAR











CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD Opposite St Raphael's Higher Secondary School , Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel : 0731 2490008

30.0 and Above: Obese

PATIENT NAME	IE: NIDHI ROSHAN (10000292) PATIENT ID: NIDHF1412872		PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE: 35 Years SEX: Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	FOR: DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
»»	BOTH THE COSTOP	HRENIC AND CARIOPHRENIC ANGELS ARE CLEAR
»»	BOTH THE HILA AR	E NORMAL
»»	CARDIAC AND AOR	TTC SHADOWS APPEAR NORMAL
»»	BOTH THE DOMES	OF THE DIAPHRAM ARE NORMAL
»»	VISUALIZED BONY	THORAX IS NORMAL
IMPRESSION	NO ABNORMALITY I	DETECTED
	Dr G.S. Saluja, (M (Consultant Radiolo	
TMT OR ECHO		
TMT OR ECHO	NEGATIVE	
ECG		
ECG	WITHIN NORMAL L	IMITS
MEDICAL HISTORY		
RELEVANT PRESENT HISTORY	NOT SIGNIFICANT	
RELEVANT PAST HISTORY	NOT SIGNIFICANT	
RELEVANT PERSONAL HISTORY	NOT SIGNIFICANT	
RELEVANT FAMILY HISTORY	DM/HT - MOTHER	
OCCUPATIONAL HISTORY	NOT SIGNIFICANT	
HISTORY OF MEDICATIONS	NOT SIGNIFICANT	
ANTHROPOMETRIC DATA & BMI		
HEIGHT IN METERS	1.47	mts
WEIGHT IN KGS.	55	Kgs
BMI	25	BMI & Weight Status as follows: kg/sqmts Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight

### **GENERAL EXAMINATION**

MENTAL / EMOTIONAL STATE
PHYSICAL ATTITUDE
GENERAL APPEARANCE / NUTRITIONAL STATUS
BUILT / SKELETAL FRAMEWORK
FACIAL APPEARANCE
SKIN
UPPER LIMB

NORMAL NORMAL OVERWEIGHT AVERAGE NORMAL NORMAL NORMAL











RL LTD	
pposite St Raphael's Higher Secondary School, Old Seshore Road	ı,
esidency Area	
NDORE, 452001	
ladhya Pradesh, India	
el : 0731 2490008	

PATIENT NAME	TIENT NAME : NIDHI ROSHAN (10000292) PATIENT ID : NIDHF14128		PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	TOR: DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
LOWER LIMB	NORMAL	
NECK LYMPHATICS / SALIVARY GLANDS	NOT ENLARGED OR TENDI	EK
THYROID GLAND	NOT ENLARGED	
CAROTID PULSATION	NORMAL	
BREAST (FOR FEMALES)	NORMAL	
TEMPERATURE	AFEBRILE	
PULSE	67/MIN REGULAR, ALL PE BRUIT HEARD	RIPHERAL PULSES WELL FELT, NO CAROTID
RESPIRATORY RATE	NORMAL	
CARDIOVASCULAR SYSTEM		
BP	110/70	mm/Hg
PERICARDIUM	NORMAL	
APEX BEAT	NORMAL	
HEART SOUNDS	NORMAL	
MURMURS	ABSENT	
RESPIRATORY SYSTEM		
SIZE AND SHAPE OF CHEST	NORMAL	
MOVEMENTS OF CHEST	SYMMETRICAL	
BREATH SOUNDS INTENSITY	NORMAL	
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)	
ADDED SOUNDS	ABSENT	
PER ABDOMEN		
APPEARANCE	NORMAL	
VENOUS PROMINENCE	ABSENT	
LIVER	NOT PALPABLE	
SPLEEN	NOT PALPABLE	
HERNIA	ABSENT	
CENTRAL NERVOUS SYSTEM		
HIGHER FUNCTIONS	NORMAL	
CRANIAL NERVES	NORMAL	
CEREBELLAR FUNCTIONS	NORMAL	
SENSORY SYSTEM	NORMAL	
MOTOR SYSTEM	NORMAL	
REFLEXES	NORMAL	











CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

RL LTD	
pposite St Raphael's Higher Secondary School , Old Seshore Road	d,
esidency Area	
NDORE, 452001	
ladhya Pradesh, India	
el : 0731 2490008	
	_

PATIENT NAME	ME: NIDHI ROSHAN (10000292)		PATIENT ID : NIDHF141287290
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	<b>TOR :</b> DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID :

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
MUSCULOSKELETAL SYSTEM		
SPINE	NORMAL	
JOINTS	NORMAL	
BASIC EYE EXAMINATION		
CONJUNCTIVA	NORMAL	
EYELIDS	NORMAL	
EYE MOVEMENTS	NORMAL	
CORNEA	NORMAL	
DISTANT VISION RIGHT EYE WITH GLASSES	6/6 WITH GLASSES	S NORMAL
DISTANT VISION LEFT EYE WITH GLASSES	6/6 WITH GLASSES	S NORMAL
NEAR VISION RIGHT EYE WITH GLASSES	N6 WITHIN NORMA	AL LIMIT
NEAR VISION LEFT EYE WITH GLASSES	SION LEFT EYE WITH GLASSES N6 WITHIN NORMAL LIMIT	
COLOUR VISION	NORMAL	
BASIC ENT EXAMINATION		
EXTERNAL EAR CANAL	NORMAL	
TYMPANIC MEMBRANE	NORMAL	
NOSE	NO ABNORMALITY DETECTED	
SINUSES	NORMAL	
THROAT	NO ABNORMALITY	DETECTED
TONSILS	NOT ENLARGED	
BASIC DENTAL EXAMINATION		
TEETH	NORMAL	
GUMS	HEALTHY	
SUMMARY		
RELEVANT HISTORY	NOT SIGNIFICANT	
RELEVANT GP EXAMINATION FINDINGS	OVERWEIGHT	
REMARKS / RECOMMENDATIONS	NONE	
FITNESS STATUS		
EITNESS STATUS		I ADVICE) (AS PER REQUESTED PANEL OF TESTS)

FITNESS STATUS

FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)











**CLIENT'S NAME AND ADDRESS :** ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

Test Report Status <u>Final</u> Results		Biological Reference Interval Units
REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )		CLIENT PATIENT ID :
DRAWN :	RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
ACCESSION NO : 0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
PATIENT NAME : NIDHI ROSHA	N (10000292)	PATIENT ID : NIDHF141287290

Comments

CLINICAL FINDINGS:-

LOW HB.

DYSLIPIDEMIA.

OVER WEIGHT STATUS.

FITNESS STATUS :-

FITNESS STATUS : FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

ADVICE: WEIGHT REDUCTION, LOW FAT& CARBOHYDRATE DIET AND REGULAR PHYSICAL EXERCISE FOR OVERWEIGHT STATUS AND DYSLIPIDEMIA.

ADD TAKE FOOD STUFFS RICH IN IRON i.e. BEATROOT & SPINACH WITH IRON SUPPLEMENTS IN DIET. (NEEDS PHYSICIAN CONSULTATION IF HB < 8 gms%.)

NEED PHYSICIAN CONSULTATION FOR LIFE STYLE MODIFICATIO

#### Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

diagnosing a case of beta thalassaemia trait. WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR <

3.3, COVID-19 patients tend to show mild disease. (Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-**TEST DESCRIPTION** :-Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall

(sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging,

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

#### LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia False Decreased : Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine,











**CLIENT'S NAME AND ADDRESS :** ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

RL LTD	
pposite St Raphael's Higher Secondary School, Old Seshore	Road,
esidency Area	
NDORE, 452001	
ladhya Pradesh, India	
el : 0731 2490008	

Test Report Status Final Results		Biological Reference Interval Units		
REFERRING DOCTOR : DR. ACR	OFEMI HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID:		
DRAWN :	RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27		
ACCESSION NO : 0290WB006	<b>05</b> AGE : 35 Years SEX : Female	ABHA NO :		
PATIENT NAME : NIDHI ROS	5HAN (10000292)	PATIENT ID : NIDHF141287290		

#### salicvlates)

REFERENCE

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition. GLUCOSE FASTING, FLUORIDE PLASMA-**TEST DESCRIPTION** 

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

#### Increased in

Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

#### Decreased in

Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical,

stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin, ethanol, propranolol sulfonylureas,tolbutamide, and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within Individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control. High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.

2.Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes). The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for

well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.
1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
2. eAG gives an evaluation of blood glucose levels for the last couple of months.
3. eAG is calculated as eAG (mg/dl) = 28.7 \* HbA1c - 46.7

#### HbA1c Estimation can get affected due to :

anemia) will falsely lower HbA1c test results.Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II.Vitamin C & E are reported to falsely lower test results (possibly by inhibiting glycation of hemoglobin. III.Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.

IV.Interference of hemoglobinopathies in HbA1c estimation is seen in a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.) c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is

CLUCP > 25% of alternate partorni (boronate animity chronatography) is recommended for testing of HDATC.Abformal Henoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobin electrophoresis (HPLC method) is GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-LIVER FUNCTION PROFILE Bilirubin is a vellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give

yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin viral hepatitis). there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured

clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget"""'s disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson"""'s disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom"""s disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.Human serum albumin is the most abundant protein in human blood plasma.It is produced in the liver.Albumin constitutes about









**CLIENT'S NAME AND ADDRESS :** ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
Dpposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
NDORE, 452001
Madhya Pradesh, India
Tel : 0731 2490008

Test Report Status Final	Results	Biological Reference Interval Units
REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD ( MEDIWHEEL )		CLIENT PATIENT ID :
DRAWN :	RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
ACCESSION NO : 0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
PATIENT NAME : NIDHI ROSHA	N (10000292)	PATIENT ID : NIDHF141287290

half of the blood serum protein.Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol,

Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism) Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

Blockage in the urinary tract

Kidney problems, such as kidney damage or failure, infection, or reduced blood flow

Loss of body fluid (dehydration)
Muscle problems, such as breakdown of muscle fibers

• Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to: Myasthenia Gravis

, Muscular dystrophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic svndrome

Causes of decreased levels-Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM-Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum...Protein in the plasma is made up of albumin and globulin

syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc. ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface

of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

MEDICAL

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

FITNESS STATUS-Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for .These are then further correlated with details of the job under consideration to eventually fit the right man to the right job. Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:

• Fit (As per requested panel of tests) - SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.

• Fit (with medical advice) (As per requested panel of tests) - This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary Iffestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician's consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.
 Fitness on Hold (Temporary Unfit) (As per requested panel of tests) - Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal

the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.

Unfit (As per requested panel of tests) - An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.











CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD ( MEDIWHEEL ) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRLLID
Opposite St Raphael's Higher Secondary School , Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

PATIENT NAME: NIDHI ROSHAN (10000292)		PATIENT ID : NIDHF141287290	
ACCESSION NO :	0290WB00605	AGE : 35 Years SEX : Female	ABHA NO :
DRAWN :		RECEIVED : 27/02/2023 08:36	REPORTED : 28/02/2023 12:27
REFERRING DOCT	OR: DR. ACROFEM	I HEALTHCARE LTD ( MEDIWHEEL )	CLIENT PATIENT ID :
ſ			

Test Report Status Final

Results

Units

#### MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

#### **ULTRASOUND ABDOMEN**

ULTRASOUND ABDOMEN

### NO ABNORMALITIES DETECTED

#### Comments

U.S.G OF WHOLE ABDOMEN

Liver is normal in size, shape with with smooth outline. Parenchymal echotexture is homogeneous. Intra & Extra hepatic biliary radicals are normal. Portal vein and C.B.D are normal in caliber.

Gall Bladder is normal, thin walled & its lumen is echo free.

Spleen is normal in size, shape & echotexture.

Pancreas is normal in size, shape & echotexture.

Both Kidneys are normal in size, shape and echotexture. Central pelvicalyceal system is normal. Corticomedullary differentiation is maintained.

IVC and AO is normal in caliber.No lymphadenopathy. Urinary Bladder is normal thin walled,there is no calculus.

Uterus is anteverted and normal in size. Myometrial echotexture is homogeneous Endometrial echo reflection is normal. Cervix and endocervical canal appears normal.

Bilateral Ovaries are normal in size, shape and echotexture.

IMPRESSION- No Significant abnormality seen in USG of Whole Abdomen.

Dr G S Saluja MBBS, DMRD (Consultant Radiologist)

\*\*End Of Report\*\* Please visit www.srlworld.com for related Test Information for this accession

Dr.Arpita Pasari, MD Consultant Pathologist









SRL LTD
Opposite St Raphael's Higher Secondary School, Old Seshore Road,
Residency Area
INDORE, 452001
Madhya Pradesh, India
Tel: 0731 2490008

Test Report Sta	tus <u>Final</u>	F	Results			Units
REFERRING DOCT	OR: DR. ACROFEM	11 HEALTHCARE LTD	( MEDIWHEEL )	CLIEN	T PATIENT ID	:
DRAWN :		RECEIVED : 27/0	2/2023 08:36	REPORTED :	28/02/20	23 12:27
ACCESSION NO :	0290WB00605	AGE: 35 Years	SEX : Female	ABHA NO :		
PATIENT NAME	: NIDHI ROSHAN	N (10000292)		PA	TIENT ID:	NIDHF141287290

CONDITIONS OF LABORAT	ORY TESTING & REPORTING
<ol> <li>It is presumed that the test sample belongs to the patient named or identified in the test requisition form.</li> <li>All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.</li> <li>Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.</li> <li>A requested test might not be performed if:         <ol> <li>Specimen received is insufficient or inappropriate</li> <li>Specimen quality is unsatisfactory</li> <li>Incorrect specimen type</li> <li>Discrepancy between identification on specimen container label and test requisition form</li> </ol> </li> </ol>	<ol> <li>SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety &amp; technical integrity.</li> <li>Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.</li> <li>Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.</li> <li>Test results cannot be used for Medico legal purposes.</li> <li>In case of queries please call customer care (91115 91115) within 48 hours of the report.</li> </ol>
	SRL Limited
	Fortis Hospital, Sector 62, Phase VIII, Mohali 160062



