

PATIENT NAME: ASHWINI RATHOD REF. DOCTOR: SELF

CODE/NAME & ADDRESS : C000138382

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: **0075WA002029**PATIENT ID : ASHWF04068875

CLIENT PATIENT ID: ABHA NO : DRAWN :28/01/2023 09:59:51
RECEIVED :28/01/2023 10:01:12
REPORTED :30/01/2023 11:25:40

:34 Years

AGE/SEX

Test Report Status <u>Final</u> Results Biological Reference Interval Units

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

XRAY-CHEST

»» BOTH THE LUNG FIELDS ARE CLEAR

»»
BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR

»» BOTH THE HILA ARE NORMAL

»» CARDIAC AND AORTIC SHADOWS APPEAR NORMAL»» BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL

»» VISUALIZED BONY THORAX IS NORMAL

IMPRESSION NO ABNORMALITY DETECTED

METHOD: MICROSCOPIC EXAMINATION

ECG

ECG WITHIN NORMAL LIMITS

MEDICAL HISTORY

RELEVANT PRESENT HISTORY

RELEVANT PAST HISTORY

RELEVANT PERSONAL HISTORY

RELEVANT FAMILY HISTORY

NOT SIGNIFICANT

NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.75 mts WEIGHT IN KGS. 85 Kgs

BMI 28 BMI & Weight Status as follows/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE NORMAL PHYSICAL ATTITUDE NORMAL TEMPERATURE NORMAL

PULSE REGULAR, ALL PERIPHERAL PULSES WELL FELT

RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM

BP 110/90 mm/Hg

PERICARDIUM NORMAL

Dr. Anamika Pal Lab Head



Page 1 Of 16

View Details

View Report

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email: wellness.itpl@srl.in



CODE/NAME & ADDRESS: C000138382

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: **0075WA002029**PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

AGE/SEX :34 Years Female
DRAWN :28/01/2023 09:59:51
RECEIVED :28/01/2023 10:01:12
REPORTED :30/01/2023 11:25:40

Test Report Status <u>Final</u> Results Biological Reference Interval Units

NORMAL

APEX BEAT NORMAL

HEART SOUNDS S1, S2 HEARD NORMALLY

BASIC EYE EXAMINATION

DISTANT VISION RIGHT EYE WITH GLASSES NORMAL DISTANT VISION LEFT EYE WITH GLASSES NORMAL

NEAR VISION RIGHT EYE WITH GLASSES WITHIN NORMAL LIMIT
NEAR VISION LEFT EYE WITH GLASSES WITHIN NORMAL LIMIT

BASIC DENTAL EXAMINATION

TEETH NORMAL GUMS HEALTHY

SUMMARY

COLOUR VISION

RELEVANT HISTORY

RELEVANT GP EXAMINATION FINDINGS

RELEVANT LAB INVESTIGATIONS

PRE DIABETIC
DYSLIPEDIMIA

DYSLIPEDIMIA
ADVICE:TAB TONACT TG 0-0-1x3MONTHS

RELEVANT NON PATHOLOGY DIAGNOSTICS NO ABNORMALITIES DETECTED REMARKS / RECOMMENDATIONS NONE

FITNESS STATUS

FITNESS STATUS FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

Comments

*NOTE: NON PATHOLOGY TESTS ARE REVIEWED BY Consultant Physician: Dr.RITESH RAJ MBBS,CCEBDM

Radiologist : Dr.THILAK BABU Dental Doctor:Dr Ashish sinha BDS,

Interpretation(s)

MEDICAL

Dr. Anamika Pal

(対数回 回 なな(な 59





Page 2 Of 16

View Details

View Report



REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0075WA002029 PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN

RECEIVED: 28/01/2023 10:01:12

REPORTED :30/01/2023 11:25:40

Test Report Status Biological Reference Interval <u>Final</u> Results Units

FITNESS STATUS-Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history; as well as the comprehensiveness of the diagnostic panel which has been requested for .These are then further correlated with details of the job under consideration to eventually fit the right man to the right job.

Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:

- Fit (As per requested panel of tests) SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.
- Fit (with medical advice) (As per requested panel of tests) This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician's consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.

 • Fitness on Hold (Temporary Unfit) (As per requested panel of tests) - Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal
- the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.
- Unfit (As per requested panel of tests) An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.

Dr. Anamika Pal Lab Head



Page 3 Of 16

View Report

PERFORMED AT:

SRL Ltd 7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956



CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0075WA002029 PATIENT ID : ASHWF04068875

CLIENT PATIENT ID: ABHA NO

AGE/SEX :34 Years Female DRAWN :28/01/2023 09:59:51 RECEIVED: 28/01/2023 10:01:12 REPORTED :30/01/2023 11:25:40

Biological Reference Interval Test Report Status Results Units <u>Final</u>

| н | AEMATOLOGY - CBC | | |
|---|--------------------------------|---------------------------|---------|
| MEDI WHEEL FULL BODY HEALTH CHECKUP BE | LOW 40FEMALE | | |
| BLOOD COUNTS, EDTA WHOLE BLOOD | | | |
| HEMOGLOBIN (HB) | 12.9 | 12.0 - 15.0 | g/dL |
| RED BLOOD CELL (RBC) COUNT | 4.75 | 3.8 - 4.8 | mil/μL |
| WHITE BLOOD CELL (WBC) COUNT | 9.60 | 4.0 - 10.0 | thou/µL |
| PLATELET COUNT | 281 | 150 - 410 | thou/µL |
| RBC AND PLATELET INDICES | | | |
| HEMATOCRIT (PCV) | 40.0 | 36 - 46 | % |
| MEAN CORPUSCULAR VOLUME (MCV) | 84 | 83 - 101 | fL |
| MEAN CORPUSCULAR HEMOGLOBIN (MCH) | 27.5 | 27.0 - 32.0 | pg |
| MEAN CORPUSCULAR HEMOGLOBIN | 32.2 | 31.5 - 34.5 | g/dL |
| CONCENTRATION (MCHC) | 14.0 | 11.6 14.0 | % |
| RED CELL DISTRIBUTION WIDTH (RDW) | 14.0 17.7 | 11.6 - 14.0 | 90 |
| MENTZER INDEX | 7.4 | 6.0. 10.0 | fL |
| MEAN PLATELET VOLUME (MPV) WBC DIFFERENTIAL COUNT | 7.4 | 6.8 - 10.9 | IL |
| NEUTROPHILS | 64 | 40 - 80 | % |
| LYMPHOCYTES | 27 | 20 - 40 | % |
| MONOCYTES | 6 | 2 - 10 | % |
| EOSINOPHILS | 2 | 1 - 6 | % |
| BASOPHILS | 1 | 0 - 2 | % |
| ABSOLUTE NEUTROPHIL COUNT | 6.14 | 2.0 - 7.0 | thou/µL |
| ABSOLUTE LYMPHOCYTE COUNT | 2.59 | 1.0 - 3.0 | thou/µL |
| ABSOLUTE MONOCYTE COUNT | 0.58 | 0.2 - 1.0 | thou/µL |
| ABSOLUTE EOSINOPHIL COUNT | 0.19 | 0.02 - 0.50 | thou/µL |
| ABSOLUTE BASOPHIL COUNT | 0.10 | 0.02 - 0.10 | thou/µL |
| NEUTROPHIL LYMPHOCYTE RATIO (NLR) | 2.4 | 0.02 0.10 | |
| MORPHOLOGY | | | |
| RBC | NORMOCYTIC NORMOCH | HROMIC | |
| WBC | | RPHOLOGY AND DISTRIBUTION | |
| PLATELETS | ADEQUATE NO HEMOPARASITES SEEN | | |

Dr. Anamika Pal Lab Head





Page 4 Of 16

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email: wellness.itpl@srl.in



REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

:28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12

:34 Years

AGE/SEX

REPORTED :30/01/2023 11:25:40

Test Report Status Results **Biological Reference Interval** <u>Final</u> Units

ABHA NO

IMPRESSION

NORMOCYTIC NORMOCHROMIC BLOOD PICTURE

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

Dr. Anamika Pal Lab Head





Page 5 Of 16

View Report

SRL Ltd 7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA

Tel: 9111591115, Fax:





mm at 1 hr

REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12

REPORTED :30/01/2023 11:25:40

Test Report Status Biological Reference Interval <u>Final</u> Results Units

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD

0 - 20E.S.R

METHOD: MODIFIED WESTERGREN

Interpretation(s)
ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION:

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine, salicylates)

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

Dr. Anamika Pal Lab Head





Page 6 Of 16

PERFORMED AT:

SRL Ltd

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email: wellness.itpl@srl.in



CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

Test Report Status

8800465156

ACCESSION NO: 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12

REPORTED :30/01/2023 11:25:40

Biological Reference Interval Units

IMMUNOHAEMATOLOGY

Results

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

<u>Final</u>

TYPE B **ABO GROUP** RH TYPE **POSITIVE**

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

Dr. Anamika Pal Lab Head





Page 7 Of 16

SRL Ltd

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email: wellness.itpl@srl.in



CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 RECEIVED: 28/01/2023 10:01:12 REPORTED :30/01/2023 11:25:40

Biological Reference Interval Test Report Status Results Units **Final**

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR)

112 High

74 - 99

mg/dL

METHOD: SPECTROPHOTOMETRY HEXOKINASE

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE

BLOOD HBA1C

6.0 High

Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 %

Diabetics: > or = 6.5ADA Target: 7.0

Action suggested: > 8.0

METHOD: PARTICLE-ENHANCED TURBIDIMETRIC INHIBITION IMMUNOASSAY(PETINIA)

ESTIMATED AVERAGE GLUCOSE(EAG)

125.5 High < 116.0 mg/dL

METHOD: PARTICLE-ENHANCED TURBIDIMETRIC INHIBITION IMMUNOASSAY(PETINIA)

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR)

120

70 - 139

mg/dL

METHOD: SPECTROPHOTOMETRY HEXOKINASE

LIPID PROFILE, SERUM

238 High CHOLESTEROL, TOTAL

< 200 Desirable

mg/dL

200 - 239 Borderline High >/= 240 High

METHOD: SPECTROPHOTOMETRY, CHOLESTEROL OXIDASE ESTERASE PEROXIDASE

TRIGLYCERIDES 400 High

< 150 Normal

mg/dL

150 - 199 Borderline High

200 - 499 High

>/=500 Very High

METHOD: LIPOPROTEIN LIPASE (LPL), GLYCEROL KINASE (GK)

HDL CHOLESTEROL 32 Low < 40 Low

mg/dL

>/=60 High

METHOD: DIRECT HDL, PEGME

CHOLESTEROL LDL 126 High < 100 Optimal

mg/dL

100 - 129

Near optimal/ above optimal

130 - 159 Borderline High 160 - 189 High

>/= 190 Very High

Dr. Anamika Pal Lab Head



Page 8 Of 16

PERFORMED AT:

SRL Ltd

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA





PATIENT NAME: ASHWINI RATHOD REF. DOCTOR: SELF CODE/NAME & ADDRESS: C000138382 ACCESSION NO: 0075WA002029 AGE/SEX :34 Years Female ACROFEMI HEALTHCARE LTD (MEDIWHEEL) PATIENT ID : ASHWF04068875 DRAWN :28/01/2023 09:59:51 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 28/01/2023 10:01:12 DELHI REPORTED :30/01/2023 11:25:40 **NEW DELHI 110030** ABHA NO 8800465156

| Test Report Status <u>Final</u> | Results | Biological Reference | Interval Units |
|--|-----------|---|----------------|
| | | | |
| METHOD: DIRECT ENZYME CLEARANCE | 206 High | Darkeller Lare Hear | 120 |
| NON HDL CHOLESTEROL | 206 High | Desirable: Less than Above Desirable: 13 | |
| | | Borderline High: 160 | |
| | | High: 190 - 219 | |
| | | Very high: $>$ or $= 22$ | 20 |
| METHOD: CALCULATED PARAMETER | | | |
| VERY LOW DENSITY LIPOPROTEIN | 80.0 High | = 30.0</td <td>mg/dL</td> | mg/dL |
| CHOL/HDL RATIO | 7.4 High | 3.3 - 4.4 | |
| | | Low Risk 4.5 - 7.0 | |
| | | Average Risk | |
| | | 7.1 - 11.0 | |
| | | Moderate Risk | |
| | | > 11.0 | |
| | | High Risk | |
| LDL/HDL RAΠO | 3.9 High | 0.5 - 3.0 Desirable/L | |
| | | 3.1 - 6.0 Borderline/ Risk | Moderate |
| | | >6.0 High Risk | |
| Interpretation(s) | | J | |
| LIVER FUNCTION PROFILE, SERUM | | | |
| BILIRUBIN, TOTAL | 0.40 | 0.2 - 1.0 | mg/dL |
| METHOD : SPECTROPHOTOMETRY | | | |
| BILIRUBIN, DIRECT | 0.1 | 0.0 - 0.2 | mg/dL |
| METHOD: SPECTROPHOTOMETRY | | | |
| BILIRUBIN, INDIRECT | 0.3 | 0.1 - 1.0 | mg/dL |
| METHOD: CALCULATED PARAMETER | | | |
| TOTAL PROTEIN | 7.9 | 6.4 - 8.2 | g/dL |
| METHOD: SPECTROPHOTOMETRY, MODIFIED BIURET | | | |
| ALBUMIN | 3.8 | 3.4 - 5.0 | g/dL |
| METHOD: SPECTROPHOTOMETRIC - BROMOCRESOL GREEN (| | | |
| GLOBULIN | 4.1 | 2.0 - 4.1 | g/dL |
| METHOD: CALCULATED PARAMETER | | | |
| ALBUMIN/GLOBULIN RATIO | 0.9 Low | 1.0 - 2.1 | RATIO |
| METHOD: CALCULATED PARAMETER | | | |

Dr. Anamika Pal Lab Head





Page 9 Of 16

SRL Ltd

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956 Email: wellness.itpl@srl.in



CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0075WA002029 PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN RECEIVED : 28/01/2023 10:01:12 REPORTED :30/01/2023 11:25:40

| | <u> </u> | I | |
|---|---------------|----------------------|------------------|
| Test Report Status <u>Final</u> | Results | Biological Reference | e Interval Units |
| | | | |
| ASPARTATE AMINOTRANSFERASE (AST/SGOT) METHOD: SPECTROPHOTOMETRY, UV WITH PYRIDOXAL -5-PHO | 15 | 15 - 37 | U/L |
| | 22 | < 34.0 | U/L |
| ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: SPECTROPHOTOMETRY, UV WITH PYRIDOXAL -5-PHO | | < 34.0 | U/L |
| ALKALINE PHOSPHATASE METHOD: SPECTROPHOTOMETRY | 83 | 30 - 120 | U/L |
| GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: SPECTROPHOTOMETRY, G-GLUTAMYL-CARBOXY-NITR | 51 ONILIDE | 5 - 55 | U/L |
| LACTATE DEHYDROGENASE METHOD: SPECTROPHOTOMETRY | 159 | 100 - 190 | U/L |
| BLOOD UREA NITROGEN (BUN), SERUM | | | |
| BLOOD UREA NITROGEN | 7 | 6 - 20 | mg/dL |
| CREATININE, SERUM | | | |
| CREATININE | 0.6 | 0.60 - 1.10 | mg/dL |
| METHOD: SPECTROPHOTOMETRIC, JAFFE'S KINETICS | | | |
| BUN/CREAT RATIO | | | |
| BUN/CREAT RATIO | 11.67 | 5.00 - 15.00 | |
| URIC ACID, SERUM | | | |
| URIC ACID | 5.0 | 2.6 - 6.0 | mg/dL |
| METHOD: SPECTROPHOTOMETRY | | | |
| TOTAL PROTEIN, SERUM | | | |
| TOTAL PROTEIN | 7.9 | 6.4 - 8.2 | g/dL |
| METHOD : SPECTROPHOTOMETRY, MODIFIED BIURET | | | |
| ALBUMIN, SERUM | | | |
| ALBUMIN | 3.8 | 3.4 - 5.0 | g/dL |
| METHOD: SPECTROPHOTOMETRIC - BROMOCRESOL GREEN (BC | .G) | | |
| GLOBULIN | 4.1 | 2.0 4.1 | الماري |
| GLOBULIN METHOD: CALCULATED PARAMETER | 4.1 | 2.0 - 4.1 | g/dL |
| ELECTROLYTES (NA/K/CL), SERUM | | | |
| SODIUM, SERUM | 141.6 | 137 - 145 | mmol/L |
| POTASSIUM, SERUM | 4.76 | 3.6 - 5.0 | mmol/L |
| CHLORIDE, SERUM | 108.7 High | 98 - 107 | mmol/L |
| Interpretation(s) | ingn | JO 107 | |
| interpretation(s) | | | |

Dr. Anamika Pal





Page 10 Of 16

Patient Ref. No. 775000002221426

Lab Head



REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO : 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID: ABHA NO

:34 Years :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12

AGE/SEX

REPORTED :30/01/2023 11:25:40

Test Report Status Results **Biological Reference Interval Final** Units

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

Increased in

Diabetes mellitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs:corticosteroids, phenytoin, estrogen, thiazides.

Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency, hypopituitarism,diffuse liver disease, malignancy (adrenocortical, stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin, ethanol, propranolol; sulfonylureas,tolbutamide, and other oral hypoglycemic agents

NOTE:

While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals.Thus,

glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1.Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2.Diagnosing diabetes.
- 3.Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- 1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
 2. eAG gives an evaluation of blood glucose levels for the last couple of months.
 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :

1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II.Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

III.Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results. IV.Interference of hemoglobinopathies in HbA1c estimation is seen in

a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.

b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is

C.HDF > 25% on atternate pattorni (boroliate animity circumatography) is recommended for detecting a hemoglobinopathy
GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE

LIVER FUNCTION PROFILE, SERUM-LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis,

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction,

Dr. Anamika Pal Lab Head



Page 11 Of 16

View Report

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956



REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO : 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID: ABHA NO

:34 Years :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12

AGE/SEX

REPORTED :30/01/2023 11:25:40

Test Report Status Biological Reference Interval Final Results Units

Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget''''s disease,Rickets,Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia,Malnutrition,Protein deficiency,Wilson'''s disease.GGT is an enzyme found in cell membranes of many tissues mainly in the liver,kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection,including HIV and hepatitis B or C,Multiple myeloma,Waldenstrom'''s disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.Human serum albumin is the most abundant protein in human blood plasma.It is produced in the liver.Albumin constitutes about half of the blood serum protein.Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
 Loss of body fluid (dehydration)
- · Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis
- Muscular dystroph

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome

Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis
TOTAL PROTEIN, SERUM-Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum..Protein in the plasma is made up of albumin and globulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom''''''''''''' disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc.

Dr. Anamika Pal

Lab Head

Page 12 Of 16





View Report

PERFORMED AT:

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA





CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0075WA002029

PATIENT ID : ASHWF04068875

CLIENT PATIENT ID: ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12

REPORTED :30/01/2023 11:25:40

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

APPEARANCE CLEAR

CHEMICAL EXAMINATION, URINE

PH 5.5 4.7 - 7.51.003 - 1.035 SPECIFIC GRAVITY 1.005 **PROTEIN** NOT DETECTED NOT DETECTED **GLUCOSE** NOT DETECTED NOT DETECTED **KETONES** NOT DETECTED NOT DETECTED **BLOOD** NOT DETECTED NOT DETECTED **BILIRUBIN** NOT DETECTED NOT DETECTED UROBILINOGEN **NORMAL NORMAL NITRITE** NOT DETECTED NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

LEUKOCYTE ESTERASE

/HPF RED BLOOD CELLS NOT DETECTED NOT DETECTED /HPF PUS CELL (WBC'S) 2-3 0-5 EPITHELIAL CELLS 3-5 0-5 /HPF

NOT DETECTED

NOT DETECTED

NOT DETECTED **CASTS** CRYSTALS NOT DETECTED

BACTERIA NOT DETECTED NOT DETECTED YEAST NOT DETECTED NOT DETECTED

Interpretation(s)

Dr. Anamika Pal

Page 13 Of 16





PERFORMED AT:

SRL Ltd

Lab Head

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA





REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0075WA002029 PATIENT ID : ASHWF04068875

CLIENT PATIENT ID: ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12 REPORTED :30/01/2023 11:25:40

Test Report Status Results **Biological Reference Interval** Units **Final**

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Dr. Anamika Pal Lab Head





Page 14 Of 16

SRL Ltd

7/3, SRINARAYANI ARCADE 1ST FLOOR, ABOVE BATA SHOWROOM BROOKEFIELD MAIN ROAD, KUNDALAHALLI BANGALORE, 560066 KARNATAKA, INDIA





REF. DOCTOR: SELF PATIENT NAME: ASHWINI RATHOD

CODE/NAME & ADDRESS: C000138382 ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0075WA002029 PATIENT ID : ASHWF04068875

CLIENT PATIENT ID:

ABHA NO

AGE/SEX :34 Years Female :28/01/2023 09:59:51 DRAWN RECEIVED: 28/01/2023 10:01:12 REPORTED :30/01/2023 11:25:40

Biological Reference Interval Test Report Status Results Units **Final**

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

THYROID PANEL, SERUM

ng/dL T3 111.0 Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester: 105.0 - 230.0 2nd Trimester: 129.0 - 262.0 3rd Trimester: 135.0 - 262.0 T4 8.66 Non-Pregnant Women µg/dL 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70 9.360 High Non Pregnant Women µIU/mL TSH (ULTRASENSITIVE) 0.27 - 4.20

Pregnant Women

1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyporthyroidism, TSH levels are low. owidctlparowidctlparBelow mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

| Sr. No. | TSH | Total T4 | FT4 | Total T3 | Possible Conditions |
|---------|------|----------|-----|----------|--|
| 1 | High | Low | Low | Low | (1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3) Post Thyroidectomy (4) Post Radio-Iodine treatment |



Dr. Prajwal A, MD CONSULTANT BIOCHEMIST (SECTION HEAD)





Page 15 Of 16



154/9, BANNERGHATTA ROAD, OPP. IIM-B, BANGALORE, 560076 KARNATAKA, INDIA Tel: 80-66214444, Fax: CIN - U74899PB1995PLC045956





PATIENT NAME: ASHWINI RATHOD REF. DOCTOR: SELF CODE/NAME & ADDRESS: C000138382 ACCESSION NO: 0075WA002029 AGE/SEX :34 Years Female ACROFEMI HEALTHCARE LTD (MEDIWHEEL) PATIENT ID : ASHWF04068875 :28/01/2023 09:59:51 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 28/01/2023 10:01:12 DELHI REPORTED :30/01/2023 11:25:40 **NEW DELHI 110030** ABHA NO 8800465156

Test Report Status <u>Final</u> Results Biological Reference Interval Units

| 2 | High | Normal | Normal | Normal | (1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid |
|---|--|---|--------|--------|--|
| | | 350000000000000000000000000000000000000 | | | hormone replacement therapy (3) In cases of Autoimmune/Hashimoto |
| | | | | | thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical |
| | | | | | inflammation, drugs like amphetamines, Iodine containing drug and |
| | | | | | dopamine antagonist e.g. domperidone and other physiological reasons. |
| 3 | Normal/Low | Low | Low | Low | (1) Secondary and Tertiary Hypothyroidism |
| 4 | Low | High | High | High | (1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre |
| | | | | 1 2 | (3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid |
| | | | | | hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4 |
| | | | | | replacement therapy (7) First trimester of Pregnancy |
| 5 | Low | Normal | Normal | Normal | (1) Subclinical Hyperthyroidism |
| 6 | High | High | High | High | (1) TSH secreting pituitary adenoma (2) TRH secreting tumor |
| 7 | Low | Low | Low | Low | (1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent |
| | 0.00, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, 0.000, | | | | treatment for Hyperthyroidism |
| 8 | Normal/Low | Normal | Normal | High | (1) T3 thyrotoxicosis (2) Non-Thyroidal illness |
| 9 | Low | High | High | Normal | (1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies |

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

End Of Report
Please visit www.srlworld.com for related Test Information for this accession



Dr. Prajwal A, MD CONSULTANT BIOCHEMIST (SECTION HEAD)





Page 16 Of 16

View Details



PERFORMED AT:

SKL Ltd 154/9, BANNERGHATTA ROAD, OPP. IIM-B, BANGALORE, 560076 KARNATAKA, INDIA Tel: 80-66214444, Fax: CIN - U74899PB1995PLC045956

