







## LAB DIVISION

Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Gender / Age Male / 30 Yrs

Refd. By

. Apollo Health & Lifestyle Ltd Client

Collected On 22/07/2023 10:06:37 22/07/2023 10:06:39 Received On Released On 22/07/2023 17:53:07

24/07/2023 18:03:37 Printed On

Investigation Value Unit Biological Ref. Range

# **HAEMATOLOGY**

# Peripheral Blood Smear

RBC:-RBC are Normocytic Normochromic.

WBC:-WBC Shows normal morphology.

PLATELET:- Platelets are adequate with normal morphology.

PARASITES:- Malaria parasites are not detected.

REMARKS:- Unremarkable P/S

Dr. Vishal Bhuva MBBS, MD Pathologist, HAM

- Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.
   2. The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently
   3. These reports are not valid for medico legal purposes.







Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Gender / Age Male / 30 Yrs

Refd. By

Client . Apollo Health & Lifestyle Ltd



Collected On	22/07/2023 10:06:37
Received On	22/07/2023 10:06:39
Released On	22/07/2023 17:27:38
Printed On	24/07/2023 18:03:40

Investigation	Value	Unit	Biological Ref. Range
Glucose (Fasting)	95	mg/dL	60 - 110

Fasting Plasma Glucose (mg/dl)	2 hr plasma Glucose (mg/dl) Post Glucose load	Diagnosis
99 or below	139 or below	Normal
100 to 125	140 to 199	Pre-Diabetes (IGT)
126 or above	200 or above	Diabetes

Reference: American Diabetes Association.

Comment:

Impaired glucose tolerance (IGT) fasting, means a person has an increased risk of developing type 2 diabetes but does not

it yet. A level of 126 mg/dL or above, confirmed by repeating the test on another day, means a person has diabetes. IGT (2 hrs Post meal), means a person has an increased risk of developing type 2 diabetes but does not have it yet. A 2-hour glucose level of 200 mg/dL or above, confirmed by repeatig the test on another day, means a person has diabetes

Plasma Glucose Goals	For people with Diabetes
Before meal	70-130 mg/dL
2 Hours after meal	Less than 180 mg/dL
HbA1c	Less than 7%

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.

The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently.
 These reports are not valid for medico legal purposes.





# LAB DIVISION

Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Gender / Age Male / 30 Yrs

Refd. By

Client . Apollo Health & Lifestyle Ltd



Collected On	22/07/2023 10:06:37
Received On	22/07/2023 10:06:39
Released On	22/07/2023 17:27:38
Printed On	24/07/2023 18:03:42

Investigation	Value	Unit	Biological Ref. Range
Glucose, Post Prandial (PP)	131.0	mg/dL	70.0 - 140.0

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.
 The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently.
 These reports are not valid for medico legal purposes.

Page No: 3 of 11







Patient ID 1223240 Collected On 22/07/2023 10:06:37 Patient Name Mr. CHIRAG AMRUTBHAI GOHEL Received On 22/07/2023 10:06:39 Gender / Age Male / 30 Yrs Released On 22/07/2023 17:27:38 24/07/2023 18:03:43 Refd. By Printed On

Investigation	Value	Unit	Biological Ref. Range
Glycosylated Hb	5.6	%	
Average Plasma Glucose	114		

# Interpretation:

## HbA1c %

Client

<=5.6	Normal
5.7-6.4	At Risk for
	Diabetes
>=6.5	Diabetes

. Apollo Health & Lifestyle Ltd

Estimated Average Glucose (eAG) is a new way to understand how well you are managing your diabetes. Using eAG may help you get a better idea of how well you are taking care of your diabetes. And that can help you and your health care provider know what changes you may need to make to be as healthy as possible.

HbA1c %	5	5.5	6	6.5	7	7.5	8	8.5	9	10		12
I(A) I ma/ai	u /	111	126	140	154	169	183	197	212	240	269	298

The HbA1c goal for people with diabetes is less than 7 percent. A 3 to 6 monthly monitoring is recommended in diabetics. People with diabetes should get the test done more often if their blood sugar stays too high or if their healthcare provider makes any change in the treatment plan. HbA1c concentration represents the integrated values for blood glucose over the preceding 6 -10 wks and is not affected by daily glucose fluctuation, exercise & recent food intake. It is a more useful tool for clinical management of Diabetes mellitus through routine monitoring & assesses compliance with therapeutic regimen.

> Dr. Vishal Bhuva MBBS, MD Pathologist, HAM

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.

The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgentlyThese reports are not valid for medico legal purposes.







Patient ID 1223240 Collected On 22/07/2023 10:06:37 Patient Name Mr. CHIRAG AMRUTBHAI GOHEL Received On 22/07/2023 10:06:39 Gender / Age Male / 30 Yrs Released On 24/07/2023 18:01:01 24/07/2023 18:03:45 Refd. By Printed On Client . Apollo Health & Lifestyle Ltd

Investigation	Value	Unit	Biological Ref. Range
Blood group Gel Technique	"A" Positive		

Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB. The test is performed by both forward as well as reverse grouping methods.

The report is of sample received. It is presumed that the sample belongs to the patient. In case of any discrepancy related to this report, contact lab.

# COMPLETE BLOOD COUNT

Hemoglobin Cynmeth Photometric Measurement	14.5	gm/dL	13.0 - 17.0
Erythrocyte RBC Count	5.37	millions/cu.mm	4.50 - 5.50
Total Leukocyte Count (TLC) Electrical Impedance	12.2	X10^3/uL	4.0 - 11.0
Platelet Count Electrical Impedance	415	x10^3/uL	150 - 450
HCT Electrical Impedance	44.7	%	40.0 - 50.0
Mean Cell Volume (MCV)  Electrical Impedance	83.3	fL	80.0 - 100.0
Mean Cell Haemoglobin (MCH) Electrical Impedance	27.0	pg	27.0 - 32.0
Mean Corpuscular Hb Concn. (MCHC)	32.4	gm/dL	32.0 - 35.0
Red Cell Distribution Width (RDW-CV)	13.7	%	11.5 - 14.5
Differential Leukocyte Count (DLC)			
Neutrophils	61	%	40 - 80
Lymphocytes vcs	31	%	20 - 40
Eosinophils vcs	03	%	01 - 06
Monocytes vcs	05	%	02 - 08

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.

Page No: 5 of 11

The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgentlyThese reports are not valid for medico legal purposes.







Patient ID	1223240		Collected On	22/07/2023 10:06:37
Patient Name	Mr. CHIRAG AMRUTBHAI GOHEL		Received On	22/07/2023 10:06:39
Gender / Age	Male / 30 Yrs		Released On	24/07/2023 18:01:01
Refd. By			Printed On	24/07/2023 18:03:49
Client	. Apollo Health & Lifestyle Ltd	回道政治學院		

Investigation	Value	Unit	Biological Ref. Range
Basophils vcs	00	%	00 - 02
Erythrocyte Sedimentation Rate (ESR) Westergren's	30	mm in 1hr	00 - 10

- \* Test conducted on EDTA whole blood at 37 degree Celsius.
- \* ESR is an index of the presence of the active diseases of many types.
- \* Increased- in most infections, anaemias, injection of foreign proteins, auto-immune disorders, conditions accompanied by hyperglobunemia and hypercholesterolaemia.
- \* A rising ESR suggests a progressive disease.
- \* Decreased- in polycythemia, congestive heart failure.
- \* ESR is a useful but nonspecific marker of underlying inflammation. C-Reactive Protein(CRP) is the recommended test in a acute inflammatory conditions.

Dr. Vishal Bhuva MBBS, MD Pathologist, HAM

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.

The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently.
 These reports are not valid for medico legal purposes.

Page No: 6 of 11







Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Gender / Age Male / 30 Yrs

Refd. By

Client . Apollo Health & Lifestyle Ltd



Collected On 22/07/2023 10:06:37 Received On 22/07/2023 10:06:39 Released On 22/07/2023 17:27:38 24/07/2023 18:03:51 Printed On

Investigation	Value	Unit	Biological Ref. Range	
	Liver Function Test +	<u>GGT</u>		
Billirubin — Total Diazonium Salt	0.52	mg/dL	0.50 - 1.90	
Billirubin – Direct Diazo Reaction	0.23	mg/dL	0.00 - 0.50	
Bilirubin, Indirect	0.29	mg/dL	0.10 - 1.00	
Gultamic Oxaloacetic Transaminase (SGOT, AST)	17	U/L	10 - 37	
Gultamic Pyruvic Transaminase (SGPT, ALT)	20	U/L	0 - 41	
ALP (Alkaline Phosphatase)	106	U/L	40 - 150	
Total Protien Biuret method	6.7	g/dL	6.6 - 8.7	
Albumin Bromcresol Green	3.7	g/dL	3.5 - 5.2	
Globulin Calculated	3.0	g/dL	2.3 - 3.5	
A:G (Albumin:Globulin) Ratio	1.23		1.20 - 2.00	
Gamma Glutamyle Transpeptidas	34	U/L	0 - 55	

Page No: 7 of 11

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.
 The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently.
 These reports are not valid for medico legal purposes.







Patient ID 1223240 Collected On 22/07/2023 10:06:37 Patient Name Mr. CHIRAG AMRUTBHAI GOHEL Received On 22/07/2023 10:06:39 Gender / Age Male / 30 Yrs Released On 22/07/2023 17:27:38 24/07/2023 18:03:54 Refd. By Printed On Client

Investigation	Value	Unit	Biological Ref. Range
	Kidney Function Tes	<u>t</u>	
Urea, Serum	33	mg/dL	13 - 43
Creatinine Modified jaffe's	0.90	mg/dL	0.60 - 1.30
Uric Acid, Serum	4.50	mg/dL	3.50 - 7.20
Calcium Arsenazo III	10.00	mg/dl	8.40 - 10.20
Phosphorus uv photometric	3.32	mg/dL	2.60 - 4.50

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is awaste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. Inblood, it is a marker of GFR .in urine, it can remove the need for 24-hour collections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not workingproperly. As kidney disease progresses, the level of creatinine and urea in the blood increases. Certain drugs are nephrotoxichence KFT is done before and after initiation of treatment with these drugs.Low serum creatinine values are rare; they almost always reflect low muscle mass. Apart from renal failure Blood Urea can increase in dehydration and GI bleed.Reference ranges vary between laboratories.

Note: The result obtained relate only to the sample given/received & tested. A single test result is not always indicative of a disease, it has to be correlated with clinical data for interpretation.

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.

. Apollo Health & Lifestyle Ltd

The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently
 These reports are not valid for medico legal purposes.

Page No: 8 of 11





## LAB DIVISION

Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Gender / Age Male / 30 Yrs

Refd. By

Client . Apollo Health & Lifestyle Ltd



Collected On 22/07/2023 10:06:37 Received On 22/07/2023 10:06:39 22/07/2023 17:27:38 Released On 24/07/2023 18:03:57 Printed On

Investigation	Value	Unit	Biological Ref. Range
	<u>Lipid Profile</u>		
Cholesterol TOTAL	160	mg/dL	Desirable < 200 Borderline 200 - 239 High Risk >= 240
Triglycerides Glycerol Phosphate Oxidase	156	mg/dL	Normal <150 Borderline 150-199 High 200 -499 Very High >=500
DIRECT HDL Accelerator Selective Detergent	49	mg/dL	Major risk factor for heart disease < 40 Negative risk factor for heart disease =>60
VLDL Cholesterol	31	mg/dL	0 - 30
DIRECT LDL Calculated	80	mg/dL	Recommended <130 Moderate Risk 130-159 High Risk >160

In High Risk Patients including patient with Diabetes (Ref.NCEP III update 2017, the desirable LDL goal is < 100mg/dl.

Total / HDL Cholesterol Ratio	3.3		Low Risk 3.3-4.4 Average Risk 4.4-7.1 Moderate Risk 7.1-11.0 High Risk >11.0
Non HDL Cholesterol Calculated	111.0	mg/dL	Adult Optimal <130 Above Optimal 130 -159 Borderline High 160-189 High 190 -219 Very High >=220

Lipid profile is a panel of blood tests that serves as an initial screening tool for abnormalities in lipids, such as cholesterol and triglycerides. The results of this test can identify certain genetic diseases and can determine approximate risks for cardiovascular disease, certain forms of pancreatitis. Hypertriglyceridemia is indicative of insulin resistance when present with low high-density lipoprotein (HDL) and elevated low-density lipoprotein (LDL), while elevated triglyceride is a clinical risk factor for coronary artery disease (CAD), especially when low HDL is present. Very high levels of triglycerides are defined by serum levels of 500mg/dL or greater and can be concerning for development of pancreatitis. Reference ranges vary between laboratories.

> Dr. Vishal Bhuva MBBS, MD Pathologist, HAM

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.

The values are to be comborated with clinical finding and any alarming or unexpected result should be referred to this lab urgentlyThese reports are not valid for medico legal purposes.

Page No: 9 of 11







Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Gender / Age Male / 30 Yrs

Refd. By

Client . Apollo Health & Lifestyle Ltd



Collected On 22/07/2023 10:06:37 Received On 22/07/2023 10:06:39 Released On 24/07/2023 17:56:39 24/07/2023 18:04:04 Printed On

Investigatio	n		Value	Unit	Biological Ref. Range
			Thyroid Function T	est	
Triiodoth Chemilumineso	yronine (T3) cent Microparticle Immunoass	say (CMIA)	1.45	ng/dl	0.69 - 2.15
Thyroxine	e (T4) cent Microparticle Immunoass	say (CMIA)	116.00	ng/mL	52.00 - 127.00
Thyroid Stimulating Hormone (TSH) Chemiluminescent Microparticle Immunoassay (CMIA)		1.50	ulU/ml	0.45 - 5.60	
					Euthyroid 0.25 - 5.00 Hyperthyroid < 0.15 Hypothyroid > 7.00
TSH	Т3	T4	Suggested Interpretatio	n for the Thyroid Funct	ion Tests Pattern
Raised	Within range	Within range	Raised Within Range Within Range .Isolated High TSHespecially in the range of 4.7 to 15 m1U/m1 is commonly associated with Physiological & Biological TSH Variability. Subclinical Autoimmune Hypothyroidism.Intermittent 14 therapy for		

TSH	T3	T4	Suggested Interpretation for the Thyroid Function Tests Pattern
Raised	Within range	Within range	Raised Within Range Within Range .Isolated High TSHespecially in the range of 4.7 to 15 m1U/m1 is commonly associated with Physiological & Biological TSH Variability. Subclinical Autoimmune Hypothyroidism.Intermittent 14 therapy for hypothyroidism .Recovery phase after Non-Thyroidal illness"
Raised	Raised	Decreased	Chronic Autoimmune Thyroiditis Post thyroidectomy,Post radioiodine Hypothyroid phase of transient thyroiditis"
Raised or within range	Raised	Raised or within range	Interfering antibodies to thyroid hormones (anti-TPO antibodies)Intermittent 14 therapy or T4 overdose •Drug interference Amiodarone, Heparin,Beta blockers,steroids, anti-epileptics
Decreased	Raised or within range	Raised or within range	Isolated Low TSH -especially in the range of 0.1 to 0.4 often seen in elderly & Range Range associated with Non-Thyroidal illness .Subclinical Hyperthyroidism .Thyroxine ingestion'
Decreased	Decreased	Decreased	Central Hypothyroidism .Non-Thyroidal illness .Recent treatment for Hyperthyroidism (TSH remains suppressed)"
Decreased	Raised	Raised	Primary Hyperthyroidism (Graves' disease).Multinodular goitre, Toxic nodule •Transient thyroiditis:Postpartum, Silent (lymphocytic), Postviral (granulomatous,subacute, DeQuervain's),Gestational thyrotoxicosis with hyperemesis gravidarum"
Decreased Within Rang	Raised	Within range	T3 toxicosis •Non-Thyroidal illness
Within range	Decreased	Within range	Isolated Low T3-often seen in elderly & associated Non-Thyroidal illness In elderly the drop in 13 level can be upto 25%.

Dr. Vishal Bhuva MBBS, MD Pathologist, HAM

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.
 The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently.
 These reports are not valid for medico legal purposes.

Page No: 10 of 11







Patient ID 1223240

Patient Name Mr. CHIRAG AMRUTBHAI GOHEL

Male / 30 Yrs

Gender / Age

Refd. By

Client . Apollo Health & Lifestyle Ltd

Collected On 22/07/2023 10:06:37 Received On 22/07/2023 10:06:39 Released On 22/07/2023 17:27:38 24/07/2023 18:04:08 Printed On

Investigation	<b>V</b> alue	Unit	Biological Ref. Range
	Urine Examination (Rout	ine)	
Physical Examination			
Volume	30	mL	
Colour	PALE YELLOW		STRAW
Appearance	Clear		Turbid
pH	6.0		8.0
Specific Gravity	1.015		1.001-1.035
Chemical Examination			
Urine Protein	Nil		Nil
Urine Glucose	Nil		Negative
Ketone	Negative		Negative
Nitrite	Negative		Negative
Urobilinogen	Not Increased		Not Increased
Bilirubin	Nil		Nil
Microscopic Examination.			
Red Blood Cells	0-1	/hpf	Nil
Pus Cells (WBC)	1-2	/hpf	NIL
Epithelial Cells	2-3	/hpf	Nil
Casts	Nil	/hpf	
Crystals	Nil	·	Nil
Bacteria	Nil		Nil
Yeast	Nil		Nil
Mucous Threads	Nil		Nil
Trichomonas	Nil		Nil
Amorphous Material	Nil		Nil

\*\*\* End of Report \*\*\*

Dr. Vishal Bhuva MBBS, MD Pathologist, HAM

Note: 1. These report are mere estimation and are liable to vary / Change in different conditions in different laboratories.
 The values are to be corroborated with clinical finding and any alarming or unexpected result should be referred to this lab urgently.
 These reports are not valid for medico legal purposes.



# MER- MEDICAL EXAMINATION REPORT

Date of Examination	22/7/2023
NAME	Chivad Amoutbley Golil
AGE	Gender M
HEIGHT(cm)	169 WEIGHT (kg) 99.7
B.P.	118/ 72 M,
ECG	MSA
X Ray	N
Vision Checkup	Color Vision: Neo Coren such fel
	Far Vision Ratio: 6/6
	Near Vision Ratio:
Present Ailments	M.
Details of Past ailments (If Any)	
Comments / Advice : She /He is Physically Fit	Fil
BMI:34.9	

Signature with Stamp of Medical Examiner

Dr. Ninad J. Gor

Reg. No.: G-64033




# **CERTIFICATE OF MEDICAL FITNESS**

is is to certify that I have conducted the clinical examination	
Chiray (xohi) on 22/7/23	
(/ ter reviewing the medical history and on clinical examination it has been found that /she is	
	7
Medically Fit	,
• Fit with restrictions/recommendations	
Though following restrictions have been revealed, in my opinion, these are not impediments to the job.	
1	
2	
3	
However, the employee should follow the advice/medication that has been communicated to him/her.	
Review after	
Currently Unfit.	+
Review afterrecommended	
◆ Unfit	
	-
Dr	
Med <del>ical</del> Officer The Apollo Clinic, (Location)	
This certificate is not meant for medico-legal purposes	
Dr. Ninad J. Gor	<b>~</b>

M.B.B.S.

Reg. No.: G-64033







DATE:

# SWAMINARAYAN HOSPITAL

NCD CLINIC & COUNSELLING

Dr. Jagdish Dhanji Hala AGE/SEXS, 30 YEAB/ MALE DIABETOLOG

CENTER

REF BY: ROHA HELATHCARE

# **2D ECHO AND COLOUR DOPPLER STUDY**

FINAL IMPRESSION:

NAME: CHIRAG GOHEL

22.07.2023

- NORMAL LV SYSTOLIC FUNCTION WITH NORMAL LV SIZE.
- LVEF: 77.00 %, NO RWMA AT REST. TRIVIAL MR, TRIVIAL TR/AR, NO PR.
- NORMAL PAH, NORMAL RA/RV. NORMAL LA.
- NORMAL LV COMPLIANCE. LVH +
- NORMAL'RV FUNCTION.
- NO ASD, NO VSD, NO PDA. NO PE.

• IVC: NORMAL.

NOTE:

NO CHD, NO SHD, NO VHD

Reg.No.G 42676 IG 27-200855/ Sr.No.D-19188

ਗਮ ਗੇधामा माटे Appointment : 74074 98098

Education is Foundation For Prevention, Prevention is better than cure સ્વતઃસ્કુર્ણા : સ્વાદ આઝાદીનો - Swata:Sfurna : The taste of Freedom





# SWAMINARAYAN HOSPITAL

NCD CLINIC & COUNSELLING

**CENTER** 

Dr. Jagdish Dhanji Halai

NAME: CHIRAG GOHEL

MBBG, D. CARDIÓLOGY & DIABETOLOGY
AGE/SEX: 30 YEAR/FEMALE

DATE: 22.07.2023

**REF BY: ROHA HELATHCARE** 

# **2D ECHO AND COLOUR DOPPLER STUDY**

MITRAL VALVE

: NORMAL

**AORTIC VALVE** 

: NORMAL.

**PULMONARY VALVE** 

: NORMAL.

TRICUSPID VALVE

: NORMAL.

**AORTA** 

: ROOT: 18.00 MM AND AORTA ST JUNCTION: 22.00 MM.

NORMAL ARCH AND WHOLE AORTA FROM ORIGINE TO

BIFURCATION.

LA

: 33 MM

LV- D/LV-S

: 44/29 MM.

LVEF

: 77.00 %, RWMA PRESENT AT REST.

IVS

: INTACT, IVS: 11.30 MM.

IAS

: INTACT, PW: 11.30 MM.

AOVP

: 1.68 M/SEC. PVP: 0.82 M/SEC.

RA AND RV

: NORMAL, PA: NORMAL.

**RVSP** 

: TR JET + RA MEAN PRESSURE: 31 MM HG TAPSE: 21.60 MM

COLOR DOPPLER STUDY

: TRIVIAL MR, TRIVAL TR, PR: NO, TRIVIAL AR.

NO AS, NO MS, NO TS, NO PS.

ALL VLVES ARE NORMAL WITH NORMAL HEMODYNAMIC.

**MVIS** 

: VE/VA > 1.


NO PERICARDIAL EFFUSION.

NO VSR, NO SCAR, NO CLOT, NO VEGETATION.

NO THROMBUS IN LV/LVA.

ਗਮ ਗੇधामा माटे Appointment : 74074 98098

Education is Foundation For Prevention, Prevention is better than cure સ્વતઃસ્કુર્ણા : સ્વાદ આઝાદીનો - Swata:Sfurna : The taste of Freedom





(A Division of KRIC Radio Diagnostic Pvt. Ltd.)

• Email : kric2008@gmail.com • Website : www.kric.in

# Dr. Kripalsinh Jadeja

Dr. Bhaven Shah

M.D.

Consultant Radiologist

M.B., D.M.R.E.

Consultant Radiologist Name: CHIRAG GOHIL

MR No: D91126 Modality: US Gender: M Age: 30YY Date: 22/07/2023

Referred By : ROHA HEALTH CARE

# **USG: ABDOMEN & PELVIS**

**LIVER:** appears enlarged in size and measures 17 cm and show raised parenchymal echotexture. No evidence of focal or diffuse lesion. No evidence of dilated IHBR. PV and CBD appear normal in calibre.

GALL BLADDER: appears normal. No intrinsic lesion seen.

**PANCREAS:** appears normal in size and echotexture. No focal mass lesion or changes of pancreatitis seen.

SPLEEN: appears normal in size and echotexture. No evidence of focal or diffuse lesion.

**BOTH KIDNEYS:** appear normal in size and echotexture with preservation of corticomedullary differentiation. No evidence of calculus, hydronephrosis or mass lesion involving either kidney.

RK: 10.7 x 5.3 cm LK: 10.8 x 5.9 cm

**URINARY BLADDER:** appears normal. No intrinsic lesion seen.

PROSTATE: Appears normal in size.

No e/o Ascites or paraaortic lymphadenopathy seen.

# **CONCLUSION:**

- \* Moderate changes of fatty liver Grade II with Mild hepatomegaly.
- \* NORMAL SONOGRAPHY OF GB, SPLEEN, PANCREAS, BOTH KIDNEYS, U.BLADDER & PROSTATE.


ADV: Clinical correlation and further investigation. Thanks for ref...

Dr.KRIPALSINH JADEJA M.B,D.M.R.E RADIOLOGIST

**KRICBHUJ** 

1.5 TESLA 196 CHANNEL MRI | 16 SLICE MDCT SCAN | 3D & 4D USG | COLOUR DOPPLER | DIGITAL X-RAY | MAMMOGRAPHY | CBCT | OPG







Patient Name: MR CHIRAG AMRUT BHAI GOHEL

MR No: 220703 Modality: DX Gender: M Age: 30YY

Date: 22/07/2023

# X RAY CHEST (PA)

Both the lung fields do not reveal any parenchymal abnormality except minimal bilateral apical pleural thickening noted.

Both CP angles are clear.

Cardiac size is within normal limits.

Both domes of the diaphragm appear normal.

Bony thoracic cage appears normal.

# **CONCLUSION:**

NO SIGNIFICANT ABNORMALITY DETECTED.

ADV: Clinical correlation and further investigation. Thanks for ref...

Dr.KRIPALSINH JADEJA M.B,D.M.R.E RADIOLOGIST