



Lab No. : KLP/11-03-2023/SR7392127
 Patient Name : AVISHEK DASGUPTA
 Age : 35 Y 1 M 12 D
 Gender : M

Lab Add. : Newtown, Kolkata-700156
 Ref Dr. : Dr.MEDICAL OFFICER
 Collection Date: 11/Mar/2023 09:29AM
 Report Date : 11/Mar/2023 05:08PM



Test Name	Result	Unit	Bio Ref. Interval	Method
-----------	--------	------	-------------------	--------

[PDF Attached](#)

GLYCATED HAEMOGLOBIN (HBA1C) , EDTA WHOLE BLOOD

GLYCATED HEMOGLOBIN (HBA1C)	5.1	%	***FOR BIOLOGICAL REFERENCE INTERVAL DETAILS , PLEASE REFER TO THE BELOW MENTIONED REMARKS/NOTE WITH ADDITIONAL CLINICAL INFORMATION ***	
HbA1c (IFCC)	32.0	mmol/mol		HPLC

Clinical Information and Laboratory clinical interpretation on Biological Reference Interval:

Low risk / Normal / non-diabetic : <5.7% (NGSP) / < 39 mmol/mol (IFCC)
 Pre-diabetes/High risk of Diabetes : 5.7%- 6.4% (NGSP) / 39 - < 48 mmol/mol (IFCC)
 Diabetics-HbA1c level : >= 6.5% (NGSP) / > 48 mmol/mol (IFCC)

Analyzer used : Bio-Rad-VARIANT TURBO 2.0
 Method : HPLC Cation Exchange

Recommendations for glycemic targets

- Ø Patients should use self-monitoring of blood glucose (SMBG) and HbA1c levels to assess glycemic control.
- Ø The timing and frequency of SMBG should be tailored based on patients' individual treatment, needs, and goals.
- Ø Patients should undergo HbA1c testing at least twice a year if they are meeting treatment goals and have stable glycemic control.
- Ø If a patient changes treatment plans or does not meet his or her glycemic goals, HbA1c testing should be done quarterly.
- Ø For most adults who are not pregnant, HbA1c levels should be <7% to help reduce microvascular complications and macrovascular disease . Action suggested >8% as it indicates poor control.
- Ø Some patients may benefit from HbA1c goals that are stringent.

Result alterations in the estimation has been established in many circumstances, such as after acute/ chronic blood loss, for example, after surgery, blood transfusions, hemolytic anemia, or high erythrocyte turnover; vitamin B₁₂/ folate deficiency, presence of chronic renal or liver disease; after administration of high-dose vitamin E / C; or erythropoietin treatment.

Reference: Glycated hemoglobin monitoring BMJ 2006; 333:586-8

References:

1. Chamberlain JJ, Rhinehart AS, Shaefer CF, et al. Diagnosis and management of diabetes: synopsis of the 2016 American Diabetes Association Standards of Medical Care in Diabetes. Ann Intern Med. Published online 1 March 2016. doi:10.7326/M15-3016.
2. Mosca A, Goodall I, Hoshino T, Jeppsson JO, John WG, Little RR, Miedema K, Myers GL, Reinauer H, Sacks DB, Weykamp CW. International Federation of Clinical Chemistry and Laboratory Medicine, IFCC Scientific Division. Global standardization of glycated hemoglobin measurement: the position of the IFCC Working Group. Clin Chem Lab Med. 2007;45(8):1077-1080.

Dr NEEPA CHOWDHURY
 MBBS MD (Biochemistry)
 Consultant Biochemist

Lab No. : SR7392127 Name : AVISHEK DASGUPTA Age/G : 35 Y 1 M 12 D / M Date : 11-03-2023

SODIUM, BLOOD , GEL SERUM					
SODIUM,BLOOD	141.00	mEq/L	132 - 146 mEq/L	ISE INDIRECT	
SGOT/AST , GEL SERUM					
SGOT/AST	36.00	U/L	13-40 U/L	Modified IFCC	
BILIRUBIN (DIRECT) , GEL SERUM					
BILIRUBIN (DIRECT)	0.20	mg/dL	<0.2 mg/dL	Vanadate oxidation	
URIC ACID, BLOOD , GEL SERUM					
URIC ACID,BLOOD	8.30	mg/dL	3.5-7.2 mg/dL	Uricase/Peroxidase	
ALKALINE PHOSPHATASE , GEL SERUM					
ALKALINE PHOSPHATASE	99.00	U/L	46-116 U/L	IFCC standardization	
BILIRUBIN (TOTAL) , GEL SERUM					
BILIRUBIN (TOTAL)	0.80	mg/dL	0.3-1.2 mg/dL	Vanadate oxidation	
POTASSIUM, BLOOD , GEL SERUM					
POTASSIUM,BLOOD	4.60	mEq/L	3.5-5.5 mEq/L	ISE INDIRECT	
UREA,BLOOD , GEL SERUM					
UREA,BLOOD	30.0	mg/dL	19-49 mg/dL	Urease with GLDH	
CREATININE, BLOOD					
CREATININE, BLOOD	1.06	mg/dL	0.7-1.3 mg/dL	Jaffe, alkaline picrate, kinetic	
GLUCOSE, FASTING , BLOOD, NAF PLASMA					
GLUCOSE,FASTING	85	mg/dL	Impaired Fasting-100-125 .-Diabetes- >= 126.-Fasting is defined as no caloric intake for at least 8 hours.	Gluc Oxidase Trinder	

In the absence of unequivocal hyperglycemia, diagnosis requires two abnormal test results from the same sample or in two separate test samples.

Reference :
ADA Standards of Medical Care in Diabetes – 2020. *Diabetes Care* Volume 43, Supplement 1.

THYROID PANEL (T3, T4, TSH) , GEL SERUM					
T3-TOTAL (TRI IODOTHYRONINE)	0.86	ng/ml	0.60-1.81 ng/ml	CLIA	
T4-TOTAL (THYROXINE)	5.4	µg/dL	3.2-12.6 µg/dL	CLIA	
TSH (THYROID STIMULATING HORMONE)	2.63	µIU/mL	0.55-4.78 µIU/mL	CLIA	

Serum TSH levels exhibit a diurnal variation with the peak occurring during the night and the nadir, which approximates to 50% of the peak value, occurring between 1000 and 1600 hours.[1,2]

References:

- Bugalho MJ, Domingues RS, Pinto AC, Garrao A, Catarino AL, Ferreira T, Limbert E and Sobrinho L. Detection of thyroglobulin mRNA transcripts in peripheral blood of individuals with and without thyroid glands: evidence for thyroglobulin expression by blood cells. *Eur J Endocrinol* 2001;145:409-13.
- Bellantone R, Lombardi CP, Bossola M, Ferrante A, Princi P, Boscherini M et al. Validity of thyroglobulin mRNA assay in peripheral blood of postoperative thyroid carcinoma patients in predicting tumor recurrence varies according to the histologic type: results of a prospective study. *Cancer* 2001;92:2273-9.

BIOLOGICAL REFERENCE INTERVAL: [ONLY FOR PREGNANT MOTHERS]

Trimester specific TSH LEVELS during pregnancy:

FIRST TRIMESTER: 0.10 – 3.00 μ IU/mL

SECOND TRIMESTER: 0.20 -3.50 μ IU/mL

THIRD TRIMESTER : 0.30 -3.50 μ IU/mL

References:

1. Erik K. Alexander, Elizabeth N. Pearce, Gregory A. Brent, Rosalind S. Brown, Herbert Chen, Chrysoula Dosiou, William A. Grobman, Peter Laurberg, John H. Lazarus, Susan J. Mandel, Robin P. Peeters, and Scott Sullivan. *Thyroid*. Mar 2017. 315-389. <http://doi.org/10.1089/thy.2016.0457>

2. Kalra S, Agarwal S, Aggarwal R, Ranabir S. Trimester-specific thyroid-stimulating hormone: An indian perspective. *Indian J Endocr Metab* 2018;22:1-4.

PHOSPHORUS-INORGANIC, BLOOD , GEL SERUM

PHOSPHORUS-INORGANIC,BLOOD	3.1	mg/dL	2.4-5.1 mg/dL	Phosphomolybdate/UV
----------------------------	-----	-------	---------------	---------------------

***CHLORIDE, BLOOD , .**

CHLORIDE,BLOOD	105.00	mEq/L	99-109 mEq/L	ISE INDIRECT
----------------	--------	-------	--------------	--------------

□



Dr NEEPA CHOWDHURY
MBBS MD (Biochemistry)
Consultant Biochemist



Lab No. : SR7392127 Name : AVISHEK DASGUPTA Age/G : 35 Y 1 M 12 D / M Date : 11-03-2023

SGPT/ALT , GEL SERUM

SGPT/ALT **66.00** U/L 7-40 U/L Modified IFCC

CALCIUM, BLOOD

CALCIUM,BLOOD 9.80 mg/dL 8.7-10.4 mg/dL Arsenazo III


TOTAL PROTEIN [BLOOD] ALB:GLO RATIO , .

TOTAL PROTEIN 7.20 g/dL 5.7-8.2 g/dL BIURET METHOD
 ALBUMIN 4.7 g/dL 3.2-4.8 g/dL BCG Dye Binding
 GLOBULIN 2.50 g/dl 1.8-3.2 g/dl Calculated
 AG Ratio 1.88 1.0 - 2.5 Calculated

LIPID PROFILE , GEL SERUM

CHOLESTEROL-TOTAL 151.00 mg/dL Desirable: < 200 mg/dL
 Borderline high: 200-239 mg/dL
 High: > or =240 mg/dL Enzymatic
 TRIGLYCERIDES 82.00 mg/dL Normal: < 150,
 BorderlineHigh::150-199,
 High:: 200-499,
 VeryHigh:: >500 GPO-Trinder
 HDL CHOLESTEROL **28.00** mg/dl < 40 - Low
 40-59- Optimum Elimination/catalase
 60 - High
 LDL CHOLESTEROL DIRECT **116.0** mg/dL OPTIMAL : <100 mg/dL,
 Near optimal/ above optimal :
 100-129 mg/dL,
 Borderline high : 130-159 mg/dL,
 High : 160-189 mg/dL,
 Very high : >=190 mg/dL Elimination / Catalase
 VLDL 7 mg/dl < 40 mg/dl Calculated
 CHOL HDL Ratio 5.4 LOW RISK 3.3-4.4 AVERAGE
 RISK 4.47-7.1 MODERATE RISK
 7.1-11.0 HIGH RISK >11.0 Calculated

Reference: National Cholesterol Education Program. Executive summary of the third report of The National Cholesterol Education Program (NCEP) Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults (Adult Treatment Panel III). JAMA. May 16 2001;285(19):2486-97.


Dr. SUPARBA CHAKRABARTI
 MBBS, MD(BIOCHEMISTRY)
 Consultant Biochemist



Lab No. : SR7392127 Name : AVISHEK DASGUPTA Age/G : 35 Y 1 M 12 D / M Date : 11-03-2023

URINE ROUTINE ALL, ALL , URINE

PHYSICAL EXAMINATION

COLOUR PALE YELLOW
 APPEARANCE SLIGHTLY HAZY

CHEMICAL EXAMINATION

pH	6.0	4.6 - 8.0	Dipstick (triple indicator method)
SPECIFIC GRAVITY	1.020	1.005 - 1.030	Dipstick (ion concentration method)
PROTEIN	NOT DETECTED	NOT DETECTED	Dipstick (protein error of pH indicators)/Manual
GLUCOSE	NOT DETECTED	NOT DETECTED	Dipstick (glucose-oxidase-peroxidase method)/Manual
KETONES (ACETOACETIC ACID, ACETONE)	NOT DETECTED	NOT DETECTED	Dipstick (Legals test)/Manual
BLOOD	NOT DETECTED	NOT DETECTED	Dipstick (pseudoperoxidase reaction)
BILIRUBIN	NEGATIVE	NEGATIVE	Dipstick (azo-diazo reaction)/Manual
UROBILINOGEN	NEGATIVE	NEGATIVE	Dipstick (diazonium ion reaction)/Manual
NITRITE	NEGATIVE	NEGATIVE	Dipstick (Griess test)
LEUCOCYTE ESTERASE	NEGATIVE	NEGATIVE	Dipstick (ester hydrolysis reaction)

MICROSCOPIC EXAMINATION

LEUKOCYTES (PUS CELLS)	0-1	/hpf	0-5	Microscopy
EPITHELIAL CELLS	1-2	/hpf	0-5	Microscopy
RED BLOOD CELLS	NOT DETECTED	/hpf	0-2	Microscopy
CAST	NOT DETECTED		NOT DETECTED	Microscopy
CRYSTALS	NOT DETECTED		NOT DETECTED	Microscopy
BACTERIA	NOT DETECTED		NOT DETECTED	Microscopy
YEAST	NOT DETECTED		NOT DETECTED	Microscopy

Note:

- All urine samples are checked for adequacy and suitability before examination.
- Analysis by urine analyzer of dipstick is based on reflectance photometry principle. Abnormal results of chemical examinations are confirmed by manual methods.
- The first voided morning clean-catch midstream urine sample is the specimen of choice for chemical and microscopic analysis.
- Negative nitrite test does not exclude urinary tract infections.
- Trace proteinuria can be seen in many physiological conditions like exercise, pregnancy, prolonged recumbency etc.
- False positive results for glucose, protein, nitrite, urobilinogen, bilirubin can occur due to use of certain drugs, therapeutic dyes, ascorbic acid, cleaning agents used in urine collection container.
- Discrepancy between results of leukocyte esterase and blood obtained by chemical methods with corresponding pus cell and red blood cell count by microscopy can occur due to cell lysis.
- Contamination from perineum and vaginal discharge should be avoided during collection, which may falsely elevate epithelial cell count and show presence of bacteria and/or yeast in the urine.

CBC WITH PLATELET (THROMBOCYTE) COUNT , EDTA WHOLE BLOOD

HEMOGLOBIN	15.2	g/dL	13 - 17	PHOTOMETRIC
WBC	7.0	*10 ³ /μL	4 - 10	DC detection method
RBC	5.00	*10 ⁶ /μL	4.5 - 5.5	DC detection method
PLATELET (THROMBOCYTE) COUNT	180	*10 ³ /μL	150 - 450*10 ³ /μL	DC detection method/Microscopy

DIFFERENTIAL COUNT

NEUTROPHILS	56	%	40 - 80 %	Flowcytometry/Microscopy
LYMPHOCYTES	34	%	20 - 40 %	Flowcytometry/Microscopy
MONOCYTES	07	%	2 - 10 %	Flowcytometry/Microscopy
EOSINOPHILS	03	%	1 - 6 %	Flowcytometry/Microscopy
BASOPHILS	00	%	0-0.9%	Flowcytometry/Microscopy

CBC SUBGROUP

Lab No. : KLP/11-03-2023/SR7392127

Page 5 of 8



Lab No. : SR7392127	Name : AVISHEK DASGUPTA	Age/G : 35 Y 1 M 12 D / M	Date : 11-03-2023	
HEMATOCRIT / PCV	45.7	%	40 - 50 %	Calculated
MCV	91.4	fl	83 - 101 fl	Calculated
MCH	30.5	pg	27 - 32 pg	Calculated
MCHC	33.4	gm/dl	31.5-34.5 gm/dl	Calculated
RDW - RED CELL DISTRIBUTION WIDTH	14.9	%	11.6-14%	Calculated
PDW-PLATELET DISTRIBUTION WIDTH	32.0	fL	8.3 - 25 fL	Calculated
MPV-MEAN PLATELET VOLUME	13.8		7.5 - 11.5 fl	Calculated

ESR (ERYTHROCYTE SEDIMENTATION RATE) , EDTA WHOLE BLOOD

1stHour	10	mm/hr	0.00 - 20.00 mm/hr	Westergren
---------	----	-------	--------------------	------------

BLOOD GROUP ABO+RH [GEL METHOD] , EDTA WHOLE BLOOD

ABO	O			Gel Card
RH	POSITIVE			Gel Card

TECHNOLOGY USED: GEL METHOD

ADVANTAGES :

- Gel card allows simultaneous forward and reverse grouping.
- Card is scanned and record is preserved for future reference.
- Allows identification of Bombay blood group.
- Daily quality controls are run allowing accurate monitoring.

Historical records check not performed.

DR. NEHA GUPTA
MD, DNB (Pathology)
Consultant Pathologist

Lab No. : SR7392127 Name : AVISHEK DASGUPTA Age/G : 35 Y 1 M 12 D / M Date : 12-03-2023

URIC ACID, URINE, SPOT URINE

URIC ACID, SPOT URINE 50.00 mg/dL 37-92 mg/dL URICASE

□



DR. ANANNYA GHOSH
MBBS, MD (Biochemistry)
Consultant Biochemist

Lab No. : KLP/11-03-2023/SR7392127
Patient Name : AVISHEK DASGUPTA
Age : 35 Y 1 M 12 D
Gender : M

Lab Add. :
Ref Dr. : Dr.MEDICAL OFFICER
Collection Date:
Report Date : 11/Mar/2023 02:43PM




X-RAY REPORT OF CHEST (PA)

FINDINGS :

No active lung parenchymal lesion is seen.
Both the hila are normal in size, density and position.
Mediastinum is in central position. Trachea is in midline.
Domes of diaphragm are smoothly outlined. Position is within normal limits.
Lateral costo-phrenic angles are clear.
The cardio-thoracic ratio is normal.
Bony thorax reveals no definite abnormality.

IMPRESSION :

Normal study.


Dr. Anoop Sastry
MBBS, DMRT(CAL)
CONSULTANT RADIOLOGIST
Registration No.: WB-36628

Patient Data

Sample ID: C02135026473
 Patient ID: SR7392127
 Name:
 Physician:
 Sex:
 DOB:

Analysis Data

Analysis Performed: 11/MAR/2023 16:45:20
 Injection Number: 5589U
 Run Number: 131
 Rack ID: 0005
 Tube Number: 9
 Report Generated: 11/MAR/2023 16:51:42
 Operator ID: ASIT

Comments:

Peak Name	NGSP %	Area %	Retention Time (min)	Peak Area
A1a	---	1.2	0.157	22177
A1b	---	0.9	0.220	16353
F	---	0.7	0.275	13214
LA1c	---	1.6	0.403	30359
A1c	5.1	---	0.510	78528
P3	---	3.4	0.791	64482
P4	---	1.2	0.869	22735
Ao	---	87.1	0.989	1671941

Total Area: 1,919,789

HbA1c (NGSP) = 5.1 % HbA1c (IFCC) = 32 mmol/mol

