

9438 8795 9944

मेरा आधार, मेरी पहचान

Dr. GOYAL'S Path Lab & Imaging Center, Jaipus

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Tele: 0141-2293346, 4049787, 9887049787





Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID: -122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :- EDTA

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 14:57:50

#### HAEMATOLOGY

Test Name Value Unit Biological Ref Interval

**BOB PACKAGEFEMALE BELOW 40** 

GLYCOSYLATED HEMOGLOBIN (HbA1C)

5.9

%

Non-diabetic: < 5.7 Pre-diabetics: 5.7-6.4 Diabetics: = 6.5 or higher ADA Target: 7.0 Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose overthe period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasmaglucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHbdepends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to themean of HbA1C.Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1cmeasurements. The effects vary depending on the specific Hb vatiant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE Method:- Calculated Parameter

mg/dL

Non Diabetic < 100 mg/dL Prediabetic 100- 125 Diabetic 126 mg/dL or

**AJAYSINGH Technologist** 

Page No: 1 of 15



DR.TANURUNGTA M.D (Path) RMC No.-17226

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





:- 12/03/2022 11:08:18 Date

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company:- MediWheel

Sample Type :- EDTA

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 14:57:50

HAEMA	OTA	LOGY	ľ
-------	-----	------	---

Test Name	est Name Value Unit		Biological Ref Interva
HAEMOGARAM			Ŷ
HAEMOGLOBIN (Hb)	11.8 L	g/dL	12.0 - 15.0
TOTAL LEUCOCYTE COUNT	9.25	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	56.3	%	40.0 - 80.0
LYMPHOCYTE	37.0	%	20,0 - 40,0
EOSINOPHIL	3.4	%	1.0 - 6.0
MONOCYTE	3.0	%	2.0 - 10.0
BASOPHIL	0.3	%	0.0 - 2.0
NEUT#	5.21	10^3/uL	1.50 - 7.00
LYMPH#	3.43	10^3/uL	1.00 - 3.70
EO#	0.31	10^3/uL	0.00 - 0.40
MONO#	0.27	10^3/uL	0.00 - 0.70
BASO#	0.03	10^3/uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.73	x10^6/uL	3.80 - 4.80
HEMATOCRIT (HCT)	36.40	%	36.00 - 46.00
MEAN CORP VOLUME (MCV)	77.0 L	fL	83,0 - 101,0
MEAN CORP HB (MCH)	25.0 L	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	32.5	g/dL	31.5 - 34.5
PLATELET COUNT	267	x10^3/uL	150 - 410
RDW-CV	14.0	%	11.6 - 14.0
MENTZER INDEX	16.28		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

**AJAYSINGH Technologist** 

Page No: 2 of 15



DR.TANURUNGTA M.D (Path) RMC No.-17226

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Sample Type :- EDTA

Patient ID: -122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 14:57:50

Biological Ref Interval

#### HAEMATOLOGY

**Test Name** 

Value

Unit

Erythrocyte Sedimentation Rate (ESR)

24 H

mm/hr.

00 - 20

(ESR) Methodology: Measurment of ESR by cells aggregation.

Instrument Name : Indepedent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator ofinflammatory disease and abnormal protein states.

The test in used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnency due to hyperfibrinogenaemia.

The "3-figure ESR " x>100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia (FBC) the the disease of the content of the the disease of the content of

AJAYSINGH **Technologist** 

Page No: 3 of 15



Jany DR.TANURUNGTA M.D (Path) RMC No.-17226

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date :- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel
Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 16:18:06

### BIOCHEMISTRY

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp :-

	BIOCHEMISTRY			
Test Name	Value	Unit	Biological Ref Interval	
LIPID PROFILE				
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	164.78	mg/dl	Desirable <200 Borderline 200-239 High> 240	
TRIGLYCERIDES Method:- GPO-PAP	121.33	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500	
VLDL CHOLESTEROL Method:- Calculated	24.27	mg/dl	0.00 - 80.00	

MUKESHSINGH

Page No: 4 of 15



DR.TANURUNGTA
M.D (Path) RMC No.-17226

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Patient ID :-122127659 Ref. By Dr:- BOB

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Lab/Hosp :-

Company :- MediWheel Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 16:18:06

### BIOCHEMISTRY

BIOCHEMISTRY					
Test Name	Value	Unit	Biological Ref Interval		
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	37.23	mg/dl	Low < 40 High > 60		
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	107.33	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190		
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	4.43		0.00 - 4.90		
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	2.88	5:	0.00 - 3.50		
TOTAL LIPID Method:- CALCULATED	512.82	mg/dl	400.00 - 1000.00		

TOTAL CHOLESTEROL InstrumentName: Randox Rx Imola Interpretation; Cholesterol measurements are used in the diagnosis and treatments of lipid lipop

TRIGLYCERIDES InstrumentName: Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid measurements.

DIRECT HDLCHOLESTERO InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serion and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing pattern risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROL Instrument Name: Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on high reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.

TOTAL LIPID AND VLDL ARE CALCULATED

MUKESHSINGH

Page No: 5 of 15



DR.TANURUNGTA M.D (Path) RMC No.-17226

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 16:18:06

BI	О	С	Н	Е	M	II	S	Г	R	Y

	DIOCHEN	ISIKI	
Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.66	mg/dl	Up to - 1.0 Cord blood <2 mg/dL Premature < 6 days <16mg/dL Full-term < 6 days= 12 mg/dL 1month - <12 months <2 mg/dL 1-19 years <1.5 mg/dL Adult - Up to - 1.2 Ref-(ACCP 2020)
SGOT Method:- IFCC	20.8	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	17.6	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:-AMP Buffer	51.30	IU/L	30.00 - 120.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	8.28	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.51	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	3.77 H	gm/dl	2.20 - 3.50
A/G RATIO	1.20 └		1.30 - 2.50

MUKESHSINGH

Page No: 6 of 15



### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Sample Type :- PLAIN/SERUM

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date NAME :- Mrs. NEHA TAILOR

:- 12/03/2022 11:08:18

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 16:18:06

#### RIOCHEMISTRY

	DIOCHEMISTRI				
Test Name	Value	Unit	Biological Ref Interval		
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.31	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL		
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.35	mg/dl	0.30-0.70		
SERUM GAMMA GT Method:- IFCC	8.00	U/L	7.00 - 32.00		

Total BilirubinMethodology: Colorimetric method InstrumentName:Randox Rx Imola Interpretation An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC InstrumentName:Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric nuccess, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCCInstrumentName:Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer InstrumentName:Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobilary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Biuret Reagent InstrumentName:Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green InstrumentName:Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous disprimarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola Interpretation: Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra-or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

MUKESHSINGH

Page No: 7 of 15



### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel Sample Type :- PLAIN/SERUM Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 13:55:02

**IMMUNOASSAY** 

**Test Name** 

Value

Biological Ref Interval

TOTAL THYROID PROFILE

SERUM TSH Method:- Enhanced Chemiluminescence Immunoassay

2.200

μIU/mL

Unit

0.465 - 4.680

**ANANDSHARMA Technologist** 

Page No: 8 of 15



DR.TANURUNGTA M.D (Path) RMC No.-17226

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Ref. By Dr:- BOB

Patient ID: -122127659

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Lab/Hosp :-

Company :- MediWheel
Sample Type :- PLAIN/SERUM

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 13:55:02

#### **IMMUNOASSAY**

Test Name	est Name Value		Biological Ref Interval
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.250	ng/ml	0.970 - 1.690
SERUM TOTAL T4	8.120	ug/dl	5.500 - 11.000

Method:-Chemiluminescence(Competitive immunoassay)
InstrumentName: VITROS ECI Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

InstrumentName: VITROS ECI Interpretation: The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT41) and estimate the concentration of free T4.Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

InstrumentName: VITROS ECI Interpretation: TSH stimulates the production of thyroxine (T4) and triodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

ANANDSHARMA Technologist

Page No: 9 of 15



Janu

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sample Type :- URINE

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 15:58:13

#### **CLINICAL PATHOLOGY**

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	3-5	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT	51	ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		

SAPNA **Technologist** 

Page No: 10 of 15



### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :- URINE

Sample Collected Time 12/03/2022 11:13:26

Final Authentication: 12/03/2022 15:58:13

#### **CLINICAL PATHOLOGY**

Test Name	Value Unit	Biological Ref Interval
PHYSICAL EXAMINATION		
COLOUR	PALE YELLOW	PALE YELLOW
APPEARANCE	Clear	Clear
CHEMICAL EXAMINATION		
REACTION(PH)	6.0	5.0 - 7.5
SPECIFIC GRAVITY	1.005	1,010 - 1,030
PROTEIN	NIL	NIL
SUGAR	NIL	NIL
BILIRUBIN	NEGATIVE	NEGATIVE
UROBILINOGEN	NORMAL	NORMAL
KETONES	NEGATIVE	NEGATIVE
NITRITE	NEGATIVE	NEGATIVE

SAPNA Technologist

Page No: 11 of 15



Janu

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel Sample Type :- STOOL

Sample Collected Time 12/03/2022 14:30:34

Patient ID :-122127659 Ref. By Dr:- BOB

Lab/Hosp :-

Final Authentication: 12/03/2022 15:55:55

CLINICAL PATHOLOGY

**Test Name** 

Value

Unit

Biological Ref Interval

STOOL ANALYSIS

PHYSICAL EXAMINATION

MUCUS

BLOOD

MICROSCOPIC EXAMINATION

RBC's

WBC/HPF

OVA

**CYSTS** 

OTHERS Collected Sample Received

/HPF

/HPF

SAPNA **Technologist** 

Page No: 12 of 15



DR.TANURUNGTA M.D (Path) RMC No.-17226

### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID :-122127659

Ref. By Dr:- BOB

Lab/Hosp:-

Sample Type :- KOx/Na FLUORIDE-F, KOx/Na Sabbor IOE FOR PRIMARY SERSION 22 14:30:50 Final Authentication: 12/03/2022 16:18:06

BIOCHEMISTRY

**Test Name** Value Unit Biological Ref Interval

FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP

91.2

mg/dl

Impaired glucose tolerance (IGT) 111 - 125 mg/dL Diabetes Mellitus (DM) > 126 mg/dL

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, panereatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases

BLOOD SUGAR PP (Plasma) Method:- GOD PAP

75.0 - 115.0

Instrument Name: Randox Rx Imola Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancrentic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels(hypoglycemia) may result from excessive insulin therapy or various liver diseases .

SERUM CREATININE Method:- Colorimetric Method

SERUM URIC ACID Method:- Enzymatic colorimetric

0.74

5.02

94.7

mg/dl

mg/dl

Men - 0.6-1.30 Women - 0,5-1.20

Men - 3.4-7.0 Women - 2.4-5.7

MUKESHSINGH

Page No: 13 of 15



DR.TANURUNGTA M.D (Path) RMC No.-17226

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019

Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date

:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID: -122127659

Ref. By Dr:- BOB

Lab/Hosp :-

### **HAEMATOLOGY**

**Test Name** 

Value

Unit

Biological Ref Interval

AJAYSINGH, ANANDSHARMA, BILAL, KOMAL, MUKESHSINGH, SAPNA

Page No: 14 of 15



### Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur-302019 Tele: 0141-2293346, 4049787, 9887049787 Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



:- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID: -122127659

Ref. By Dr:- BOB

Lab/Hosp :-

Sample Type :- EDTA, PLAIN/SERUM, URINE, SERNINE-PBIlected Time 12/03/2022 14:30:34

Final Authentication: 12/03/2022 16:18:06

#### HAEMATOLOGY

Test Name

Value

Unit

Biological Ref Interval

**BLOOD GROUP ABO** 

"O"POSITIVE

BLOOD GROUP ABO Methodology: Haemagglutination reaction Kit Name: Monoclonal agglutinating antibodies (Span clone)

URINE SUGAR (FASTING)
Collected Sample Received

Nil

Nil

URINE SUGAR PP Collected Sample Received

Nil

Nil

BLOOD UREA NITROGEN (BUN)

11.0

mg/dl

0.0 - 23.0

\*\*\* End of Report \*\*\*

AJAYSINGH, MUKESHSINGH, SAPNA Technologist

Page No: 15 of 15



Dr. Piyush Goyal DR.TANURUNGTA

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com





Date :- 12/03/2022 11:08:18

NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID :-122127659 Ref. By Doctor:-BOB

Lab/Hosp :-

Final Authentication: 12/03/2022 12:24:52

**BOB PACKAGEFEMALE BELOW 40** 

### X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

Impression :- Normal Study

(Please correlate clinically and with relevant further investigations)

\*\*\* End of Report \*\*\*

Page No: 1 of 1

Dr. Piyush Goyal

M.B.B.S., D.M.R.D.

RMC Reg No. 017996

Dr. Poo Ram Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Tej Prakash Gupta DMRD (RADIO DIAGNOSIS) RMC No. 24436

Dr. Hitesh Kumar Sharma M.B.B.S., D.M.R.D. RMC Reg No. 27380

Dr. Piyush Goyal (D.M.R.D.) BI

Transcript by.

BILAL

This report is not valid for medico-legal purpos

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

MITRAL VALVE

AORTIC VALVE

Web

NORMAN

bsite : www.drg	oyalspathlab.com   E-mail : drgoyalpiyush@gmail.com			W-W-W-1
NAME:	MRS.NEHA TAILOR	AGE	28 YRS	A
REF.BY	74	DATE	12.03.2022	00

DITEMONARY WALVE

### 2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

FAIR TRANSTHORACIC ECHOCARIDIOGRAPHIC WINDOW MORPHOLOGY: NORMAL TRICUSPID VALVE

AURTIC VALVE		NORIVIAL		POLIVIONARY VALVE			NORMAL					
		×	M.M	MODE EXAMIT	ATION:	*2	. 9					
AO	21	mm	LA	22	Mm	IVS-D	9	mm				
IVS-S	14	mm	LVID	42	Mm	LVSD	28	mm				
LVPW-D	9	mm	LVPW-S	16	Mm	RV		mm				
RVWT		mm	EDV		MI	LVVS		ml				
LVEF	61%			RWMA		ABSENT						

#### CHAMBERS:

		CHITCHE			
LA	NORMAL -	RA		NORMAL	
LV	NORMAL	RV	100	NORMAL	
PERICARDIL	M	NORMAL	114		

#### 

			(	COLOUR DO	PPLER:			
47	MITRA	AL VALVE					- 0.	
E VELOCITY	1.0	m/sec	PEAK GRADIENT			Mm/hg		
A VELOCITY		m/sec	MEA	MEAN GRADIENT			Mm/hg	
MVA BY PHT		Cm2	MVA BY PLANIMETRY				Cm2	
MITRAL REGURGITAT	ION			8.	ABSENT.			
(3)	AORT	IC VALVE						
PEAK VELOCITY	1.0	m/	sec	PEAK GRADIENT			mm/hg	
AR VMAX		m/	sec MEAN GRADIENT		RADIENT		mm/hg	
AORTIC REGURGITATI	ON			ABSENT		1		
	TRICUS	PID VALVE						
PEAK VELOCITY	0.5	5	m/sec	PEAK GRADIENT		7	mm/hg	
MEAN VELOCITY			m/sec	MEAN GRADIENT			mm/hg	
VMax VELOCITY								
TRICUSPID REGURGITATION			ABSENT					
	PULM	ONARY VAL	.VE					
PEAK VELOCITY		1.2	1.2		PEAK GRADIENT			Mm/hg
MEAN VALOCITY					MEAN GRAD	IENT		Mm/hg
PULMONARY REGURGITATION				ABSENT				

### Impression--

Normal LV size & contractility No RWMA, LVEF 61 %. Normal cardiac chamber. Normal valve

No clot, no vegetation, no pericardial effusion.

(Cardiologist)

NORMAL

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996

Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Tej Prakash Gupta DMRD (RADIO DIAGNOSIS) RMC No. 24436

Dr. Hitesh Kumar Sharma M.B.B.S., D.M.R.D. RMC Reg No. 27380

Transcript by.

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur

Tele: 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com



Date :- 12/03/2022 11:08:18
NAME :- Mrs. NEHA TAILOR

Sex / Age :- Female 28 Yrs 3 Mon 8 Days

Company :- MediWheel

Patient ID :-122127659 Ref. By Doctor:-BOB

Lab/Hosp :-



**BOB PACKAGEFEMALE BELOW 40** 

#### **ULTRA SOUND SCAN OF ABDOMEN**

Liver is mild enlarge in size(16.6cm). Echo-texture is normal. No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

**Gall bladder** is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

Pancreas is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

Spleen is of normal size and shape. Echotexture is normal. No focal lesion is seen.

**Kidneys** are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

**Urinary Bladder**: is well distended and showing smooth wall with normal thickness. Urinary bladder does not show any calculus or mass lesion.

**Uterus** is anteverted and normal in size and measures 83 X 53 X 43mm. Myometrium shows normal echo - pattern. No focal space occupying lesion is seen. Endometrial echo is normal. Endometrial thickness is 9.6mm.

**Both ovaries** are visualised and are normal. No adnexal mass is seen. No enlarged nodes are visualised. No retro-peritoneal lesion is identified. Minimal free fluid is seen in pouch of douglas.

Note: A well defined mixed echogenic SOL measuring 21 x 18mm on left side of pelvis in subcutenous tissues near lateral end of scar likely granuloma.

IMPRESSION: \*Mild hepatomegaly .

\*Minimal free fluis in POD.

Needs clinical correlation & further evaluation

\*\*\* End of Report \*\*\*

KOMAL

Dr. Piyush Goyal M.B.B.S., D.M.R.D. RMC Reg No. 017996 Dr. Poonam Gupta MBBS, MD (Radio Diagnosis) RMC No. 32495

Dr. Tej Prakash Gupta DMRD (RADIO DIAGNOSIS) RMC No. 24436 Dr. Hitesh Kumar Sharma M.B.B.S., D.M.R.D. RMC Reg No. 27380

Transcript by.