TENET DIAGNOSTICS

Customer Name	Mr. Prahlad Kremal	Customer ID	BIL 4768009
Age & Gender	1151	Visit Date	28/9/24

Eye screening

with spectactes / with out spectacles(strike out whichever is not applicable)


Near Vision		
Distance Vision	6 (Plano)	O (Plano)
Colour Vision		-

Right eye

Left eye

observasion / comments

Esc Sight is True (NOVHOU).





Good to Know		450000	
	MR.PRAHLAD KUMAR	Req NO:4768009	
Name		Registered on:28.09.2024	
Age & Gender	43Y/MALE		
		Reported on:28.09.2024	
Ref Doctor	CREDIT CLIENTS	Reported on Zeror	

2D ECHOCARDIOGRAPHY & COLOUR DOPPLER REPORT

M-mode:

Value	Normal range
	(1.9 – 4.0 cm)
	(2.5 - 3.7 cm)
	(0.6 – 1.1 cm)
1.2	(0.6- 1.1 cm)
4.0	(3.5 – 5.5 cm)
2.7	(2.4 – 4.2 cm)
73	ml
26	ml
	50 – 70 %
	4.0 2.7

CHAMBERS:

LEFT ATRIUM: Normal

RIGHT ATRIUM: Normal

LEFT VENTRICLE: CONCENTRIC LVH+

RIGHT VENTRICLE: Normal

VALVES:

MITRAL VALVE: Normal

AORTIC VALVE: Normal

TRICUSPID VALVE: Normal

PULMONARY VALVE: Normal

GREAT ARTERIES:

AORTA: Normal

PULMONARY ARTERY: Normal

od to Know

SEPTAE:

IAS/IVS: Intact

WALL MOTION ABNORMALITIES:

REGIONAL: No RWMA

GLOBAL: Normal

COLOUR DOPPLER:

MITRAL VALVE: TRIVIAL MR, E/A: 1.31

AORTIC VALVE: TRIVIAL AR

TRICUSPID VALVE: TRIVIAL TR, PASP-22 mmHg

PULMONARY VALVE: Normal

CLOT/ VEGETATION: Nil

PERICARDIUM: No effusion

IVC: NORMAL &COLLAPSING

CONCLUSION:

- CONCENTRIC LVH+
- NORMAL VALVES
- NO REGIONAL WALL MOTION ABNORMALITIES
- NORMAL LV SYSTOLIC FUNCTION (EF: 63%)
- NORMAL PA PRESSURE
- NO CLOT/ VEG / PERICARDIAL EFFUSION

Dr. MAHADEV SWAMY B

MBBS, MD, DM Cardiology (JIPMER), FSCAI, FICC Consultant & Interventional Cardiologist KMC No 75242



Mr. PRAHLAD KUMAR	Visit Date	28.09.2024
43 Years/Male	Customer ID	BIL4768009
ARCOFEMI HEALTH CARE LTD -		DID-100009
	43 Years/Male	Visit Date

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size with uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is not visualized, consistent with h/o cholecystectomy.

CBD measures 6mm. No evidence of intraluminal calculi.

PANCREAS is normal in size and echopattern. No evidence of ductal dilatation or calcification.

SPLEEN is normal in size and echopattern. It measures 10.3cms in long axis and 3.1cms in short axis.

KIDNEYS move well with respiration and are normal in size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.0	1.2
Left Kidney	11.2	1.3

URINARY BLADDER is well distended with normal wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE is normal in size and echopattern. It measures 3.9 x 2.8 x 2.7cms (Vol. 16cc).

No evidence of ascites / pleural effusion / para -aortic lymphadenopathy.

IMPRESSION:

- > POST CHOLECYSTECTOMY STATUS.
- > NO SIGNIFICANT ABNORMALITY.

Dr Meera Krishnan C20023 Consultant Radiologist

Tenet Diagnostics Pvt. Ltd.







Name

: MR.PRAHLAD KUMAR

TID/SID

:UMR2016331/ 28325141

Age / Gender : 43 Years / Male

Registered on: 28-Sep-2024 / 09:08 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Reported on : 28-Sep-2024 / 14:19 PM

: BIL4768009 Req.No

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL F	PATHOLOGY
Complete Urine Examination	(CUE), Urine

Investigation	Observed Value	Biological Reference Intervals
Physical Examination		
Colour	Pale Yellow	Straw to Yellow
Method:Physical		
Appearance	Clear	Clear
Method:Physical		
Chemical Examination		
Reaction and pH	6.0	4.6-8.0
Method:pH- Methyl red & Bromothymol blue		
Specific gravity	1.015	1.003-1.035
Method:Bromothymol Blue		
Protein	Negative	Negative
Method:Tetrabromophenol blue	NI P	N
Glucose	Negative	Negative
Method:Glucose oxidase/Peroxidase	NI 1°	Negative
Blood	Negative	Negative
Method:Peroxidase	Negotivo	Negotivo
Ketones	Negative	Negative
Method:Sodium Nitroprusside	Negative	Negative
Bilirubin Method:Dichloroanilinediazonium	Negative	Negative
	Negative	Negative
Leucocytes Method:3 hydroxy5 phenylpyrrole + diazonium	Negative	Negative
Nitrites	Negative	Negative
Nitrites Method:Diazonium + 1,2,3,4 tetrahydrobenzo (h) qui		Nogalivo
3-ol	1101111	
Urobilinogen	0.2	0.2-1.0 mg/dl
Method:Dimethyl aminobenzaldehyde		
Microscopic Examination		
Pus cells (leukocytes)	0-1	2 - 3 /hpf
Method:Microscopy		
Epithelial cells	1-2	2 - 5 /hpf
Method:Microscopy		
RBC (erythrocytes)	Absent	Absent
Method:Microscopy		
Casts	Absent	Occasional hyaline casts may be seen
Method:Microscopy		







Name Ref.By Reg.No : MR.PRAHLAD KUMAR

TID/SID

: UMR2016331/ 28325141

: 43 Years / Male Age / Gender

Registered on: 28-Sep-2024 / 09:08 AM

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Reported on : 28-Sep-2024 / 14:19 PM

: BIL4768009

TEST REPORT

Reference : Arcofemi Health Care Ltd -

Crystals

Absent

Phosphate, oxalate, or urate crystals may

be seen

Method:Microscopy Others

Nil

Nil

Method:Microscopy

Method: Semi Quantitative test ,For CUE

Reference: Godkar Clinical Diagnosis and Management by Laboratory Methods, First South Asia edition. Product kit literature.

Interpretation:

The complete urinalysis provides a number of measurements which look for abnormalities in the urine. Abnormal results from this test can be indicative of a number of conditions including kidney disease, urinary tract infecation or elevated levels of substances which the body is trying to remove through the urine. A urinalysis test can help identify potential health problems even when a person is asymptomatic. All the abnormal results are to be correlated clinically.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.Kavya S N **Consultant Pathologist**









Name

: MR.PRAHLAD KUMAR

TID/SID

: UMR2016331/ 28325142

Age / Gender : 43 Years / Male

Registered on: 28-Sep-2024 / 09:08 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Reported on : 28-Sep-2024 / 16:07 PM

: BIL4768009 Reg.No

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF HEMATOPATHOLOGY

Blood Grouping ABO And Rh Typing, EDTA Whole Blood

Results Parameter Blood Grouping (ABO) O Rh Typing (D) **NEGATIVE** Note Suggested further comfirmation by gel card method.

Method: Hemagglutination Tube Method by Forward & Reverse Grouping

Reference: Tulip kit literature

Interpretation: The ABO grouping and Rh typing test determines blood type grouping (A,B, AB, O) and the Rh factor (positive or negative). A person's blood type is based on the presence or absence of certain antigens on the surface of their red blood cells and certain antibodies in the plasma. ABO antigens are poorly expresses at birth, increase gradually in strength and become fully expressed around 1 year of age.

Note: Records of previous blood grouping/Rh typing not available. Please verify before transfusion.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debluena Thakur

Dr Debleena Thakur **Consultant Pathologist**









Name Age / Gender Ref.By

: MR.PRAHLAD KUMAR

TID/SID

:UMR2016331/ 28325142

: 43 Years / Male : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Registered on: 28-Sep-2024 / 09:08 AM

Req.No : BIL4768009

Reported on : 28-Sep-2024 / 14:54 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF HEMATOPATHOLOGY

Erythrocyte Sedimentation Rate (ESR), Whole Blood

Biological Reference Intervals Investigation Observed Value 06 <=15 mm/hour ESR 1st Hour

Method:Modified Westergren

Complete Blood Count (CBC), EDTA Whole Blood

Investigation	Observed Value	Biological Reference Interval
Hemoglobin Method:Spectrophotometry	14.6	13.0-18.0 g/dL
Packed Cell Volume Method:Derived from Impedance	43.1	40-54 %
Red Blood Cell Count. Method:Impedance Variation	4.98	4.3-6.0 Mill/Cumm
Mean Corpuscular Volume Method:Derived from Impedance	86.4	78-100 fL
Mean Corpuscular Hemoglobin Method:Derived from Impedance	29.3	27-32 pg
Mean Corpuscular Hemoglobin Concentration Method:Derived from Impedance	33.9	31.5-36 g/dL
Red Cell Distribution Width - CV Method:Derived from Impedance	13.3	11.5-16.0 %
Red Cell Distribution Width - SD Method:Derived from Impedance	43.7	39-46 fL
Total WBC Count. Method:Impedance Variation	5550	4000-11000 cells/cumm
Neutrophils Method:Impedance Variation, Flowcytometry	55.0	40-75 %
Lymphocytes Method:Microscopy	33.7	20-45 %
Eosinophils Method:Impedance Variation,Method_Desc= Flow Cytometry	1.7	01-06 %
Monocytes Method:Impedance Variation, Flowcytometry	8.9	01-10 %
Basophils. Method:Impedance Variation,Method_Desc= Flow Cytometry	0.7	00-02 %







Name : MR.PRAHLAD KUMAR

Registered on: 28-Sep-2024 / 09:08 AM

:UMR2016331/ 28325142

Age / Gender

: 43 Years / Male

Ref.By

Note

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

TID/SID

Reported on : 28-Sep-2024 / 14:54 PM

Req.No : BIL4768009

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

Absolute Neutrophils Count. Method:Calculated	3053	1500-6600 cells/cumm
Absolute Lymphocyte Count Method:Calculated	1870	1500-3500 cells/cumm
Absolute Eosinophils count. Method:Calculated	94	40-440 cells/cumm
Absolute Monocytes Count. Method:Calculated	494	<1000 cells/cumm
Absolute Basophils count. Method:Calculated	39	<200 cells/cumm
Platelet Count. Method:Impedance Variation	1.13	1.4-4.4 lakhs/cumm
Mean Platelet Volume. Method:Derived from Impedance	12.3	7.9-13.7 fL
Plateletcrit. Method:Derived from Impedance	0.14	0.18-0.28 %
Platelets	Decreased in number, ma	acroplatelets seen. Manual platelet

Method: Automated Hematology Analyzer, Microscopy

Reference: Dacie and Lewis Practical Hematology, 12th Edition

Interpretation: A Complete Blood Picture (CBP) is a screening test which can aid in the diagnosis of a variety of conditions and diseases such as anemia, leukemia, bleeding disorders and infections. This test is also useful in monitoring a person's reaction to treatment when a condition which affects blood cells has been diagnosed. All the abnormal results are to be correlated clinically.

count=1.3 lakhs/cumm

Kindly correlate clinically

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Debluena Thakua

Dr Debleena Thakur **Consultant Pathologist**







Name Age / Gender : MR.PRAHLAD KUMAR

TID/SID

: UMR2016331/ 28325144F

: 43 Years / Male

Registered on: 28-Sep-2024 / 09:08 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Reported on : 28-Sep-2024 / 14:01 PM

Reg.No : BIL4768009

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Blood Urea Nitrogen (BUN), Serum

Investigation	Observed Value	Biological Reference Interval
Blood Urea Nitrogen.	10	6-20 mg/dL

Method:Kinetic, Urease - GLDH, Calculated

Interpretation: Urea is a waste product formed in the liver when protein is metabolized. Urea is released by the liver into the blood and is carried to the kidneys, where it is filtered out of the blood and released into the urine. Since this is a continuous process, there is usually a small but stable amount of urea nitrogen in the blood. However, when the kidneys cannot filter wastes out of the blood due to disease or damage, then the level of urea in the blood will rise. The blood urea nitrogen (BUN) evaluates kidney function in a wide range of circumstances, to diagnose kidney disease, and to monitor people with acute or chronic kidney dysfunction or failure. It also may be used to evaluate a person's general health status as well.

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics

Creatinine, Serum

Investigation	Observed Value	Biological Reference Interval
Creatinine.	0.97	0.7-1.3 mg/dL
Method:Spectrophotometry Jaffe - IDMS Traceable		

Interpretation:

Creatinine is a nitrogenous waste product produced by muscles from creatine. Creatinine is majorly filtered from the blood by the kidneys and released into the urine, so serum creatinine levels are usually a good indicator of kidney function. Serum creatinine is more specific and more sensitive indicator of renal function as compared to BUN because it is produced from muscle at a constant rate and its level in blood is not affected by protein catabolism or other exogenous products. It is also not reabsorbed and very little is secreted by tubules making it a reliable marker. Serum creatinine levels are increased in pre renal, renal and post renal azotemia, active acromegaly and gigantism. Decreased serum creatinine levels are seen in pregnancy and increasing age.

Biological reference interval changed; Reference: Tietz Textbook of Clinical Chemistry & Molecular Diagnostics, Fifth Edition.

Glucose Fasting (FBS), Sodium Fluoride Plasma

	3 (//	
Investigation	Observed Value	Biological Reference Interval
Glucose Fasting Method:Hexokinase	86	Normal: <100 mg/dL Impaired FG: 100-125 mg/dL Diabetes mellitus: >/=126 mg/dL

Interpretation: It measures the Glucose levels in the blood with a prior fasting of 9-12 hours. The test helps screen a symptomatic/ asymptomatic person who is at risk for Diabetes. It is also used for regular monitoring of glucose levels in people with Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2022





: Arcofemi Health Care Ltd -

 Name
 : MR.PRAHLAD KUMAR
 TID/SID
 : UMR2016331/28325143

 Age / Gender
 : 43 Years / Male
 Registered on : 28-Sep-2024 / 09:08 AM

 Ref.By
 : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS
 Collected on : 28-Sep-2024 / 09:08 AM

 Req.No
 : BIL4768009
 Reported on : 28-Sep-2024 / 14:19 PM

Glucose Post Prandial (PPBS), Sodium Fluoride Plasma

Reference

Investigation	Observed Value	Biological Reference Interval
Glucose Post Prandial Method:Hexokinase	90	Normal : <140 mg/dL Impaired PG: 140-199 mg/dL Diabetes mellitus: >/=200 mg/dL

TEST REPORT

Interpretation: This test measures the blood sugar levels 2 hours after a normal meal. Abnormally high blood sugars 2 hours after a meal reflect that the body is not producing sufficient insulin which is indicative of Diabetes.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2020.

Glycosylated Hemoglobin (HbA1C), EDTA Whole Blood

Investigation	Observed Value	Biological Reference Interval
Glycosylated Hemoglobin (HbA1c) Method:High-Performance Liquid Chromatography	5.2	Non-diabetic: <= 5.6 % Pre-diabetic: 5.7 - 6.4 % Diabetic: >= 6.5 %
Estimated Average Glucose (eAG) Method:High-Performance Liquid Chromatography	103	mg/dL

Interpretation: It is an index of long-term blood glucose concentrations and a measure of the risk for developing microvascular complications in patients with diabetes. Absolute risks of retinopathy and nephropathy are directly proportional to the mean HbA1c concentration. In persons without diabetes, HbA1c is directly related to risk of cardiovascular disease.

In known diabetic patients, HbA1c can be considered as a tool for monitoring the glycemic control.

Excellent Control - 6 to 7 %,

Fair to Good Control - 7 to 8 %,

Unsatisfactory Control - 8 to 10 %

and Poor Control - More than 10 %.

Reference: American Diabetes Association. Standards of Medical Care in Diabetes-2018.

Bun/Creatinine Ratio. Serum

	•	
Investigation	Observed Value	
BUN/Creatinine Ratio	10	
Method:Calculated		

Reference:

A Manual of Laboratory Diagnostic Tests. Edition 7, Lippincott Williams and Wilkins, By Frances Talaska Fischbach, RN, BSN, MSN, and Marshall Barnett Dunning 111, BS, MS, Ph.D.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---





Name : MR.PRAHLAD KUMAR TID/SID : UMR2016331/

Age / Gender : 43 Years / Male Registered on : 28-Sep-2024 / 09:08 AM

Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on :

Req.No : BIL4768009 Reported on :

TEST REPORT Reference : Arcofemi Health Care Ltd -

איניעס \ Dr.Kavya S N Consultant Pathologist







Name Age / Gender Ref.By

: MR.PRAHLAD KUMAR

TID/SID

:UMR2016331/ 28325143

: 43 Years / Male

Registered on: 28-Sep-2024 / 09:08 AM

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Reported on : 28-Sep-2024 / 14:19 PM

: BIL4768009 Reg.No

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Lipid Profile, Serum

Lipid Frome, Octum			
Investigation	Observed Value	Biological Reference Interval	
Total Cholesterol Method:Spectrophotometry , CHOD - POD	133	Desirable: < 200 mg/dL Borderline: 200-239 mg/dL High: >/= 240 mg/dL	
HDL Cholesterol Method:Spectrophotometry , Direct Measurement	47	Optimal : >=60 mg/dL Borderline : 40-59 mg/dL High Risk <40 mg/dL	
Non HDL Cholesterol Method:Calculated	86	Optimal: <130 mg/dL Above Optimal: 130-159 mg/dL Borderline: 160-189 mg/dL High Risk: 190-219 mg/dL Very high Risk: >=220 mg/dL	
LDL Cholesterol Method:Calculated	71.4	Optimum: <100 mg/dL Near/above optimum: 100-129 mg/dL Borderline: 130-159 mg/dL High: 160-189 mg/dL Very high: >/=190 mg/dL	
VLDL Cholesterol Method:Calculated	14.60	<30 mg/dL	
Total Cholesterol/HDL Ratio Method:Calculated	2.83	Optimal: <3.3 Low Risk: 3.4-4.4 Average Rsik: 4.5-7.1 Moderate Risk: 7.2-11.0 High Risk: >11.0	
LDL/HDL Ratio Method:Calculated	1.52	Optimal : 0.5-3.0 Borderline : 3.1-6.0 High Risk : >6.0	
Triglycerides Method:Spectrophotometry, Enzymatic - GPO/POD	73	Normal:<150 mg/dL Borderline: 150-199 mg/dL High: 200-499 mg/dL Very high: >/=500 mg/dL mg/dl #	

Interpretation: Lipids are fats and fat-like substances which are important constituents of cells and are rich sources of energy. A lipid profile typically includes total cholesterol, high density lipoproteins (HDL), low density lipoprotein (LDL), chylomicrons, triglycerides, very low density lipoproteins (VLDL), Cholesterol/HDL ratio .The lipid profile is used to assess the risk of developing a heart disease and to monitor its treatment. The results of the lipid profile are evaluated along with other known risk factors associated with heart disease to plan and monitor treatment. Treatment options require clinical correlation. Reference: Third Report of the National Cholesterol Education program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III), JAMA 2001.

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore





Name : MR.PRAHLAD KUMAR TID/SID : UMR2016331/

Age / Gender : 43 Years / Male Registered on : 28-Sep-2024 / 09:08 AM

Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on :

Req.No : BIL4768009 Reported on :

TEST REPORT Reference : Arcofemi Health Care Ltd -

Your SM

Dr.Kavya S N Consultant Pathologist





: MR.PRAHLAD KUMAR Name

TID/SID

:UMR2016331/ 28325143

Age / Gender

: 43 Years / Male

Registered on: 28-Sep-2024 / 09:08 AM

Ref.By

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Req.No

: BIL4768009

Reported on : 28-Sep-2024 / 14:56 PM

TEST REPORT

Reference

: Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Liver Function Test (LFT), Serum

Investigation	Result	Biological Reference Interval
Total Bilirubin. Method:Spectrophotometry, Diazo method	1.33	Neonates: <=15.0 mg/dL Adults: <=1.2 mg/dL
Direct Bilirubin. Method:Spectrophotometry, Diazo method	0.74	<=0.30 mg/dL
Indirect Bilirubin. Method:Calculated	0.59	Neonates: <= 14.7 mg/dL Adults: <= 1.0 mg/dL
Alanine Aminotransferase ,(ALT/SGPT) Method: IFCC without pyridoxal phosphate activation	33	<=41 U/L
Aspartate Aminotransferase,(AST/SGOT) Method: IFCC without pyridoxal phosphate activation	32	<=40 U/L
ALP (Alkaline Phosphatase). Method:Spectrophotometry, IFCC	95	40-129 U/L
Gamma GT. Method:Spectrophotometry , IFCC	20	<60 U/L
Total Protein. Method:Spectrophotometry, Biuret	7.2	6.4-8.3 g/dL
Albumin. Method:Spectrophotometry, Bromcresol Green	4.6	3.5-5.2 g/dL
Globulin. Method:Spectrophotometry, Bromcresol Green	2.60	2.0-3.5 g/dL
A/GRatio. Method:Calculated	1.77	1.1-2.5

Interpretation: Liver functions tests help to identify liver disease, its severity, and its type. Generally these tests are performed in combination, are abnormal in liver disease, and the pattern of abnormality is indicative of the nature of liver disease. An isolated abnormality of a single liver function test usually means a non-hepatic cause. If several liver function tests are simultaneously abnormal, then hepatic etiology is likely.

--- End Of Report ---

Dr.Kavya S N **Consultant Pathologist**

^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore



Age / Gender



TO VERIFY THE REPORT ONLINE

:UMR2016331/ 28325143

Name : MR.PRAHLAD KUMAR

TID/SID : 43 Years / Male Registered on: 28-Sep-2024 / 09:08 AM

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM Ref.By

: BIL4768009 Req.No

Reported on : 28-Sep-2024 / 13:35 PM

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Prostate Specific Antigen (PSA) Total Serum

riostate opcomo Antigen (rioA) rotan			
Investigation	Observed Value	Biological Reference Interval	
Prostate Specific Antigen (PSA) Total	0.599	0.0-4.0 ng/mL	
Method:ECLIA			

Interpretation: PSA is a protein produced by cells in the prostate and is used to screen men for prostate cancer. PSA levels are elevated in Prostate cancer, and other conditions such as benign prostatic hyperplasia (BPH) and inflammation of the prostate. An elevated PSA may be followed by a biopsy and other tests like urinalysis and ultrasound to rule out urinary tract infections and for an accurate diagnosis. PSA levels are vital to determine the effectiveness of treatment and to detect recurrence in diagnosed cases of prostate cancer.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.M.G.Satish **Consultant Pathologist**









Name : MR.PRAHLAD KUMAR TID/SID : UMR2016331/ 28325143 Age / Gender : 43 Years / Male Registered on : 28-Sep-2024 / 09:08 AM

Ref.By : ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Reg.No : BIL4768009 Reported on : 28-Sep-2024 / 13:27 PM

TEST REPORT Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I

Thyroid Profile (T3,T4,TSH), Serum

111yrola 1 rollic (10,14,1011), Octain		
Investigation	Observed Value	Biological Reference Interval
Triiodothyronine Total (T3) Method:ECLIA	1.07	0.80-2.00 ng/mL Note: Biological Reference Ranges are changed due to change in method of testing.
Thyroxine Total (T4) Method:ECLIA	8.46	4.6-12.0 μg/dL
Thyroid Stimulating Hormone (TSH) Method:ECLIA	2.42	0.27-4.20 μIU/mL

Interpretation: A thyroid profile is used to evaluate thyroid function and/or help diagnose hypothyroidism and hyperthyroidism due to various thyroid disorders. T4 and T3 are hormones produced by the thyroid gland. They help control the rate at which the body uses energy, and are regulated by a feedback system. TSH from the pituitary gland stimulates the production and release of T4 (primarily) and T3 by the thyroid. Most of the T4 and T3 circulate in the blood bound to protein. A small percentage is free (not bound) and is the biologically active form of the hormones

Reference: Tietz Fundamentals of Clinical Chemistry and Molecular Diagnostics, Carl A. Burtis, David E. Bruns.

--- End Of Report ---

Debleena Thakua

Dr Debleena Thakur Consultant Pathologist



^{*} Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore







Name Age / Gender : MR.PRAHLAD KUMAR : 43 Years / Male

Registered on: 28-Sep-2024 / 09:08 AM

:UMR2016331/ 28325143

: ARCOFEMI HEALTH CARE LTD - MEDI WHEELS Collected on : 28-Sep-2024 / 09:08 AM

Ref.By

TID/SID

Reported on : 28-Sep-2024 / 14:19 PM

: BIL4768009 Req.No

TEST REPORT

Reference : Arcofemi Health Care Ltd -

DEPARTMENT OF CLINICAL CHEMISTRY I Uric Acid, Serum Observed Value Biological Reference Interval Investigation 6.5 3.4-7.0 mg/dL Uric Acid. Method:Enzymatic

Interpretation: It is the major product of purine catabolism. Hyperuricemia can result due to increased formation or decreased excretion of uric acid which can be due to several causes like metabolic disorders, psoriasis, tissue hypoxia, pre-eclampsia, alcohol, lead poisoning, acute or chronic kidney disease, etc. Hypouricemia may be seen in severe hepato cellular disease and defective renal tubular reabsorption of uric acid.

* Sample processed at Regional Reference Laboratory, Tenet Diagnostics, Bangalore

--- End Of Report ---

Dr.Kavya S N **Consultant Pathologist**







PLEASE SCAN OR CODE

Name : Mr. PRAHLAD KUMAR TID : UMR2016331

Age/Gender: 43 Years/MaleRegistered On: 28-Sep-2024 09:08 AMRef By: ARCOFEMI HEALTH CARE LTD - MEDI WHEELSReported On: 28-Sep-2024 11:44 AMReg.No: BIL4768009Reference: Arcofemi Health Care Ltd

- Medi Whe

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in size with uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is not visualized, consistent with h/o cholecystectomy.

CBD measures 6mm. No evidence of intraluminal calculi.

PANCREAS is normal in size and echopattern. No evidence of ductal dilatation or calcification.

SPLEEN is normal in size and echopattern. It measures 10.3cms in long axis and 3.1cms in short axis.

KIDNEYS move well with respiration and are normal in size and echopattern.

Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	10.0	1.2
Left Kidney	11.2	1.3

URINARY BLADDER is well distended with normal wall thickness. It has clear contents. No evidence of diverticula.

PROSTATE is normal in size and echopattern. It measures 3.9 x 2.8 x 2.7cms (Vol: 16cc).

No evidence of ascites / pleural effusion / para -aortic lymphadenopathy.

IMPRESSION:

- POST CHOLECYSTECTOMY STATUS.
- NO SIGNIFICANT ABNORMALITY.

*** End Of Report ***

Dr Meera Krishnan Consultant Radiologist





PLEASE SCAN OR CODE

Name : Mr. PRAHLAD KUMAR TID : UMR2016331

Age/Gender: 43 Years/MaleRegistered On: 28-Sep-2024 09:08 AMRef By: ARCOFEMI HEALTH CARE LTD - MEDI WHEELSReported On: 28-Sep-2024 10:24 AM

Reg.No : BIL4768009 Reference : Arcofemi Health Care Ltd

- Medi Whe

X-Ray Chest PA View

FINDINGS AND IMPRESSION:

Lung fields appear normal.

Cardiomegaly is noted.

Hila is normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

*** End Of Report ***

Dr Suhas C MConsultant Radiologist





PLEASE SCAN OR CODE

Name : Mr. PRAHLAD KUMAR TID : UMR2016331

Age/Gender: 43 Years/MaleRegistered On: 28-Sep-2024 09:08 AMRef By: ARCOFEMI HEALTH CARE LTD - MEDI WHEELSReported On: 28-Sep-2024 10:24 AM

Reg.No : BIL4768009 Reference : Arcofemi Health Care Ltd

- Medi Whe

X-Ray Chest PA View

FINDINGS AND IMPRESSION:

Lung fields appear normal.

Cardiomegaly is noted.

Hila is normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

*** End Of Report ***

Dr Suhas C MConsultant Radiologist