

Name : MR. JAISWAL SACHIN

Age / Gender : 38 Years / Male

Consulting Dr. : -

Reg. Location : Malad West (Main Centre)



Use a QR Code Scanner Application To Scan the Code

Collected : 31-Mar-2022 / 09:23

Reported :31-Mar-2022 / 13:52

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

CBC (Complete Blood Count), Blood			
<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
RBC PARAMETERS			
Haemoglobin	12.9	13.0-17.0 g/dL	Spectrophotometric
RBC	4.32	4.5-5.5 mil/cmm	Elect. Impedance
PCV	40.7	40-50 %	Measured
MCV	94.3	80-100 fl	Calculated
MCH	29.9	27-32 pg	Calculated
MCHC	31.7	31.5-34.5 g/dL	Calculated
RDW	14.5	11.6-14.0 %	Calculated
WBC PARAMETERS			
WBC Total Count	6410	4000-10000 /cmm	Elect. Impedance
WBC DIFFERENTIAL AND ABSO	LUTE COUNTS		
Lymphocytes	37.0	20-40 %	
Absolute Lymphocytes	2370	1000-3000 /cmm	Calculated
Monocytes	8.9	2-10 %	
Absolute Monocytes	570	200-1000 /cmm	Calculated
Neutrophils	52.7	40-80 %	
Absolute Neutrophils	3370	2000-7000 /cmm	Calculated
Eosinophils	1.0	1-6 %	
Absolute Eosinophils	60	20-500 /cmm	Calculated
Basophils	0.4	0.1-2 %	
Absolute Basophils	20	20-100 /cmm	Calculated

WBC Differential Count by Absorbance & Impedance method/Microscopy.

PLATELET PARAMETERS

Platelet Count	221000	150000-400000 /cmm	Elect. Impedance
MPV	11.9	6-11 fl	Calculated
PDW	23.0	11-18 %	Calculated

RBC MORPHOLOGY

Immature Leukocytes

Hypochromia	-
Microcytosis	-

Page 1 of 9

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343



Name : MR.JAISWAL SACHIN

:38 Years / Male Age / Gender

Consulting Dr. Collected :31-Mar-2022 / 09:23

Reported :31-Mar-2022 / 12:27 Reg. Location : Malad West (Main Centre)

Macrocytosis

Anisocytosis

Poikilocytosis

Polychromasia

Target Cells

Basophilic Stippling

Normoblasts

Others Normocytic, Normochromic

WBC MORPHOLOGY

PLATELET MORPHOLOGY

COMMENT

Specimen: EDTA Whole Blood

ESR, EDTA WB 2-15 mm at 1 hr. Westergren

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West *** End Of Report ***









Page 2 of 9

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code



Name : MR.JAISWAL SACHIN

Age / Gender : 38 Years / Male

Consulting Dr. :

Reg. Location: Malad West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Collected : 31-Mar-2022 / 09:23

Reported :31-Mar-2022 / 12:51

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
GLUCOSE (SUGAR) FASTING, Fluoride Plasma	83.2	Non-Diabetic: < 100 mg/dl Impaired Fasting Glucose: 100-125 mg/dl Diabetic: >/= 126 mg/dl	Hexokinase
GLUCOSE (SUGAR) PP, Fluoride Plasma PP/R	66.3	Non-Diabetic: < 140 mg/dl Impaired Glucose Tolerance: 140-199 mg/dl Diabetic: >/= 200 mg/dl	Hexokinase
BILIRUBIN (TOTAL), Serum	0.40	0.1-1.2 mg/dl	Colorimetric
BILIRUBIN (DIRECT), Serum	0.15	0-0.3 mg/dl	Diazo
BILIRUBIN (INDIRECT), Serum	0.25	0.1-1.0 mg/dl	Calculated
TOTAL PROTEINS, Serum	7.4	6.4-8.3 g/dL	Biuret
ALBUMIN, Serum	4.2	3.5-5.2 g/dL	BCG
GLOBULIN, Serum	3.2	2.3-3.5 g/dL	Calculated
A/G RATIO, Serum	1.3	1 - 2	Calculated
SGOT (AST), Serum	24.8	5-40 U/L	NADH (w/o P-5-P)
SGPT (ALT), Serum	39.8	5-45 U/L	NADH (w/o P-5-P)
GAMMA GT, Serum	24.4	3-60 U/L	Enzymatic
ALKALINE PHOSPHATASE, Serum	97.5	40-130 U/L	Colorimetric
BLOOD UREA, Serum	26.5	12.8-42.8 mg/dl	Kinetic
BUN, Serum	12.4	6-20 mg/dl	Calculated
CREATININE, Serum	0.77	0.67-1.17 mg/dl	Enzymatic
eGFR, Serum	120	>60 ml/min/1.73sqm	Calculated
URIC ACID, Serum	5.3	3.5-7.2 mg/dl	Enzymatic

Page 3 of 9



Name : MR. JAISWAL SACHIN

Age / Gender : 38 Years / Male

Consulting Dr. : -

Reg. Location: Malad West (Main Centre)

Authenticity Check

Use a QR Code Scanner Application To Scan the Code

Collected

: 31-Mar-2022 / 12:16

Reported :31-Mar-2022 / 16:11

Urine Sugar (Fasting)AbsentAbsentUrine Ketones (Fasting)AbsentAbsent

Urine Sugar (PP) Absent Absent Urine Ketones (PP) Absent Absent

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***







Dr.ANUPA DIXIT
M.D.(PATH)
Consultant Pathologist & Lab
Director

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com

Page 4 of 9



Name : MR.JAISWAL SACHIN

Age / Gender : 38 Years / Male

Consulting Dr. : - Collected : 31-Mar-2022 / 09:23

Reg. Location: Malad West (Main Centre) Reported: 31-Mar-2022 / 13:10

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE GLYCOSYLATED HEMOGLOBIN (HbA1c)

<u>PARAMETER</u> <u>RESULTS</u> <u>BIOLOGICAL REF RANGE</u> <u>METHOD</u>

Glycosylated Hemoglobin 5.2 Non-Diabetic Level: < 5.7 % HPLC (HbA1c), EDTA WB - CC Prediabetic Level: 5.7-6.4 %

Prediabetic Level: 5.7-6.4 % Diabetic Level: >/= 6.5 %

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code

Estimated Average Glucose 102.5 mg/dl Calculated

(eAG), EDTA WB - CC

Intended use:

• In patients who are meeting treatment goals, HbA1c test should be performed at least 2 times a year

• In patients whose therapy has changed or who are not meeting glycemic goals, it should be performed quarterly

• For microvascular disease prevention, the HbA1C goal for non pregnant adults in general is Less than 7%.

Clinical Significance:

• HbA1c, Glycosylated hemoglobin or glycated hemoglobin, is hemoglobin with glucose molecule attached to it.

• The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of glycosylated hemoglobin in the blood.

Test Interpretation:

- The HbA1c test evaluates the average amount of glucose in the blood over the last 2 to 3 months by measuring the percentage of Glycosylated hemoglobin in the blood.
- HbA1c test may be used to screen for and diagnose diabetes or risk of developing diabetes.
- To monitor compliance and long term blood glucose level control in patients with diabetes.
- Index of diabetic control, predicting development and progression of diabetic micro vascular complications.

Factors affecting HbA1c results:

Increased in: High fetal hemoglobin, Chronic renal failure, Iron deficiency anemia, Splenectomy, Increased serum triglycerides, Alcohol ingestion, Lead/opiate poisoning and Salicylate treatment.

Decreased in: Shortened RBC lifespan (Hemolytic anemia, blood loss), following transfusions, pregnancy, ingestion of large amount of Vitamin E or Vitamin C and Hemoglobinopathies

Reflex tests: Blood glucose levels, CGM (Continuous Glucose monitoring)

References: ADA recommendations, AACC, Wallach's interpretation of diagnostic tests 10th edition.

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***







Dr.JYOT THAKKER M.D. (PATH), DPB Pathologist & AVP(Medical Services)

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com

Page 5 of 9



Name : MR.JAISWAL SACHIN

:38 Years / Male Age / Gender

Consulting Dr. Collected

Reported Reg. Location : Malad West (Main Centre)

Use a OR Code Scanner

Authenticity Check

Application To Scan the Code

:31-Mar-2022 / 09:23 :31-Mar-2022 / 15:20

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE URINE EXAMINATION REPORT

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
PHYSICAL EXAMINATION			
Color	Pale yellow	Pale Yellow	-
Reaction (pH)	6.5	4.5 - 8.0	Chemical Indicator
Specific Gravity	1.005	1.001-1.030	Chemical Indicator
Transparency	Clear	Clear	-
Volume (ml)	40	-	-
CHEMICAL EXAMINATION			
Proteins	Absent	Absent	pH Indicator
Glucose	Absent	Absent	GOD-POD
Ketones	Absent	Absent	Legals Test
Blood	Absent	Absent	Peroxidase
Bilirubin	Absent	Absent	Diazonium Salt
Urobilinogen	Normal	Normal	Diazonium Salt
Nitrite	Absent	Absent	Griess Test
MICROSCOPIC EXAMINATION			
Leukocytes(Pus cells)/hpf	1-2	0-5/hpf	
Red Blood Cells / hpf	Absent	0-2/hpf	

Red Blood Cells / hpt 0-2/hpt

Epithelial Cells / hpf 0-1

Casts Absent Absent Crystals **Absent Absent** Amorphous debris Absent Absent

Bacteria / hpf 2-3 Less than 20/hpf

Others







Dr.SHASHIKANT DIGHADE M.D. (PATH) **Pathologist**

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com

Page 6 of 9

^{*}Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West *** End Of Report **



Name : MR. JAISWAL SACHIN

Age / Gender :38 Years / Male

Consulting Dr. Collected Reported

Reg. Location : Malad West (Main Centre)



Authenticity Check

Use a OR Code Scanner Application To Scan the Code

: 31-Mar-2022 / 09:23 :31-Mar-2022 / 14:11

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE BLOOD GROUPING & Rh TYPING

PARAMETER RESULTS

ABO GROUP AΒ

Rh TYPING **POSITIVE**

NOTE: Test performed by automated column agglutination technology (CAT) which is more sensitive than conventional methods.

Specimen: EDTA Whole Blood and/or serum

Clinical significance:

ABO system is most important of all blood group in transfusion medicine

Limitations:

- ABO blood group of new born is performed only by cell (forward) grouping because allo antibodies in cord blood are of maternal origin.
- Since A & B antigens are not fully developed at birth, both Anti-A & Anti-B antibodies appear after the first 4 to 6 months of life. As a result, weaker reactions may occur with red cells of newborns than of adults.
- Confirmation of newborn's blood group is indicated when A & B antigen expression and the isoagglutinins are fully developed at 2 to 4 years of age & remains constant throughout life.
- Cord blood is contaminated with Wharton's jelly that causes red cell aggregation leading to false positive result
- The Hh blood group also known as Oh or Bombay blood group is rare blood group type. The term Bombay is used to refer the phenotype that lacks normal expression of ABH antigens because of inheritance of hh genotype.

Refernces:

- Denise M Harmening, Modern Blood Banking and Transfusion Practices- 6th Edition 2012. F.A. Davis company. Philadelphia
- AABB technical manual

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West *** End Of Report *'







Dr.MILLU JAIN M.D.(PATH) **Pathologist**

Page 7 of 9

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com



Name : MR. JAISWAL SACHIN

:38 Years / Male Age / Gender

Consulting Dr.

Reg. Location : Malad West (Main Centre)



Authenticity Check

Use a OR Code Scanner

Collected

Reported

Application To Scan the Code

:31-Mar-2022 / 09:23 :31-Mar-2022 / 13:10

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE LIPID PROFILE

<u>PARAMETER</u>	<u>RESULTS</u>	BIOLOGICAL REF RANGE	<u>METHOD</u>
CHOLESTEROL, Serum	199.9	Desirable: <200 mg/dl Borderline High: 200-239mg/dl High: >/=240 mg/dl	Enzymatic
TRIGLYCERIDES, Serum	93.9	Normal: <150 mg/dl Borderline-high: 150 - 199 mg/dl High: 200 - 499 mg/dl Very high:>/=500 mg/dl	Enzymatic
HDL CHOLESTEROL, Serum	40.6	Desirable: >60 mg/dl Borderline: 40 - 60 mg/dl Low (High risk): <40 mg/dl	Enzymatic
NON HDL CHOLESTEROL, Serum	159.3	Desirable: <130 mg/dl Borderline-high:130 - 159 mg/dl High:160 - 189 mg/dl Very high: >/=190 mg/dl	Calculated
LDL CHOLESTEROL, Serum	140.0	Optimal: <100 mg/dl Near Optimal: 100 - 129 mg/dl Borderline High: 130 - 159 mg/dl High: 160 - 189 mg/dl Very High: >/= 190 mg/dl	Calculated
VLDL CHOLESTEROL, Serum	19.3	< /= 30 mg/dl	Calculated
CHOL / HDL CHOL RATIO, Serum	4.9	0-4.5 Ratio	Calculated
LDL CHOL / HDL CHOL RATIO, Serum	3.5	0-3.5 Ratio	Calculated

^{*}Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West *** End Of Report ***







Dr.JYOT THAKKER M.D. (PATH), DPB Pathologist & AVP(Medical Services)

ADDRESS: 2nd Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com

Page 8 of 9



Name : MR. JAISWAL SACHIN

Age / Gender : 38 Years / Male

Consulting Dr. : - Collected : 31-Mar-2022 / 09:23

Reg. Location : Malad West (Main Centre) Reported : 31-Mar-2022 / 12:51

AERFOCAMI HEALTHCARE BELOW 40 MALE/FEMALE THYROID FUNCTION TESTS

PARAMETER	RESULTS	BIOLOGICAL REF RANGE	<u>ME I HOD</u>
Free T3, Serum	4.6	3.5-6.5 pmol/L	ECLIA
Free T4, Serum	13.5	11.5-22.7 pmol/L	ECLIA
sensitiveTSH, Serum	3.09	0.35-5.5 microIU/ml	ECLIA

Interpretation:

A thyroid panel is used to evaluate thyroid function and/or help diagnose various thyroid disorders.

Clinical Significance:

- 1)TSH Values between 5.5 to 15 microIU/ml should be correlated clinically or repeat the test with new sample as physiological factors can give falsely high TSH.
- 2)TSH values may be trasiently altered becuase of non thyroidal illness like severe infections, liver disease, renal and heart severe burns, trauma and surgery etc.

TSH	FT4 / T4	FT3/T3	Interpretation
High	Normal	Normal	Subclinical hypothyroidism, poor compliance with thyroxine, drugs like amiodarone, Recovery phase of non-thyroidal illness, TSH Resistance.
High	Low	Low	Hypothyroidism, Autoimmune thyroiditis, post radio iodine Rx, post thyroidectomy, Anti thyroid drugs, tyrosine kinase inhibitors & amiodarone, amyloid deposits in thyroid, thyroid tumors & congenital hypothyroidism.
Low	High	High	Hyperthyroidism, Graves disease, toxic multinodular goiter, toxic adenoma, excess iodine or thyroxine intake, pregnancy related (hyperemesis gravidarum, hydatiform mole)
Low	Normal	Normal	Subclinical Hyperthyroidism, recent Rx for Hyperthyroidism, drugs like steroids & dopamine), Non thyroidal illness.
Low	Low	Low	Central Hypothyroidism, Non Thyroidal Illness, Recent Rx for Hyperthyroidism.
High	High	High	Interfering anti TPO antibodies, Drug interference: Amiodarone, Heparin, Beta Blockers, steroids & anti epileptics.

Diurnal Variation:TSH follows a diurnal rhythm and is at maximum between 2 am and 4 am, and is at a minimum between 6 pm and 10 pm. The variation is on the order of 50 to 206%. Biological variation:19.7%(with in subject variation)

Reflex Tests: Anti thyroid Antibodies, USG Thyroid , TSH receptor Antibody. Thyroglobulin, Calcitonin

Limitations: Samples should not be taken from patients receiving therapy with high biotin doses (i.e. >5 mg/day) until atleast 8 hours following the last biotin administration.

Reference:

- 1.O.koulouri et al. / Best Practice and Research clinical Endocrinology and Metabolism 27(2013)
- 2.Interpretation of the thyroid function tests, Dayan et al. THE LANCET . Vol 357
- 3. Tietz , Text Book of Clinical Chemistry and Molecular Biology -5th Edition
- 4.Biological Variation:From principles to Practice-Callum G Fraser (AACC Press)

*Sample processed at SUBURBAN DIAGNOSTICS (INDIA) PVT. LTD CPL, Andheri West
*** End Of Report ***







Dr.ANUPA DIXIT
M.D.(PATH)
Consultant Pathologist & Lab
Director

Authenticity Check

Use a OR Code Scanner

Application To Scan the Code

ADDRESS: 2" Floor, Aston, Sundervan Complex, Above Mercedes Showroom, Andheri West - 400053

HEALTHLINE - MUMBAI: 022-6170-0000 | OTHER CITIES: 1800-266-4343

For Feedback - customerservice@suburbandiagnostics.com | www.suburbandiagnostics.com

Page 9 of 9