






r. PIYUSH GOY

Dr. PIYUSH GOYAL MBBS, DMRIN Radiologist) RMC No.-037041

P3 HEALTH SOLUTIONS LLP (ASSOCIATES OF MAXCARE DIAGNOSTICS) O R-14 Vidhyadhar Enclave-II. Near Avix Rank

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023





General Physical Examination

Date of Examination: 11/03/324	
Name: SHYOTT LALYADAY Age: 3	GYRS DOB: 10/06/1987Sex: Male
Referred By: UNTON BANK	· ·
Photo ID: ADDHARCARD ID#: 3509	
Ht: 174 (cm) Wt	: <u> </u>
Chest (Expiration):38 (cm) Ab	domen Circumference: (cm)
Blood Pressure: \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	RR: 18/min Temp: Alebrile
BMI_2_5	
Eye Examination: RIETCIG, NIG, I	1 CB
Other:	
On examination he/she appears physically and mentally f	it: Yes TNo
	me of Examinee: SHYOTT LAL YADAY
Signature Medical Examiner: (Radiologist) RMC No037041	Name Medical Examiner - PR-PTYUSH COYAI



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ② maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

Sex :- Male

Patient ID :-12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mi

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

HAEMATOLOGY

HAEMOGARAM			
HAEMOGLOBIN (Hb)	13.9	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	6.00	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	48.0	%	40.0 - 80.0
LYMPHOCYTE	43.0 H	%	20.0 - 40.0
EOSINOPHIL	2.0	%	1.0 - 6.0
MONOCYTE	7.0	%	2.0 - 10.0
BASOPHIL	0.0	%	0.0 - 2.0
TOTAL RED BLOOD CELL COUNT (RBC)	4.41 L	x10^6/uL	4.50 - 5.50
HEMATOCRIT (HCT)	43.20	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	98.0	fL	83.0 - 101.0
MEAN CORP HB (MCH)	31.6	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	32.3	g/dL	31.5 - 34.5
PLATELET COUNT	158	x10^3/uL	150 - 410
RDW-CV	13.7	%	11.6 - 14.0

Technologist RAVIMEENA Page No: 2 of 12



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
 □ maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Sex :-

Age :-

36 Yrs 9 Mon 2 Days

Male

Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp:-

Company:-

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

HAEMATOLOGY

Erythrocyte Sedimentation Rate (ESR)

10

mm in 1st hr

00 - 15

The erythrocyte sedimentation rate (ESR or sed rate) is a relatively simple, inexpensive, non-specific test that has been used for many years to help detect inflammation associated with conditions such as infections, cancers, and autoimmune diseases.ESR is said to be a non-specific test because an elevated result often indicates the presence of inflammation but does not tell the health practitioner exactly where the inflammation is in the body or what is causing it. An ESR can be affected by other conditions besides inflammation. For this reason, the ESR is typically used in conjunction with other tests, such as C-reactive protein.ESR is used to help diagnose certain specific inflammatory diseases, including temporal arteritis, systemic vasculitis and polymyalgia rheumatica. (For more on these, read the article on Vasculitis.) A significantly elevated ESR is one of the main test results used to support the diagnosis. This test may also be used to monitor disease activity and response to therapy in both of the above diseases as well as





(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885

⊕ maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

Sex :- Male

Patient ID :-12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

HAEMATOLOGY

(CBC): Methodology: TLC,DLC Fluorescent Flow cytometry, HB SLS method,TRBC,PCV,PLT Hydrodynamically focused Impedance and MCH,MCV,MCHC,MENTZER INDEX are calculated. InstrumentName: Sysmex 6 part fully automatic analyzer XN-L,Japan



RAVIMEENA Page No: 4 of 12



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885

⊕ maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

Sex :- Male

Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company:- Mr.

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

BIOCHEMISTRY

FASTING BLOOD SUGAR (Plasma)

Methord: - GOD POD

89.4

mg/dl

70.0 - 115.0

Impaired glucose tolerance (IGT)	111 - 125 mg/dL	- Z4
Diabetes Mellitus (DM)	> 126 mg/dL	

Instrument Name: HORIBA CA60 Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm,

hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin

therapy or various liver diseases.

BLOOD SUGAR PP (Plasma) Methord:- GOD PAP 113.0

mg/dl

70.0 - 140.0

Instrument Name: HORIBA Interpretation: Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.

Technologist RAVIMEENA Page No: 10 of 12



 B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

36 Yrs 9 Mon 2 Days Age :-

Sex :-Male Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :-Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

GLYCOSYLATED HEMOGLOBIN (HbA1C)

Test Name	Value	Unit	Biological Ref Interval
Methord:- CAPILLARY with EDTA	5.4	mg%	Non-Diabetic < 6.0 Good Control 6.0-7.0 Weak Control 7.0-8.0 Poor control > 8.0
MEAN PLASMA GLUCOSE Methord:- Calculated Parameter	106	mg/dL	68 - 125

INTERPRETATION

AS PER AMERICAN DIABETES ASSOCIATION (ADA)

Reference Group HbA1c in %

Non diabetic adults >=18 years < 5.7

At risk (Prediabetes) 5.7 - 6.4

Diagnosing Diabetes >= 6.5

CLINICAL NOTES

In vitro quantitative determination of HbA1c in whole blood is utilized in long term monitoring of glycemia. The HbA1c level correlates with the mean glucose concentration prevailing in the course of the patient's recent history (approx - 6-8 weeks) and therefore provides much more reliable information for glycemia monitoring than do determinations of blood glucose or urinary glucose. It is recommended that the determination of HbA1c be performed at intervals of 4-6 weeks during Diabetes Mellitus therapy. Results of HbA1c should be assessed in conjunction with the patient's medical history, clinical examinations and other findings. Some of the factors that influence HbA1c and its measurement [Adapted from Gallagher et al.]

- 1. Erythropoiesis
- Increased HbA1c; iron, vitamin B12 deficiency, decreased erythropolesis,
- Decreased HbA1c: administration of erythropoietin, iron, vitamin B12, reticulocytosis, chronic liver disease.
- 2. Altered Haemoglobin-Genetic or chemical alterations in hemoglobin: hemoglobinopathies, HbF, methemoglobin, may increase or decrease HbA1c.
- 3. Glycation
- Increased HbA1c: alcoholism, chronic renal failure, decreased intraerythrocytic pH.
- Decreased HbA1c; certain hemoglobinopathies, increased intra-erythrocyte pH
- .4. Erythrocyte destruction
- Increased HbA1c: increased erythrocyte life span: Splenectomy.
 Decreased A1c: decreased RBC life span: hemoglobinopathies, splenomegaly, rheumatoid arthritis or drugs such as antiretrovirals, ribavirin & dapsone.

- Increased HbA1c: hyperbilirubinemia, carbamylated hemoglobin, alcoholism, large doses of aspirin, chronic opiate use, chronic renal failure
 Decreased HbA1c: hypertriglyceridemia, reticulocytosis, chronic liver disease, aspirin, vitamin C and E, splenomegaly, rheumatoid arthritis or drugs



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ② maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

Sex :- Male

Patient ID :-12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

HAEMATOLOGY

BLOOD GROUP ABO Methord:- Haemagglutination reaction

"B" POSITIVE



Technologist RAVIMEENA Page No: 5 of 12



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⑥ +91 141 4824885 ② maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

30 115 9 WOII 2 D

Sex :- Male

Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- 1

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

Test Name	Value	Unit	Biological Ref Interval
RFT / KFT WITH ELECTROLYTES			
SERUM UREA Methord:- Urease/GLDH	33.20	mg/dl	10.00 - 50.00
SERUM CREATININE Methord:- Jaffe's Method	1.17	mg/dl	Males : 0.6-1.50 mg/dl Females : 0.6 -1.40 mg/dl
SERUM URIC ACID	6.05	mg/dl	2.40 - 7.00
SODIUM Methord:- ISE Interpretation:	139.6	mmol/L	135.0 - 150.0

Electrolytes are minerals that are found in body tissues and blood in the form of dissolved salts. As electrically charged particles, electrolytes help move nutrients into and wastes out of the body's cells, maintain a healthy water balance, and help stabilize the body's acid/base (pH) level. The electrolyte panel measures the blood levels of the main electrolytes in the body:

POTASSIUM

Methord:- ISE

4.15

mmol/L

3.50 - 5.50

CHLORIDE

102.2

mmol/L

94.0 - 110.0

Methord:-ISE

* Chloride—this electrolyte moves in and out of the cells to help maintain electrical neutrality (concentrations of positively charged cations and negatively charged anions must be equal) and its level usually mirrors that of sodium. Due to its close association with sodium, chloride also helps to regulate the distribution of water in the body

SERUM CALCIUM Methord:- Arsenazo III Method	9.56	mg/dL	8.80 - 10.20
SERUM TOTAL PROTEIN Methord:- Direct Biuret Reagent	6.68	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- Bromocresol Green	4.21	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.47	gm/dl	2.20 - 3.50
A/G RATIO	1.70		1.30 - 2.50

RAVIMEENA Page No: 6 of 12 DR.TANU RUNGTA

MD (Pathology) RMC No. 17226

^{*} Sodium—most of the body's sodium is found in the fluid outside of the body's cells, where it helps to regulate the amount of water in the body.

^{* &}lt;u>Potassium</u>—this electrolyte is found mainly inside the body's cells. A small but vital amount of potassium is found in the plasma, the liquid portion of the blood. Potassium plays an important role in regulating muscle contraction. <u>Monitoring potassium</u> is important as small changes in the potassium level can affect the heart's rhythm and ability to contract



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

(8) +91 141 4824885 (2) maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

Sex :- Male

Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :- Mr.N

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

BIOCHEMISTRY

INTERPRETATION

Kidney function tests are group of tests that can be used to evaluate how well the kidneys are functioning. Creatinine is a waste product that comes from protein in the diet and also comes from the normal wear and tear of muscles of the body. In blood, it is a marker of GFR in urine, it can remove the need for 24-hourcollections for many analytes or be used as a quality assurance tool to assess the accuracy of a 24-hour collection Higher levels may be a sign that the kidneys are not working properly. As kidney disease progresses, the level of creatinine and urea in the bloodincreases. Certain drugs are nephrotoxic hence KFT is done before and after initiation of treatment with these drugs.

Low serum creatinine values are rare; they almost always reflect low muscle mass.

Apart from renal failure Blood Urea can increase in dehvdration and GI bleed



Technologist



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885

maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

36 Yrs 9 Mon 2 Days Age :-

Sex :-Male Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company:-

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

Borderline high 150-199

Near Optimal/above optimal

Borderline High 130-159 High 160-189

<150

200-499

>500

Desirable <200 Borderline 200-239 High> 240 Normal

High

Very high

MALE- 30-70 **FEMALE - 30-85** Optimal <100

Very High > 190 0.00 - 80.00

100-129

0.00 - 4.90

0.00 - 3.50

400.00 - 1000.00

BIOCHEMISTRY

LIPID PROFILE

TOTAL CHOLESTEROL Methord:- CHOD-PAP methodology	173.00	mg/dl
TRIGLYCERIDES Methord:- GPO-PAP	198.00 H	mg/dl
DIRECT HDL CHOLESTEROL Methord:- Direct clearance Method	43.60	mg/dl
LDL CHOLESTEROL Methord:- Calculated Method	96.40	mg/dl
VLDL CHOLESTEROL Methord:- Calculated	39.60	mg/dl
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Methord:- Calculated	3.97	
LDL / HDL CHOLESTEROL RATIO Methord:- Calculated	2.21	
TOTAL LIPID	608.15	mg/dl

608.15 mg/dl 1 Measurements in the same patient can show physiological& analytical variations. Three serialsamples 1 week apart are recommended for

DR.TANU RUNGTA

MD (Pathology) RMC No. 17226

Methord:- CALCULATED

Total Cholesterol, Triglycerides, HDL& LDL Cholesterol. 2. As per NCEP guidelines, all adults above the age of 20 years should be screened for lipid status. Selective screening of children above the age of 2 years with a family history of premature cardiovascular disease or those with at least one parent with high total cholesterol is recommended

^{3.} Low HDL levels are associated with Coronary Heart Disease due to insufficient HDL being available to participate in reverse cholesterol transport, the process by which cholesterol is eliminated fromperipheral tissues



O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age :-

36 Yrs 9 Mon 2 Days

Sex :-Male Patient ID: -12234834

Date: - 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-Company :-

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

BIOCHEMISTRY

LIVER FUNCTION TEST

SERUM BILIRUBIN (TOTAL) Methord:- DMSO/Diazo	1.18	mg/dL	Infants: 0.2-8.0 mg/dL Adult - Up to - 1.2 mg/dL
SERUM BILIRUBIN (DIRECT) Methord:- DMSO/Diazo	0.38	mg/dL	Up to 0.40 mg/dL
SERUM BILIRUBIN (INDIRECT) Methord:- Calculated	0.80	mg/dl	0.30-0.70
SGOT Methord:- IFCC	25.6	U/L	0.0 - 40.0
SGPT Methord:- IFCC	33.2	U/L	0.0 - 40.0
SERUM ALKALINE PHOSPHATASE Methord:- DGKC - SCE	95.60	U/L	53.00 - 141.00
SERUM TOTAL PROTEIN Methord:- Direct Biuret Reagent	6.68	g/dl	6.00 - 8.40
SERUM ALBUMIN Methord:- Bromocresol Green	4.21	g/dl	3.50 - 5.50
SERUM GLOBULIN Methord:- CALCULATION	2.47	gm/dl	2.20 - 3.50
A/G RATIO	1.70		1.30 - 2.50

Note: These are group of tests that can be used to detect the presence of liver disease, distinguish among different types of liver disorders, gauge the extent of known liver damage, and monitor the response to treatment. Most liver diseases cause only mild symptoms initially, but these diseases must be detected early. Some tests are associated with functionality (e.g., albumin), some with cellular integrity (e.g., transaminase), and some with conditions linked to the biliary tract (gamma-glutamyl transferase and alkaline phosphatase). Conditions with elevated levels of ALT and AST include hepatitis A,B,C ,paracetamol toxicity etc. Several biochemical tests are useful in the evaluation and management of patients with hepatic dysfunction. Some or all of these measurements are also carried out (usually about twice a year for routine cases) on those individuals taking certain medications, such as anticonvulsants, to ensure that the medications are not adversely impacting the person's liver.



B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
 □ maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age :-36 Yrs 9 Mon 2 Days

Sex :-Male Patient ID: -12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

			1 inal Authentication . 11/03/2024 17:44:07
Test Name	Value	Unit	Biological Ref Interval
TOTAL THYROID PROFILE			4
THYROID-TRIIODOTHYRONINE T3 Methord:- ECLIA	0.83	ng/mL	0.70 - 2.04
THYROID - THYROXINE (T4) Methord:- ECLIA	5.82	ug/dl	5.10 - 14.10
TSH Methord:- ECLIA	2.111	μIU/mL	0.350 - 5.500

4th Generation Assay, Reference ranges vary between laboratories

PREGNANCY - REFERENCE RANGE for TSH IN ulU/mL (As per American Thyroid Association)

1st Trimester: 0.10-2.50 uIU/mL 2nd Trimester: 0.20-3.00 uIU/mL 3rd Trimester: 0.30-3.00 uIU/mL

The production, circulation, and disintegration of thyroid hormones are altered throughout the stages of pregnancy.

NOTE-TSH levels are subject to circardian variation, reaching peak levels between 2-4 AM and min between 6-10 PM. The variation is the order of 50% hence time of the day has influence on the measures serum TSH concentration. Dose and time of drug intake also influence the test result.

INTERPRETATION

- 1.Primary hyperthyroidism is accompanied by †serum T3 & T4 values along with ↓ TSH level.
- 2. Primary hypothyroidism is accompanied by 1 serum T3 and T4 values & †serum TSH levels
- 3.Normal T4 levels accompanied by † T3 levels and low TSH are seen in patients with T3 Thyrotoxicosis
- 4.Normal or ↓ T3 & ↑T4 levels indicate T4 Thyrotoxicosis (problem is conversion of T4 to T3)
- 5.Normal T3 & T4 along with \ TSH indicate mild / Subclinical Hyperthyroidism
- . COMMENTS: Assay results should be interpreted in context to the clinical condition and associated results of other investigations. Previous treatment with corticosteroid therapy may result in lower TSH levels while thyroid hormone levels are normal. Results are invalidated if the client has undergone a radionuclide scan within 7-14 days before the test.
- Disclaimer-TSH is an important marker for the diagnosis of thyroid dysfunction. Recent studies have shown that the TSH distribution progressively shifts to a higher concentration with age ,and it is debatable whether this is due to a real change with age or an increasing proportion of unrecognized thyroid disease in the elderly
- . Reference ranges are from Teitz fundamental of clinical chemistry 8th ed (2018

Test performed by Instrument: Beckman coulter Dxi 800

Note: The result obtained relate only to the sample given/ received & tested. A single test result is not always indicative of a disease, it has to be correlated with clinical data for interpretation.



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

 B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885
 □ maxcarediagnostics1@gmail.com





NAME :- Mr. SHYOJI LAL YADAV

Age:- 36 Yrs 9 Mon 2 Days

Sex :- Male

Patient ID :-12234834

Date :- 11/03/2024

10:28:24

Ref. By Doctor:-UNION BANK

Lab/Hosp :-

Company :-

Mr.MEDIWHEEL

Final Authentication: 11/03/2024 17:44:07

TD-4 NJ	Value	TT-:4	Dialogical Definternal
Test Name	Value	Unit	Biological Ref Interval
Urine Routine PHYSICAL EXAMINATION			*
	PALE YELL	OW	PALE YELLOW
COLOUR		JOW	
APPEARANCE	Clear		Clear
CHEMICAL EXAMINATION			
REACTION(PH)	5.5		5.0 - 7.5
SPECIFIC GRAVITY	1.015		1.010 - 1.030
PROTEIN	NIL	Arthurston	NIL
SUGAR	NIL		NIL
BILIRUBIN	NEGATIVE		NEGATIVE
UROBILINOGEN	NORMAL	· A	NORMAL
KETONES	NEGATIVE		NEGATIVE
NITRITE	NEGATIVE		NEGATIVE
MICROSCOPY EXAMINATION			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT	3	ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		
Complete Control and Control a			

*** End of Report ***

Technologist



(ASSOCIATES OF MAXCARE DIAGNOSTICS)

O B-14, Vidhyadhar Enclave-II, Near Axix Bank Central Spine, Vidhyadhar Nagar, Jaipur - 302023

⊕ +91 141 4824885 maxcarediagnostics1@gmail.com



NAME:	MR. SHYOJI LAL YADAV	18.20 1 .20 1.20 1.20 1.20 1.20 1.20 1.20 1.20 1	AGE	36 YRS/M
REF.BY	UNION BANK	•	DATE	11/03/2024

CHEST X RAY (PA VIEW)

Bilateral lung fields appear clear.

Bilateral costo-phrenic angles appear clear.

Cardiothoracic ratio is normal.

Thoracic soft tissue and skeletal system appear unremarkable.

Soft tissue shadows appear normal.

IMPRESSION: No significant abnormality is detected.

Shallni !

DR.SHALINI GOEL


M.B.B.S, D.N.B (Radiodiagnosis)

RMC No.: 21954


ef.: UNION BANK Test Date: 11-Mar-2024(11:29:34) Notch: 50Hz 0.05Hz - 35Hz 10mm/m/ יems (א) בגם P3 HEALTH SOLUTIONS LLP B-14, Vidhyadhar nahar , Jaipur P-QRS-T axis: 65. -7. 58. (Deg) Comments: Vent Rate: 76 bpm; PR Interval: 120 ms; QRS Duration: 98 ms; FINDINGS: Normal Sinus Rhythm avR 12 10mm/mV QT/QTc Int: 335/378 ms BP: 25mm/Sec ... MBBS, DIP. CARDIO (ESCORTS) Dr. Naresh Kumar Wohanka mmHg RMC No.: 35703 HR: 76 bpm ٧6 **√**5 4 S

PR Interval: 120 ms
QRS Duration: 98 ms
QT/QTc: 335/378ms
P-QRS-T Axis: 65 - -7 - 58 (Deg)





DOMESTIC DESCRIPTION OF 100 1500



पुरुष/ MALE

मेरा आधार, मेरी पहचान

Issue Date: 19/03/2018

