

CERTIFICATE OF MEDICAL FITNESS
NAME: Moinal fulchas
AGE/GENDER: 354 Female
HEIGHT: 160cm WEIGHT: 71.4 Cg
IDENTIFICATION MARK: Bleucle mole on the upperly
BLOOD PRESSURE: 120 180 mm+19
PULSE: 746lm
CVS: Normal
ANY OTHER DISEASE DIAGNOSED IN THE PAST: Thyroid Thyronorm 150 mg
ALLERGIES, IF ANY:
LIST OF PRESCRIBED MEDICINES: Ni
ANY OTHER REMARKS: NO
Certify that I have carefully examined Mr/Mrs. Wind fulckar son/daughter of Mr Basanth fulckar who has signed in my presence. He/ she has no physical disease and is fit for employment.
Dr. BINDURAJ. R
Signature of Medical Officer
Place: Spectrum Daynortius & health care
Date: 09/08/04

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





Dr. Ashok S Bsc., MBBS., D.O.M.S Consultant Opthalmologist KMC No: 31827

DATE: 09-03-24

EYE EXAMINATION

		-11 /	-140	_ ^		_
NAME:	8- M	rinal 1	meg g	AGE: 35	GENDER:	F/M

RIGHT EYE LEFT EYE 616100 Bloins Vision Vision With glass Color Vision Normal Normal Anterior segment examination Normal Normal **Fundus Examination** Normal Normal Any other abnormality Nill Nill Diagnosis/ impression Normal Normal

Consultant (Opthalmologist)







NAME	AGE	GENDER	
Mx. Moral Fulekon	35771	Femile.	

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY -> None.

M: MISSING -9 None.

O: OTHERS

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:


SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR B.D.S., F.A.G.E., F.P.F.A. (USA) Reg. No: 2247/A





A Comment, March Marghall Business of Street and Supersonal Comments of the Comment of the Comme	



NAME	: MRS.MRINAL FULEKAR	DATE : 09/03/2024
AGE/SEX	: 35YEARS/FEMALE	REG NO: 0903240068
REF BY	: APOLLO CLINIC	

CHEST PA VIEW

- Visualised lungs are clear.
- Bilateral hila appears normal.
- Cardia is normal in size
- · No pleural effusion

IMPRESSION: No significant abnormality .

DR PRAVEEN B, DMRD, DNB **Consultant Radiologist**







SPECTRUM DIAGNOSTICS

Bangalore

Patient ID: 0228


Name: MRINAL FULEKAR CR Number: 20240309135655 Age: 35

Gender: Female

37,50 dB

35.00 dB

Operator: spectrum diagnostics



CI	les l	and	M	-	tes	٠
•	ш	Gall	-13	o	ue 5	¥

AIR Right

35.91 dB

Not Found

36.25 dB

36.67 dB



PATIENT NAME	MRS MRINAL FULEKAR	ID NO	0903240068
AGE	35YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	09.03.2024

2D ECHO CARDIOGRAHIC STUDY

IV	I-IVIODE	
AORTA	26mm	
LEFT ATRIUM	38mm	
RIGHT VENTRICLE	20mm	
LEFT VENTRICLE (DIASTOLE)	45mm	
LEFT VENTRICLE(SYSTOLE)	27mm	
VENTRICULAR SEPTUM (DIASTOLE)	08mm	
VENTRICULAR SEPTUM (SYSTOLE)	10mm	
POSTERIOR WALL (DIASTOLE)	12mm	
POSTERIOR WALL (SYSTOLE)	11mm	
FRACTIONAL SHORTENING	30%	
EJECTION FRACTION	60%	

DOPPLER / COLOUR FLOW

Mitral Valve Velocity : MVE- 0.84m/s MVA - 0.63m/s E/A-0.64

Tissue Doppler : e' (Septal) -10cm/s E/e'(Septal) -8

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve: 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 2.27 m/s 19mmHg







PATIENT NAME	MRS MRINAL FULEKAR	ID NO	0903240068
AGE	35YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	09.03.2024

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL	
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA	

RIGHT VENTRICLE	;	NORMAL	
LEFT ATRIUM	:	NORMAL	
RIGHT ATRIUM		NORMAL	
MITRAL VALVE	:	NORMAL	
AORTIC VALVE	:	NORMAL	
PULMONARY VALVE	:	NORMAL	
TRICUSPID VALVE	:	NORMAL	
INTER ATRIAL SEPTUM	:	INTACT	
INTER VENTRICULAR SEPTL	JM:	INTACT	
PERICARDIUM	:	NORMAL	
OTHERS	:	- NIL	

IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- NORMAL LV FUNCTION, LVEF- 60%
- TRIVIAL MR / TRIVIAL TR / TRIVIAL PAH
- NO CLOT / VEGETATION / EFFUSION

ECHO TECHNICIAN

The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correction.





NAME AND LAB NO	MRS MRINAL FULEKAR	REG -40068
AGE & SEX	35 YRS	FEMALE
DATE AND AREA OF INTEREST	09.03.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

LIVER:

Normal in size and shows diffuse increased echogenicity.

No e/o IHBR dilatation. No evidence of focal lesion

Portal vein appears normal.

CBD appears normal.

GALL BLADDER:

Partially distended . No obvious calculus in the visualised luminal portion.

SPLEEN:

Normal in size and echotexture. No focal lesion

PANCREAS:

Head and body appears normal . Tail obscured by bowel gas shadows

RETROPERITONEUM:

Suboptimal visualised due to bowel gas.

RIGHT KIDNEY:

Measures 11.9 x1.5 cm Right kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

LEFT KIDNEY:

Measures 10.8 X1.6 cm .Left kidney is normal in size & echotexture

No evidence of calculus/ hydronephrosis.

URINARY BLADDER:

Well distended. No wall thickening/ calculi.

UTERUS:

Anteverted, Normal in size 7.6 X3.8 X4.2 cm and echotexture

Endometrium is normal, ET - 9.3 mm.

OVARIES:

B/L ovaries normal in size and echotexture.

RO - 3.3 X1.4cm,

LO -4.0 X1.9 cm - Dominant follicle noted measuring 13.4 x10.0 mm

No obvious adnexal mass lesions.

No evidence of ascites/pleural effusion.

IMPRESSION:

Grade I fatty liver.

Suggested clinical / lab correlation.

EN B, DMRD, DNB CONSULTANT RADIOLOGIST



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555

info@spectrumdiagnostics.org

www.spectrumdiagnostics.org





Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068 C/o

: Apollo Clinic

Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date : 09-Mar-2024 02:39 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole E	Blood EDTA			- 1 1 1 1 1 2
Haemoglobin (HB)	13.60	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotmeter
Red Blood Cell (RBC)	4.91	million/cun	nm3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	39.90	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	81.30	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	27.60	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	33.90	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	42.70	fL	40.0-55.0	Volumetric
Red Blood Cell Distribution CV (RDW-CV)	16.20	%	Male: 11.80-14.50 Female:12.20-16.10	Impedance Volumetric Impedance
Mean Platelet Volume (MPV)	9.00	fL	8.0-15.0	Volumetric
Platelet	3.35	lakh/cumm	1.50-4.50	Impedance Volumetric
Platelet Distribution Width (PDW)	13.60	%	8.30 - 56.60	Impedance Volumetric Impedance
White Blood cell Count (WBC)	10710.00	cells/cumm	Male: 4000-11000 Female 4000-11000 Children: 6000-17500 Infants: 9000-30000	Volumetric Impedance
Veutrophils	70.0	%	40.0-75.0	Light
ymphocytes	23.0	%	20.0-40.0	scattering/Manual Light
osinophils	3.0	%	0.0-8.0	scattering/Manual Light scattering/Manual

UHID

: 0903240068

0903240068







Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068 C/o : Apollo Clinic Bill Date

: 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date : 09-Mar-2024 02:39 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Monocytes	3.0	%	0.0-10.0	Light scattering/Manua
Basophils	1.0	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	7.73	10^3/uL	2.0-7.0	Calculated
Absolute Lymphocyte Count	2.31	10^3/uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.34	10^3/uL	0.20-1.00	Calculated
Absolute Eosinophil Count	320.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.01	10^3/uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	13	mm/hr	Female: 0.0-20.0 Male: 0.0-10.0	Westergren

: 0903240068

0903240068

UHID

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : Normocytic Normochromic.

: Are normal in total number, morphology and distribution. WBC'S Platelets

: Adequate in number and normal in morphology. No abnormal cells or hemoparasites are present.

Impression: Normocytic Normochromic Blood picture.



Printed By

: spectrum

Printed On

+91 77604 97644 | 080 2337 1555

: 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalung 2560010 info@spectrumdiagnostics.org www.spectrumdiagnostics.org





Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068 C/o : Apollo Clinic UHID : 0903240068

0903240068 Report Status : Final

Bill Date : 09-Mar-2024 10:02 AM Sample Col. Date: 09-Mar-2024 10:02 AM

Result Date : 09-Mar-2024 02:39 PM

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serui	n		4	
Bilirubin Total-Serum	0.91	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.15	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.76	mg/dL	Female: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	19.00	U/L	Female: 15.0 - 37.0	UV with Pyridoxal - 5 -
Alanine Aminotransferase (ALT/SGPT)-Serum	20.00	U/L	Female: 14.0 - 59.0	Phosphate UV with Pyridoxal - 5 -
Alkaline Phosphatase (ALP)- Serum	104.00	U/L	Female: 45.0 - 117.0	Phosphate PNPP,AMP- Buffer
Protein, Total-Serum	7.26	g/dL	6.40-8.20	Biuret/Endpoint- With Blank
Albumin-Serum	4.31	g/dL	Female: 3.40 - 5.50	Bromocresol
Globulin-Serum	2.95	g/dL	2.0-3.50	Purple Calculated
Albumin/Globulin Ratio-Serum	1.46	Ratio	0.80-2.0	Calculated





Printed By


: spectrum

Printed On : 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist











Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068

C/o : Apollo Clinic Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date

: 09-Mar-2024 02:39 PM Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	239.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	162.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	61.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	178	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Scrum	160.0	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol
Very-low-density lipoprotein (VLDL) cholesterol-Serum	32	mg/dL	Female: 0.0 - 40	oxidase Calculated
Cholesterol/HDL Ratio-Serum	3.92	Ratio	Female: 0.0 - 5.0	Calculated
nterpretations				

: 0903240068

0903240068

UHID

Interpretation:

Desirable	Borderline High	Ulah	W VIII. 4
<200	-		Very High
<150	200000000	170.00	>500
<130	F. F. S. S. S.		>220
<100		100,000,000	>190
	<200 <150 <130	<200 200-239 <150 150-199 <130 160-189	<200

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atheroselerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By

: spectrum

Printed On

: 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru 560010 +91 77604 97644 | 080 2337 1555 Info@spectrumdiagnostics.org www.spectrumdiagnostics.org







Age / Gender : 35 years / Female Ref. By Dr.

Reg. No. : 0903240068 C/o : Apollo Clinic

UHID : 0903240068 : Dr. APOLO CLINIC

0903240068

Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date : 09-Mar-2024 02:39 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test)	:			
Blood Urea Nitrogen (BUN)- Serum	8.70	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.63	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	6.47	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	138.80	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.09	mmol/L	3.5 to 5.5	Ion-Selective Electrodes
Chloride(Cl-)-Serum	97.40	mmol/L	96.0-108.0	(ISE) Ion-Selective Electrodes (ISE)


Comments: Renal Function Test (RFT), also called kidney function tests, are a group of tests performed to evaluate the functions of the kidneys. The kidneys play a vital role in removing waste, toxins, and extra water from the body. They are responsible for maintaining a healthy balance of water, salts, and minerals such as calcium, sodium, potassium, and phosphorus. They are also essential for blood pressure control, maintenance of the body's pH balance, making red blood cell production hormones, and promoting bone health. Hence, keeping your kidneys healthy is essential for maintaining overall health. It helps diagnose inflammation, infection or damage in the kidneys. The test measures Uric Acid, Creatinine, BUN and electrolytes in the blood to determine the health of the kidneys. Risk factors for kidney dysfunction such as hypertension, diabetes, cardiovascular disease, obesity, elevated cholesterol or a family history of kidney disease. It may also be when has signs and symptoms of kidney disease, though in early stage often no noticeable symptoms are observed. Kidney panel is useful for general health screening; screening patients at risk of developing kidney disease; management of patients with known kidney disease. Estimated GFR is especially important in CKD patients CKD for monitoring, it helps to identify disease at early stage in those with risk factors for CKD (diabetes, hypertension, cardiovascular disease, and family history of kidney disease). Early recognition and intervention are important in slowing the progression of CKD and preventing its complications.



Printed By : spectrum

Printed On : 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist







Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068 C/o : Apollo Clinic

Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM

Result Date : 09-Mar-2024 02:39 PM Report Status : Final

Test Name Result Unit Reference Value Method Fasting Blood Sugar (FBS)-93 mg/dL 60.0-110.0 Hexo Kinase Plasma

0903240068

: 0903240068

UHID

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula C6H12O6. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high.Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes: Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol , Dietary - Intake of excessive carbohydrates and foods with high glycemic index? Exercise in between samples? Family history of Diabetes, Idiopathic, Partial / Total



Printed By


: spectrum


Printed On

: 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalurus 550010 +91 77604 97644 | 080 2337 1555











: MRS. MRINAL FULEKAR Name

Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068

C/o : Apollo Clinic Bill Date

: 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM

Result Date

: 09-Mar-2024 02:39 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA				
Glycosylated Haemoglobin	4.90	%	Non diabetic adults:<5.7	HPLC
(HbA1c)			At risk (Prediabetes): 5.7 - 6.4	
			Diagnosing Diabetes :>= 6.5	
			Diabetes	
			Excellent Control: 6-7	
			Fair to good Control: 7-8	
			Unsatisfactory Control:8-10	
			Poor Control :>10	
Estimated Average Glucose(eAG)	93.93	mg/dL		Calculated

: 0903240068

0903240068

UHID

Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

 Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1e provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemie control as compared to blood and urinary glucose determinations.



Printed By

: spectrum

Printed On

+91 77604 97644 | 080 2337 1555

: 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengalurus 700000 info@spectrumdiagnostics.org www.spectrumdiagnostics.org







Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068

C/o : Apollo Clinic Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM

Result Date : 09-Mar-2024 02:39 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TF7 Serum	Γ)-			
Tri-Iodo Thyronine (T3)-So	erum 0.77	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	7.80	μg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormo TSH)-Serum	one 4.91	μIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

0903240068

: 0903240068

UHID

Comments: Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population withouta clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester: 0.81-1.90, Second Trimester: 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG). Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments: Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males: 4.6-10.5, Females: 5.5-11.0, 60 Years: 5.0-10.70, Cord: 7.40-13.10, Children: 1-3 Days: 11.80-22.60, 1-2 Weeks: 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years: 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5,6 Days :>6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy. Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments: TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subelinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance. Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.




Printed By : spectrum


Printed On : 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru 560010













Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

: 0903240068

C/o : Apollo Clinic

Reg. No.

Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date : 09-Mar-2024 02:39 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Calcium, Total- Serum	9.40	mg/dL	8.50-10.10	Spectrophotometry (O- Cresolphthalein complexone)
Gamma-Glutamyl Transferase (GGT)-Serum	24.00	U/L	Male: 15.0-85.0	Other g-Glut-3- carboxy-4 nitro
			Female: 5.0-55.0	carboxy=4 miro

0903240068

: 0903240068

UHID

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of scrum GGT levels in detecting alcohol-induced liver disease. Elevated scrum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By

: spectrum

Printed On

: 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist

www.spectrumdiagnostics.org



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org





Name

: MRS. MRINAL FULEKAR

Age / Gender Ref. By Dr.

: 35 years / Female : Dr. APOLO CLINIC

Reg. No.

: 0903240068

C/o

: Apollo Clinic

UHID

: 0903240068

Bill Date

: 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM

Result Date

: 09-Mar-2024 03:18 PM

Report Status

: Final

Test Name

Result

Positive

Unit

Reference Value

Method

Blood Group & Rh Typing-Whole Blood EDTA

Blood Group

Rh Type

Slide/Tube

agglutination

Slide/Tube

agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (crythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type



Printed By

: spectrum

Printed On

: 09 Mar, 2024 06:03 pm

Dr. Nithun Roddy C,MD,Consultant Pathologist

Page 10 of 12 Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org



www.spectrumdiagnostics.org



Other Branch: #486/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 🌑 +91 6361 253 097 | 080-2991 6944 | 080-49511985





Age / Gender : 35 years / Female

Ref. By Dr. : Dr. APOLO CLINIC Reg. No. : 0903240068

C/o : Apollo Clinic

Bill Date : 09-Mar-2024 10:02 AM : 0903240068

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date : 09-Mar-2024 03:18 PM

Report Status : Final

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-I	Jrine .			
Physical Examination			*	
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	
Reaction (pH)	6.0		5.0-7.5	Visual
Specific Gravity	1.020		1.000-1.030	Dipstick
Biochemical Examination	107/80207		1.000-1.030	Dipstick
Albumin	Negative		Negative	B:
Glucose	Negative		Negative	Dipstick/Precipitation
Bilirubin	Negative		Negative	Dipstick/Benedicts
Ketone Bodies	Negative			Dipstick/Fouchets
Urobilinogen	Normal		Negative Normal	Dipstick/Rotheras
Nitrite	Negative			Dipstick/Ehrlichs
Microscopic Examination	riogutivo		Negative	Dipstick
Pus Cells	2-4	hpf	0050	
Epithelial Cells	1-2	100	0.0-5.0	Microscopy
RBCs	Absent	hpf	0.0-10.0	Microscopy
Casts	Absent	hpf	Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy
50000000000000000000000000000000000000	Auscill		Absent	Microscopy

UHID

0903240068

Comments: The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections,



Printed By

: spectrum


Printed On


: 09 Mar, 2024 06:03 pm

Dr. Nithun Reddy C,MD,Consultant Pathologist



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010 +91 77604 97644 | 080 2337 1555









Age / Gender : 35 years / Female Ref. By Dr. : Dr. APOLO CLINIC

Reg. No. : 0903240068 C/o : Apollo Clinic UHID : 0903240068

0903240068

Bill Date : 09-Mar-2024 10:02 AM

Sample Col. Date: 09-Mar-2024 10:02 AM Result Date : 09-Mar-2024 04:18 PM

Report Status : Final

Test Name Result Unit Reference Value Method

Fasting Urine Glucose-Urine Negative Negative Dipstick/Benedicts (Manual)



Printed By Printed On

: spectrum

: 09 Mar, 2024 06:03 pm

1. Ilum

Dr. Nithun Reddy C,MD,Consultant Pathologist



