



# SPECTRUM

DIAGNOSTICS & HEALTH CARE

## CERTIFICATE OF MEDICAL FITNESS

NAME: Ravikumar

AGE/ GENDER: 59y / m

HEIGHT: 162cm

WEIGHT: 62.3kg

IDENTIFICATION MARK: \_\_\_\_\_

BLOOD PRESSURE: 120/90 mmHg

PULSE: 75/min

CVS: Normal

RS:P

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Nil

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: Nil

ANY OTHER REMARKS: Nil

I Certify that I have carefully examined Mr/Mrs. Ravikumar son/daughter of Ms Adhinarayana Shastri who has signed in my presence. He/ she has no physical disease and is fit for employment.

Ravikumar

Signature of candidate

Signature of Medical Officer

Dr. BINDURA L. R  
Internal Medicine

Place: Spectrum Diagnostics & Health Care

Date: 27/12/23

**Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined**

SCAN FOR LOCATION



Dr. Ashok S  
Bsc., MBBS., D.O.M.S  
Consultant Ophthalmologist  
KMC No: 31827

DATE: 26.12.23

**EYE EXAMINATION**

NAME: *ms. Ravi Kumar*      AGE: *59y*      GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	<i>6/36: 2/10</i>	<i>ND PL</i> <i>6/36: 2/10</i>
Vision With glass	<i>6/6</i>	<i>6/6 ND PL</i>
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	<i>R-D, R-Ad</i> Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ impression	Normal	Normal

Consultant (Ophthalmologist)  
**Dr. ASHOK SARODHE**  
EYE CLINIC



NAME	AGE	GENDER
Mr. Parthasarathy - A	59 yrs	Male

**DENTAL EXAMINATION REPORT:**

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

- C: CAVITY** → Dental Caries on  $\frac{1}{6}$ ; needs restoration.
- M: MISSING** →  $\frac{1}{6}$ ; needs replacement.
- O: OTHERS** → Severe submucous fibrosis (??)

**ADVISED:**

CLEANING /  SCALING /  ROOTS PLANNING /  FLOSSING & POLISHING /  OTHERS

**REMARKS:**

SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

27/12/2023

**Dr. SACHDEV NAGARKAR**  
B.D.S., F.A.G.E., F.P.F.A. (USA)  
Reg. No : 2247/A






MR RAVIKUMARA  
Male 59Years

HR : 73 bpm  
P : 92 ms  
PR : 135 ms  
QRS : 95 ms  
QT/QTc : 419/463 ms  
P/QRS/T : 42/-9/-10 °  
RV5/SV1 : 1.736/0.937 mV

Diagnosis Information:  
Sinus Rhythm  
Largd PtfV1

Report Confirmed by:





<b>NAME : MR. RAVIKUMAR A</b>	<b>DATE :27/12/2023</b>
<b>AGE/SEX : 59 YEARS/MALE</b>	<b>REG NO:2230016</b>
<b>REF BY : APOLO CLINIC</b>	

## ***CHEST PA VIEW***


Lung fields are clear.

Cardiovascular shadows are within normal limits.

Both CP angles are free.

Domes of diaphragm and bony thoracic cage are normal.

**IMPRESSION: NORMAL CHEST RADIOGRAPH.**



**DR.RAM PRAKASH G MDRD  
CONSULTANT RADIOLOGIST**

*Your suggestion / feedback is a valuable input for improving our services*







# SPECTRUM DIAGNOSTICS

Bangalore

Patient ID : 0021  
 Name : RAVIKUMAR A  
 CR Number : 20231227113907  
 Registration Date : 27-Dec-2023

Age : 59  
 Gender : Male  
 Operator : spectrum diagnostics



	125 Hz	250 Hz	500 Hz	750 Hz	1000 H	1500 H	2000 H	3000 H	4000 H	6000 H	8000 H
X - Air Left	30	35	35	30	30	30	25	30	45	45	30
O - Air Right	30	30	35	35	35	35	30	35	40	45	50
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	33.18 dB	37.50 dB	28.33 dB	32.50 dB
AIR Right	36.36 dB	42.50 dB	33.33 dB	32.50 dB

**Clinical Notes :**

Right Ear: Moderate  
 Left Ear: Mild



PATIENT NAME	MR RAVIKUMAR A	ID NO	2712230016
AGE	59YEARS	SEX	MALE
REF BY	DR.APOLO CLINIC	DATE	27.12.2023

**2D ECHO CARDIOGRAHIC STUDY**

**M-MODE**

AORTA	36mm
LEFT ATRIUM	31mm
RIGHT VENTRICLE	20mm
LEFT VENTRICLE (DIASTOLE )	42mm
LEFT VENTRICLE(SYSTOLE)	27mm
VENTRICULAR SEPTUM (DIASTOLE)	12mm
VENTRICULAR SEPTUM (SYSTOLE)	13mm
POSTERIOR WALL (DIASTOLE)	12mm
POSTERIOR WALL (SYSTOLE)	11mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	58%

**DOPPLER /COLOUR FLOW**

Mitral Valve Velocity : MVE- 0.40/s MVA – 0.63m/s E/A-0.64

Tissue Doppler : e' ( Septal) -10cm/s E/e'(Septal) -4

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve : 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 2.60m/s 30mmHg





PATIENT NAME	MR RAVIKUMAR A	ID NO	2712230016
AGE	59YEARS	SEX	MALE
REF BY	DR.APOLO CLINIC	DATE	27.12.2023

### **2D ECHO CARDIOGRAHIC STUDY**

LEFT VENTRICLE	SIZE& THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE	: NORMAL
LEFT ATRIUM	: NORMAL
RIGHT ATRIUM	: NORMAL
MITRAL VALVE	: NORMAL
AORTIC VALVE	: NORMAL
PULMONARY VALVE	: NORMAL
TRICUSPID VALVE	: NORMAL
INTER ATRIAL SEPTUM	: INTACT
INTER VENTRICULAR SEPTUM	: INTACT
PERICARDIUM	: NORMAL
OTHERS	: - NIL

### **IMPRESSION**

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- NORMAL LV SYSTOLIC FUNCTION, LVEF- 58%
- LVH WITH GRADE I LVDD
- MILD MR / MILD TR/ MILD PAH
- NO CLOT / VEGETATION / EFFUSION



**ECHO TECHNICIAN**

*The science of radiology is based upon interpretation of shadows of normal and abnormal tissue. This is neither complete nor accurate; hence, findings should always be interpreted in the light of clinico-pathological correction.*





NAME AND LAB NO	MR RAVI KUMAR A	REG-30016
AGE & SEX	59YRS	MALE
DATE AND AREA OF INTEREST	27.12.2023	ABDOMEN & PELVIS
REF BY	C/ O APOLO CLINIC	

**USG ABDOMEN AND PELVIS**

- LIVER:** Measures 12.3 cm. Normal in size with increased echotexture. No e/o IHBR dilatation. No evidence of SOL. Portal vein appears normal. CBD appears normal. . No e/o calculus / SOL
- GALL BLADDER:** Well distended. Wall appears normal.No e/o calculus/ neoplasm.
- SPLEEN:** Measures 8.5 cm. Normal in size and echotexture. No e/o SOL/ calcification.
- PANCREAS:** Normal in size and echotexture. Pancreatic duct appears normal. No e/o calculus / calcifications.
- RETROPERITONEUM:** Poor window.
- RIGHT KIDNEY:** Right kidney measures 9.5x4.0cm ,is normal in size & echotexture. No evidence of calculus/ hydronephrosis. No solid / cystic lesions.
- LEFT KIDNEY:** Left kidney measures 9.8x4.2 cm ,is normal in size & echotexture. No evidence of calculus/ hydronephrosis. No solid / cystic lesions.
- URETERS:** Bilateral ureters are not dilated.
- URINARY BLADDER:** Well distended. No wall thickening/ calculi. Prevoid 170 cc Post void 107 cc
- PROSTATE:** Enlarged in size volume 30 cc and normal echotexture.
- No evidence of ascites/pleural effusion.


**IMPRESSION:**

- Grade I fatty liver.
- Prostatomegaly with significant post void residue
  - Suggested clinical and PSA correlation

  
**DR PURNIMA PUJAR**  
**MBBS MDRD**

SCAN FOR LOCATION



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male	 2712230016	<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 27-Dec-2023 12:41 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA</b>	6.80	%	Non diabetic adults : <5.7 At risk (Prediabetes) : 5.7 - 6.4 Diagnosing Diabetes : >= 6.5 Diabetes Excellent Control : 6-7 Fair to good Control : 7-8 Unsatisfactory Control : 8-10 Poor Control : >10	HPLC
<b>Estimated Average Glucose(eAG)</b>	148.46	mg/dL		Calculated

**Note:** 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

**Comments:** HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

<b>Vitamin B12-Serum</b>	>2200	pg/mL	211.0-911.0	CLIA
--------------------------	-------	-------	-------------	------

SCAN FOR LOCATION





<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male	 2712230016	<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 27-Dec-2023 12:41 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------


**Comments:** Vitamin B12 performs many important functions in the body, but the most significant function is to act as coenzyme for reducing ribonucleotides to deoxyribonucleotides, a step in the formation of genes. Inadequate dietary intake is not the commonest cause for cobalamine deficiency. The most common cause is malabsorption either due to atrophy of gastric mucosa or diseases of terminal ileum. Cobalamine deficiency leads to Megaloblastic anemia and demyelination of large nerve fibres of spinal cord. Normal body stores are sufficient to last for 3-6 years. Sources of Vitamin B12 are liver, shellfish, fish, meat, eggs, milk, cheese & yogurt.

**Decreased Levels:** Lack of Intrinsic factor: Total or partial gastrectomy, Atrophic gastritis, Intrinsic factor antibodies, Malabsorption: Regional ileitis, resected bowel, Tropical Sprue, Celiac disease, pancreatic insufficiency, bacterial overgrowth & achlorhydria, Loss of ingested vitamin B12: fish tapeworm, Dietary deficiency: Vegetarians, Congenital disorders: Orotic aciduria & transcobalamine deficiency, Increased demand: Pregnancy specially last trimester.

**Increased Levels:** Chronic renal failure, Congestive heart failure, Acute & Chronic Myeloid Leukemia, Polycythemia vera, Carcinomas with liver metastasis, Liver disease, Drug induced cholestasis & Protein malnutrition.




Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	<b>2712230016</b>	<b>Result Date</b> : 27-Dec-2023 01:50 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Complete Haemogram-Whole Blood EDTA</b>				
<b>Haemoglobin (HB)</b>	14.10	g/dL	Male: 14.0-17.0 Female:12.0-15.0 Newborn:16.50 - 19.50	Spectrophotometer
<b>Red Blood Cell (RBC)</b>	4.79	million/cumm	3.50 - 5.50	Volumetric Impedance
<b>Packed Cell Volume (PCV)</b>	41.40	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
<b>Mean corpuscular volume (MCV)</b>	86.30	fL	78.0- 94.0	Calculated
<b>Mean corpuscular hemoglobin (MCH)</b>	29.50	pg	27.50-32.20	Calculated
<b>Mean corpuscular hemoglobin concentration (MCHC)</b>	34.20	%	33.00-35.50	Calculated
<b>Red Blood Cell Distribution Width SD (RDW-SD)</b>	48.70	fL	40.0-55.0	Volumetric Impedance
<b>Red Blood Cell Distribution CV (RDW-CV)</b>	16.90	%	Male: 11.80-14.50 Female:12.20-16.10	Volumetric Impedance
<b>Mean Platelet Volume (MPV)</b>	8.30	fL	8.0-15.0	Volumetric Impedance
<b>Platelet</b>	2.52	lakh/cumm	1.50-4.50	Volumetric Impedance
<b>Platelet Distribution Width (PDW)</b>	9.00	%	8.30 - 56.60	Volumetric Impedance
<b>White Blood cell Count (WBC)</b>	4620.00	cells/cumm	Male: 4000.0-11000.0 Female 4000.0-11000.0 Children: 6000.0-17500.0 Infants : 9000.0-30000.0	Volumetric Impedance





<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	<b>2712230016</b>	<b>Result Date</b> : 27-Dec-2023 01:50 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Neutrophils	57.60	%	40.0-75.0	Light scattering/Manual
Lymphocytes	35.50	%	20.0-40.0	Light scattering/Manual
Eosinophils	2.40	%	0.0-8.0	Light scattering/Manual
Monocytes	4.30	%	0.0-10.0	Light scattering/Manual
Basophils	0.20	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	2.66	10 <sup>3</sup> /uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	1.64	10 <sup>3</sup> /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.20	10 <sup>3</sup> /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	110.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.01	10 <sup>3</sup> /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	38.00	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren


### Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

- RBC'S : Normocytic Normochromic.
- WBC'S : Are normal in total number, morphology and distribution.
- Platelets : Adequate in number and normal in morphology.  
No abnormal cells or hemoparasites are present.
- Impression : Normocytic Normochromic Blood picture.




Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 27-Dec-2023 01:50 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Thyroid function tests (TFT)- Serum</b>				
Tri-Iodo Thyronine (T3)-Serum	1.02	ng/mL	Male: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.30	µg/dL	Male: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	2.02	µIU/mL	Male: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

**Comments:** Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children: 1-3 Days: 1.0-7.40, 1-11 Months: 1.05-2.45, 1-5 Years: 1.05-2.69, 6-10 Years: 0.94-2.41, 11-15 Years: 0.82-2.13, Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester : 0.81-1.90, Second Trimester : 1.0-2.60

**Increased Levels:** Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).

**Decreased Levels:** Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

**Comments:** Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4). It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5, Females : 5.5-11.0, > 60 Years: 5.0-10.70, Cord : 7.40-13.10, Children: 1-3 Days : 11.80-22.60, 1-2 Weeks : 9.90-16.60, 1-4 Months: 7.20-14.40, 1-5 Years : 7.30-15.0, 5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70, Newborn Screen: 1-5 Days: >7.5, 6 Days : >6.5

**Increased Levels:** Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia, Increased transthyretin, estrogen therapy, pregnancy.

**Decreased Levels:** Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

**Comments:** TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester: 0.1-2.5; II -trimester: 0.2-3.0; III- trimester: 0.3-3.0

Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks: 1.7-9.1

**Increased Levels:** Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

**Decreased Levels:** Graves disease, Autonomous thyroid hormone secretion, TSH deficiency.





<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 01:50 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Prostate-Specific Antigen(PSA)-0.36 Serum		ng/mL	0.0-4.0	CLIA

Note: 1. This is a recommended test for detection of prostate cancer along with Digital Rectal Examination (DRE) in males above 50 years of age.  
 2. False negative / positive results are observed in patients receiving mouse monoclonal antibodies for diagnosis or therapy.  
 3. PSA levels may appear consistently elevated / depressed due to the interference by heterophilic antibodies & nonspecific protein binding.  
 4. Immediate PSA testing following digital rectal examination, ejaculation, prostatic massage, indwelling catheterization, ultrasonography and needle biopsy of prostate is not recommended as they falsely elevate levels  
 5. PSA values regardless of levels should not be interpreted as absolute evidence of the presence or absence of disease. All values should be correlated with clinical findings and results of other investigations  
 6. Sites of Non-prostatic PSA production are breast epithelium, salivary glands, periurethral & anal glands, cells of male urethra & breast milk  
 7. Physiological decrease in PSA level by 18% has been observed in hospitalized /sedentary patients either due to supine position or suspended sexual activity.  
 Recommended Testing Intervals: Pre-operatively ( Baseline), 2-4 days post-operatively,Prior to discharge from hospital,Monthly followup if levels are high or show a rising trend.

Clinical Use: -An aid in the early detection of Prostate cancer when used in conjunction with Digital rectal examination in males more than 50 years of age and in those with two or more affected first degree relatives.  
 -Followup and management of Prostate cancer patients  
 -Detect metastatic or persistent disease in patients following surgical or medical treatment of Prostate cancer.  
 Increased Levels : Prostate cancer,Benign Prostatic Hyperplasia,Prostatitis,Genitourinary infections.



Printed By : spectrum  
 Printed On : 27 Dec, 2023 06:13 pm




Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru-560010  
 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	151	mg/dL	70-140	Hexo Kinase

**Comments:** Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.


**Note:** Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

**Comments:** Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

**Probable causes :** Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru-560010  
+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Vitamin D Total (25 Hydroxy Cholecalciferol)</b>	10.6	ng/mL	30.0 -100.0	CLIA

Interpretation: Deficiency :<10, Insufficiency:10-30, Sufficiency:30-100, Toxicity:>100

Note: The assay measures both D2 (Ergocalciferol) and D3 (Cholecalciferol) metabolites of vitamin D.25 (OH)D is influenced by sunlight, latitude, skin pigmentation, sunscreen use and hepatic function.Optimal calcium absorption requires vitamin D 25 (OH) levels exceeding 75 nmol/L.It shows seasonal variation, with values being 40-50% lower in winter than in summer.Levels vary with age and are increased in pregnancy.A new test Vitamin D, Ultrasensitive by LC-MS/MS is also available.

Comments: Vitamin D promotes absorption of calcium and phosphorus and mineralization of bones and teeth. Deficiency in children causes Rickets and in adults leads to Osteomalacia. It can also lead to Hypocalcemia and Tetany. Vitamin D status is best determined by measurement of 25 hydroxy vitamin D, as it is the major circulating form and has longer half life (2-3 weeks) than 1,25 Dihydroxy vitamin D (5-8 hrs).

Decreased Levels:Inadequate exposure to sunlight,Dietary deficiency,Vitamin D malabsorption,Severe Hepatocellular disease,Drugs like Anticonvulsants,Nephrotic syndrome

Increased levels: Vitamin D intoxication.



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	<b>2712230016</b>	<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>LFT-Liver Function Test -Serum</b>				
Bilirubin Total-Serum	0.54	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.11	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.43	mg/dL	0.0-1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	38.00	U/L	15.0-37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	36.00	U/L	Male:16.0-63.0 Female:14.0-59.0 Adult: 45.0-117.0 Children: 48.0-445.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	69.00	U/L	Adult: 45.0-117.0 Children: 48.0-445.0 Infants: 81.90-350.30	PNPP,AMP-Buffer
Protein, Total-Serum	7.39	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	3.97	g/dL	3.40-5.00	Bromocresol Purple
Globulin-Serum	3.42	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.16	Ratio	0.80-1.20	Calculated



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist


SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru-560010  
+91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985




<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>KFT ( Kidney Function Test ) :</b>				
Blood Urea Nitrogen (BUN)-Serum	12.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.74	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	3.28	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	141.5	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.42	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	97.40	mmol/L	94.0-110.0	Ion-Selective Electrodes (ISE)



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Calcium, Total- Serum	9.30	mg/dL	8.50-10.10	Spectrophotometry (O-Cresolphthalein complexone)
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)
Fasting Blood Sugar (FBS)- Plasma	71	mg/dL	60.0-110.0	Hexo Kinase

**Comments:** Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula  $C_6H_{12}O_6$ . It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol, Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.

<b>Gamma-Glutamyl Transferase (GGT)-Serum</b>	16.00	U/L	Male: 15.0-85.0 Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro
---	-------	-----	-------------------------------------	--------------------------------






Name	: MR. RAVIKUMAR A	UHID	: 2712230016	Bill Date	: 27-Dec-2023 08:18 AM
Age / Gender	: 59 years / Male			Sample Col. Date	: 27-Dec-2023 08:18 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 27-Dec-2023 02:57 PM
Reg. No.	: 2712230016			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

**Comments:** Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm




Dr. Nithun Reddy C,MD,Consultant Pathologist

SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985

<b>Name</b> : MR. RAVIKUMAR A	<b>UHIP</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male	 2712230016	<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC		<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		


Test Name	Result	Unit	Reference Value	Method
<b>Blood Group &amp; Rh Typing-Whole Blood EDTA</b>				
Blood Group	AB			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum  
 Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist





<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 02:57 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Lipid Profile-Serum</b>				
Cholesterol Total-Serum	171.00	mg/dL	Male: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	46.00	mg/dL	Male: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	48.00	mg/dL	Male: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	123	mg/dL	Male: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	98.00	mg/dL	Male: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	9	mg/dL	Male: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	3.56	Ratio	Male: 0.0 - 5.0	Calculated


**Interpretation:**

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

**Comments:** As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 04:40 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
<b>Urine Routine Examination-Urine</b>				
<b>Physical Examination</b>				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	7.50		5.0-7.5	Dipstick
Specific Gravity	1.01		1.000-1.030	Dipstick
<b>Biochemical Examination</b>				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
<b>Microscopic Examination</b>				
Pus Cells	2-3	hpf	0.0-5.0	Microscopy
Epithelial Cells	1-2	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

**Comments:** The kidneys help infiltration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist


SCAN FOR LOCATION



Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
+91 77604 97644 | 080 2337 1555 | info@spectrumdiagnostics.org | www.spectrumdiagnostics.org

Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 | +91 6361 253 097 | 080-2991 6944 | 080-49511985



<b>Name</b> : MR. RAVIKUMAR A	<b>UHID</b> : 2712230016	<b>Bill Date</b> : 27-Dec-2023 08:18 AM
<b>Age / Gender</b> : 59 years / Male		<b>Sample Col. Date</b> : 27-Dec-2023 08:18 AM
<b>Ref. By Dr.</b> : Dr. APOLO CLINIC	2712230016	<b>Result Date</b> : 27-Dec-2023 05:29 PM
<b>Reg. No.</b> : 2712230016		<b>Report Status</b> : Final
<b>C/o</b> : Apollo Clinic		

Test Name	Result	Unit	Reference Value	Method
Post Prandial Urine Sugar	Negative		Negative	Dipstick/Benedicts(Man



Printed By : spectrum  
Printed On : 27 Dec, 2023 06:13 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Tejas Arcade, #9/1, 1st Main Road, Dr. Rajkumar Road, Rajajinagar, Opp. St. Theresa Hospital, Bengaluru - 560010  
 +91 77604 97644 | 080 2337 1555 info@spectrumdiagnostics.org www.spectrumdiagnostics.org



Other Branch: #466/A, Ideal Homes Township, 80 Feet Road, Kenchanahalli, Rajarajeshwari Nagar, Bengaluru-560098 +91 6361 253 097 | 080-2991 6944 | 080-49511985