



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor :- BOB
Lab/Hosp :-

Sample Type :- EDTA

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 13:13:49

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

BOB PACKAGE FEMALE BELOW 40

GLYCOSYLATED HEMOGLOBIN (HbA1C)

5.8

%

Method:- HPLC

Non-diabetic: < 5.7
Pre-diabetics: 5.7-6.4
Diabetics: = 6.5 or higher
ADA Target: 7.0
Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycosylated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose over the period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasma glucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHb depends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to the mean of HbA1C. Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1C measurements. The effects vary depending on the specific Hb variant or derivative and the specific HbA1C method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE

120

mg/dL

Method:- Calculated Parameter

Non Diabetic < 100 mg/dL
Prediabetic 100- 125 mg/dL
Diabetic 126 mg/dL or Higher

MUKESH SINGH

Technologist

Page No: 1 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor :- BOB
Lab/Hosp :-

Sample Type :- EDTA

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 13:13:49

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
HAEMOGARAM			
HAEMOGLOBIN (Hb)	11.6	g/dL	12.0 - 15.0
TOTAL LEUCOCYTE COUNT	4.54	/cumm	4.00 - 10.00
DIFFERENTIAL LEUCOCYTE COUNT			
NEUTROPHIL	54.0	%	40.0 - 80.0
LYMPHOCYTE	38.4	%	20.0 - 40.0
EOSINOPHIL	4.3	%	1.0 - 6.0
MONOCYTE	3.0	%	2.0 - 10.0
BASOPHIL	0.3	%	0.0 - 2.0
NEUT#	2.46	10 ³ /uL	1.50 - 7.00
LYMPH#	1.75	10 ³ /uL	1.00 - 3.70
EO#	0.19	10 ³ /uL	0.00 - 0.40
MONO#	0.13	10 ³ /uL	0.00 - 0.70
BASO#	0.01	10 ³ /uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	4.06	x10 ⁶ /uL	3.80 - 4.80
HEMATOCRIT (HCT)	35.00	%	36.00 - 46.00
MEAN CORP VOLUME (MCV)	86.2	fL	83.0 - 101.0
MEAN CORP HB (MCH)	28.7	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	33.3	g/dL	31.5 - 34.5
PLATELET COUNT	213	x10 ³ /uL	150 - 410
RDW-CV	14.0	%	11.6 - 14.0
MENTZER INDEX	21.23		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

MUKESH SINGH

Technologist

Page No: 2 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganeer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor:-BOB
Lab/Hosp :-

Sample Type :- EDTA

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 13:13:49

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

Erythrocyte Sedimentation Rate (ESR)	10	mm/hr.	00 - 20
---	----	--------	---------

(ESR) Methodology : Measurement of ESR by cells aggregation.

Instrument Name : Independent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator of inflammatory disease and abnormal protein states.

The test is used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction)

Levels are higher in pregnancy due to hyperfibrinogenaemia.

The "3-figure ESR " $x > 100$ value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia

(CBC); Methodology: FLC, DLC Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance. and

of connective tissue disease. MCH, MCV, MCHC, MENTZER INDEX are calculated. Instrument Name: Sysmex 6 part fully automatic analyzer XN-L, Japan

MUKESH SINGH

Technologist

Page No: 3 of 12

Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
 Sex / Age :- Female 37 Yrs 5 Mon 23 Days
 Company :- MediWheel

Patient ID :- 1223342
 Ref. By Doctor:-BOB
 Lab/Hosp :-

Sample Type :- PLAIN/SERUM

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 11:08:54

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

LIPID PROFILE

TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	118.82	mg/dl	Desirable <200 Borderline 200-239 High > 240
TRIGLYCERIDES Method:- GPO-PAP	89.38	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	34.77	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	69.15	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	17.88	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	3.42		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	1.99		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED	376.55	mg/dl	400.00 - 1000.00

TOTAL CHOLESTEROL InstrumentName:Radox Rx Imola **Interpretation:** Cholesterol measurements are used in the diagnosis and treatments of lipid lipoprotein metabolism disorders.

TRIGLYCERIDES InstrumentName:Radox Rx Imola **Interpretation :** Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.

DIRECT HDLCHOLESTERO InstrumentName:Radox Rx Imola **Interpretation:** An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.

DIRECT LDL-CHOLESTEROL InstrumentName:Radox Rx Imola **Interpretation:** Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.

TOTAL LIPID AND VLDL ARE CALCULATED

SURENDRAXHANGA

Page No: 4 of 12

Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
 Sex / Age :- Female 37 Yrs 5 Mon 23 Days
 Company :- MediWHEEL

Patient ID :- 1223342
 Ref. By Doctor:- BOB
 Lab/Hosp :-

Sample Type :- PLAIN/SERUM

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 11:08:54

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
LIVER PROFILE WITH GGT			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.26	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1 month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.12	mg/dL	Adult - Up to 0.25 Newborn - <0.6 mg/dL >- 1 month - <0.2 mg/dL
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.14	mg/dl	0.30-0.70
SGOT Method:- IFCC	15.1	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	20.4	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	84.00	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	25.50	U/L	7.00 - 32.00
SERUM TOTAL PROTEIN Method:- Biuret Reagent	6.63	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.16	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	2.47	gm/dl	2.20 - 3.50
A/G RATIO	1.68		1.30 - 2.50

Total Bilirubin Methodology: Colorimetric method Instrument Name: Randox Rx Imola Interpretation: An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in rhesus incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

AST Aspartate Aminotransferase Methodology: IFCC Instrument Name: Randox Rx Imola Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

ALT Alanine Aminotransferase Methodology: IFCC Instrument Name: Randox Rx Imola Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminases can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

Alkaline Phosphatase Methodology: AMP Buffer Instrument Name: Randox Rx Imola Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and intestinal disease.

TOTAL PROTEIN Methodology: Biuret Reagent Instrument Name: Randox Rx Imola Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

ALBUMIN (ALB) Methodology: Bromocresol Green Instrument Name: Randox Rx Imola Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

Instrument Name Randox Rx Imola **Interpretation:** Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra- or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal) are observed with infectious hepatitis.

SURENDRAKHANGA

Dr. Chandrika Gupta
 MBBS.MD (Path)
 RMC NO. 21021/008037





Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor:-BOB
Lab/Hosp :-

Sample Type :- PLAIN/SERUM

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 12:12:39

IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

TOTAL THYROID PROFILE

SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.210	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	8.510	ug/dl	5.500 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	3.970	μIU/mL	0.500 - 6.880

Interpretation: Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

Interpretation : The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

Interpretation : TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

AJAYKUMAR

Technologist

Page No: 6 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
 Sex / Age :- Female 37 Yrs 5 Mon 23 Days
 Company :- MediWheel

Patient ID :- 1223342
 Ref. By Doctor:-BOB
 Lab/Hosp :-

Sample Type :- URINE

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 15:33:54

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
Urine Routine			
<u>PHYSICAL EXAMINATION</u>			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
<u>CHEMICAL EXAMINATION</u>			
REACTION(PH) Method:- Reagent Strip(Double indicator blue reaction)	5.5		5.0 - 7.5
SPECIFIC GRAVITY Method:- Reagent Strip(bromthymol blue)	1.020		1.010 - 1.030
PROTEIN Method:- Reagent Strip (Sulphosalicylic acid test)	NIL		NIL
GLUCOSE Method:- Reagent Strip (Glu.Oxidase Peroxidase Benedict)	NIL		NIL
BILIRUBIN Method:- Reagent Strip (Azo-coupling reaction)	NEGATIVE		NEGATIVE
UROBILINOGEN Method:- Reagent Strip (Modified ehrlich reaction)	NORMAL		NORMAL
KETONES Method:- Reagent Strip (Sodium Nitropruside) Rothera's	NEGATIVE		NEGATIVE
NITRITE Method:- Reagent Strip (Diazotization reaction)	NEGATIVE		NEGATIVE
<u>MICROSCOPY EXAMINATION</u>			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	1-2	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT

VIJENDRAMEENA

Technologist

Page No: 7 of 12

Dr. Rashmi Bakshi
 MBBS. MD (Path)
 RMC No. 17975/008828



CONDITIONS OF REPORTING SEE OVER LEAF"

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganeer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor :- BOB
Lab/Hosp :-

Sample Type :- STOOL

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 15:33:54

CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
STOOL ANALYSIS			
PHYSICAL EXAMINATION			
MUCUS			
BLOOD			
MICROSCOPIC EXAMINATION			
RBC's		/HPF	
WBC/HPF		/HPF	
OVA			
CYSTS			
OTHERS			
Collected	Sample	Received	

VIJENDRAMEENA

Technologist

Page No: 8 of 12

Dr. Rashmi Bakshi
MBBS. MD (Path)
RMC No. 17975/008828



CONDITIONS OF REPORTING SEE OVER LEAF"



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor:-BOB
Lab/Hosp :-

Sample Type :- KOx/Na FLUORIDE-F, KOx/Na Substrate, Reagent, INHIBITORS, SALT, BUFFER
Sample ID :- 22042023080902

Final Authentication : 22/04/2023 13:08:03

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP	103.7	mg/dl	75.0 - 115.0
Impaired glucose tolerance (IGT)	111 - 125 mg/dL		
Diabetes Mellitus (DM)	> 126 mg/dL		

Instrument Name: Randox Rx Imola **Interpretation:** Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .

BLOOD SUGAR PP (Plasma) 105.1 mg/dl 70.0 - 140.0
Method:- GOD PAP

Instrument Name: Randox Rx Imola **Interpretation:** Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases .

SERUM CREATININE 0.91 mg/dl Men - 0.6-1.30
Method:- Colorimetric Method Women - 0.5-1.20
SERUM URIC ACID 5.94 mg/dl Men - 3.4-7.0
Method:- Enzymatic colorimetric Women - 2.4-5.7

SURENDRAXHANGA

Page No: 9 of 12



Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor :- BOB
Lab/Hosp :-

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

AJAYKUMAR, ANITASHARMA, MUKESHSINGH, SURENDRAKHANGA, VIJENDRAMEENA



Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor:-BOB
Lab/Hosp :-

Sample Type :- EDTA, URINE

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 15:33:54

HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BLOOD GROUP ABO	"O"POSITIVE		
BLOOD GROUP ABO Methodology : Haemagglutination reaction Kit Name : Monoclonal agglutinating antibodies (Span clone).			
URINE SUGAR (FASTING) Collected Sample Received	Nil		Nil

MUKESH SINGH, VIJENDRAMEENA

Technologist

Page No: 11 of 12

Dr. Rashmi Bakshi
MBBS. MD (Path)
RMC No. 17975/008828
Dr. Chandrika Gupta



CONDITIONS OF REPORTING SEE OVER LEAF"

Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Opp. Janpath Corner, New Sanganer Road, Jaipur
Tele : 0141-2293346, 4049787, 9887049787
Website : www.drgoyalspathlab.com | E-mail : drgoyalpiyush@gmail.com



Date :- 22/04/2023 08:09:02
NAME :- Mrs. URVASHI SOGANI
Sex / Age :- Female 37 Yrs 5 Mon 23 Days
Company :- MediWheel

Patient ID :- 1223342
Ref. By Doctor :- BOB
Lab/Hosp :-

Sample Type :- PLAIN/SERUM

Sample Collected Time 22/04/2023 08:32:08

Final Authentication : 22/04/2023 11:08:54

BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
BLOOD UREA NITROGEN (BUN)	9.2	mg/dl	0.0 - 23.0

*** End of Report ***

SURENDRAKHANGA

Page No: 12 of 12

Dr. Chandrika Gupta
MBBS.MD (Path)
RMC NO. 21021/008037



CONDITIONS OF REPORTING SEE OVER LEAF"