

CODE/NAME & ADDRESS : C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000624 PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

AGE/SEX :33 Years Female :09/02/2024 11:42:00 RECEIVED: 09/02/2024 11:43:09 REPORTED :15/02/2024 14:16:24

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

<u></u>			
ŀ	IAEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECKUP B	ELOW 40FEMALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB)	11.1 Low	12.0 - 15.0	g/dL
RED BLOOD CELL (RBC) COUNT	4.12	3.8 - 4.8	mil/μL
WHITE BLOOD CELL (WBC) COUNT	7.89	4.0 - 10.0	thou/µL
PLATELET COUNT	196	150 - 410	thou/µL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV)	34.8 Low	36 - 46	%
MEAN CORPUSCULAR VOLUME (MCV)	84.5	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	26.9 Low	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	31.9	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW)	14.4 High	11.6 - 14.0	%
MENTZER INDEX	20.5		
MEAN PLATELET VOLUME (MPV)	12.6 High	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS	63	40 - 80	%
LYMPHOCYTES	21	20 - 40	%
MONOCYTES	06	2 - 10	%
EOSINOPHILS	10 High	1 - 6	%
DA CODUTE C	0.0		0/

BASOPHILS

Sh>Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

< 1 - 2

00

Dr. Itisha Dhiman **Pathologist**





Page 1 Of 18

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India

Tel: 0291-2646000, 2644000, Fax: CIN - U74899PB1995PLC045956 Email: srl.jodhpur@gmail.com



%



PATIENT NAME: SHIVANI DUBE 43384 REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030

8800465156

ACCESSION NO: **0061XB000624**PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO : DRAWN :09/02/2024 11:42:00 RECEIVED :09/02/2024 11:43:09

:33 Years

AGE/SEX

REPORTED :15/02/2024 14:16:24

Test Report Status <u>Final</u> Results Biological Reference Interval Units

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020)

This ratio element is a calculated parameter and out of NABL scope.

grisha.

Dr. Itisha Dhiman Pathologist





Page 2 Of 18

View Details

View Report



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: **0061XB000624**PATIENT ID : SHIVF09029161

CLIENT PATIENT ID:

AGE/SEX :33 Years Female
DRAWN :09/02/2024 11:42:00
RECEIVED :09/02/2024 11:43:09
REPORTED :15/02/2024 14:16:24

Test Report Status <u>Final</u> Results Biological Reference Interval Units

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD

E.S.R 15 0 - 20

METHOD : WESTERGREN METHOD

mm at 1 hr

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD

HBA1C 5.6 Non-diabetic: < 5.7 %

Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0

Action suggested: > 8.0

ESTIMATED AVERAGE GLUCOSE(EAG) 114.0 < 116.0 mg/dL

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-TEST DESCRIPTION:

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

<br

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease

(Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

salicylates)

REFERENCE :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition.

Dr. Itisha Dhiman Pathologist





Page 3 Of 18

View Details

View Report

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School

Jodhpur, 342001 Rajasthan, India





REF. DOCTOR: SELF PATIENT NAME: SHIVANI DUBE 43384

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000624

PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

:09/02/2024 11:42:00 DRAWN

:33 Years

AGE/SEX

RECEIVED: 09/02/2024 11:43:09 REPORTED :15/02/2024 14:16:24

Test Report Status Results **Biological Reference Interval** <u>Final</u> Units

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2. Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).
The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
2. eAG gives an evaluation of blood glucose levels for the last couple of months.
3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c - 46.7

-

 anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.
 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates &
- opiates addiction are reported to interfere with some assay methods, falsely increasing results.

 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
- c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

Dr. Itisha Dhiman **Pathologist**





Page 4 Of 18

View Report

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001 Rajasthan, India





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

Test Report Status

8800465156

ACCESSION NO: 0061XB000624

PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

AGE/SEX :33 Years Female :09/02/2024 11:42:00 RECEIVED: 09/02/2024 11:43:09

REPORTED :15/02/2024 14:16:24

Biological Reference Interval Units

IMMUNOHAEMATOLOGY

Results

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

<u>Final</u>

TYPE A **ABO GROUP**

METHOD: FORWARD/REVERSE

POSITIVE RH TYPE

METHOD: FORWARD/REVERSE

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.'

The test is performed by both forward as well as reverse grouping methods.

Dr. Itisha Dhiman **Pathologist**





Page 5 Of 18

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India Tel: 0291-2646000, 2644000, Fax:

CIN - U74899PB1995PLC045956 Email: srl.jodhpur@gmail.com





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000624

PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

AGE/SEX :33 Years Female :09/02/2024 11:42:00 RECEIVED: 09/02/2024 11:43:09 REPORTED :15/02/2024 14:16:24

Test Report Status Results Biological Reference Interval Units <u>Final</u>

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

104 High Normal : < 100 mg/dL FBS (FASTING BLOOD SUGAR)

Pre-diabetes: 100-125 Diabetes: >/=126

METHOD: SPECTROPHOTOMETRY

LIPID PROFILE WITH CALCULATED LDL

CHOLESTEROL, TOTAL 179 < 200 Desirable mg/dL

200 - 239 Borderline High

>/= 240 High

148 < 150 Normal mg/dL TRIGLYCERIDES

150 - 199 Borderline High

200 - 499 High

>/=500 Very High

METHOD: SPECTROPHOTOMETRY

METHOD: SPECTROPHOTOMETRY

HDL CHOLESTEROL 34 Low < 40 Low mg/dL

>/=60 High

METHOD: SPECTROPHOTOMETRY

115 High mg/dL CHOLESTEROL LDL < 100 Optimal

100 - 129

Near optimal/ above optimal

130 - 159 Borderline High 160 - 189 High >/= 190 Very High

145 High NON HDL CHOLESTEROL mg/dL Desirable: Less than 130

Above Desirable: 130 - 159

Borderline High: 160 - 189

High: 190 - 219

Very high: > or = 220

VERY LOW DENSITY LIPOPROTEIN 29.6 </= 30.0 mg/dL

CHOL/HDL RATIO 5.3 High 3.3 - 4.4Low Risk

4.5 - 7.0 Average Risk

Dr. Itisha Dhiman **Pathologist**



Page 6 Of 18



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





PATIENT NAME: SHIVANI DUBE 43384 REF. DOCTOR: SELF CODE/NAME & ADDRESS: C000138375 ACCESSION NO: 0061XB000624 AGE/SEX :33 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID :09/02/2024 11:42:00 : SHIVF09029161 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 09/02/2024 11:43:09 DELHI ABHA NO REPORTED :15/02/2024 14:16:24 **NEW DELHI 110030** 8800465156

Test Report Status <u>Final</u> Results Biological Reference Interval Units

7.1 - 11.0 Moderate Risk > 11.0 High Risk

LDL/HDL RATIO **34.0 High** 0.5 - 3.0 Desirable/Low Risk

3.1 - 6.0 Borderline/Moderate

Risk

>6.0 High Risk

Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

	The Te (Title obere one curulo abeliar un	/ 1		
Risk Category				
Extreme risk group	A.CAD with > 1 feature of high risk group			
	B. CAD with > 1 feature of Very high risk g	group or recurrent ACS (within 1 year) despite LDL-C < or =		
	50 mg/dl or polyvascular disease			
Very High Risk	1. Established ASCVD 2. Diabetes with 2 r	najor risk factors or evidence of end organ damage 3.		
	Familial Homozygous Hypercholesterolemia	a		
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ			
	damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary			
	Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid plaque			
Moderate Risk	2 major ASCVD risk factors			
Low Risk	0-1 major ASCVD risk factors			
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors				
1. Age > or = 45 years in males and > or = 55 years in females 3. Current Cigarette smoking or tobacco use				
2. Family history of premature ASCVD 4. High blood pressure				
5. Low HDL				

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
	$\langle OR = 30 \rangle$	< OR = 60)		
Extreme Risk Group Category B	<or 30<="" =="" td=""><td><or 60<="" =="" td=""><td>> 30</td><td>>60</td></or></td></or>	<or 60<="" =="" td=""><td>> 30</td><td>>60</td></or>	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

^{*}After an adequate non-pharmacological intervention for at least 3 months.

Dr. Itisha Dhiman Pathologist





Page 7 Of 18

View Report



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School

Jodhpur, 342001 Rajasthan, India







PATIENT NAME: SHIVANI DUBE 43384 REF. DOCTOR: SELF CODE/NAME & ADDRESS: C000138375 ACCESSION NO: 0061XB000624 AGE/SEX :33 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : SHIVF09029161 :09/02/2024 11:42:00 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 09/02/2024 11:43:09 DELHI ABHA NO REPORTED :15/02/2024 14:16:24 **NEW DELHI 110030** 8800465156

Test Report Status <u>Final</u> Results Biological Reference Interval Units

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER	FUNCTION	PROFILE.	SERUM

BILIRUBIN, TOTAL METHOD: SPECTROPHOTOMETRY	0.80	0.2 - 1.0	mg/dL
BILIRUBIN, DIRECT	0.10	0.0 - 0.2	mg/dL
METHOD: SPECTROPHOTOMETRY BILIRUBIN, INDIRECT	0.70	0.1 - 1.0	mg/dL
METHOD: SPECTROPHOTOMETRY TOTAL PROTEIN	7.9	6.4 - 8.2	g/dL
METHOD: SPECTROPHOTOMETRY ALBUMIN	4.1	3.4 - 5.0	g/dL
METHOD: SPECTROPHOTOMETRY	4.1		.
GLOBULIN METHOD: CALCULATED PARAMETER	3.8	2.0 - 4.1	g/dL
ALBUMIN/GLOBULIN RATIO METHOD: CALCULATED PARAMETER	1.1	1.0 - 2.1	RATIO
ASPARTATE AMINOTRANSFERASE	26	15 - 37	U/L
(AST/SGOT) METHOD: SPECTROPHOTOMETRY			
ALANINE AMINOTRANSFERASE (ALT/SGPT) METHOD: SPECTROPHOTOMETRY	43 High	< 34.0	U/L
ALKALINE PHOSPHATASE METHOD: SPECTROPHOTOMETRY	148 High	30 - 120	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT)	102 High	5 - 55	U/L
METHOD: SPECTROPHOTOMETRY LACTATE DEHYDROGENASE	156	81 - 234	U/L
METHOD: SPECTROPHOTOMETRY			

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN	12	6 - 20	mg/dL
METHOD: SPECTROPHOTOMETRY			

CREATININE, SERUM

CREATININE **0.59 Low** 0.60 - 1.10 mg/dL

METHOD: SPECTROPHOTOMETRY

Dr. Itisha Dhiman Pathologist



Page 8 Of 18

View Details

View Report

PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030 8800465156 ACCESSION NO: **0061XB000624**PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO : AGE/SEX :33 Years Female
DRAWN :09/02/2024 11:42:00
RECEIVED :09/02/2024 11:43:09
REPORTED :15/02/2024 14:16:24

Test Report Status <u>Final</u> Results Biological Reference Interval Units

BUN/CREAT RATIO

BUN/CREAT RATIO **20.34 High** 5.00 - 15.00

METHOD: SPECTROPHOTOMETRY

URIC ACID, SERUM

URIC ACID 3.9 2.6 - 6.0 mg/dL

TOTAL PROTEIN, SERUM

METHOD: SPECTROPHOTOMETRY

TOTAL PROTEIN 7.9 6.4 - 8.2 g/dL

METHOD: SPECTROPHOTOMETRY

ALBUMIN, SERUM

ALBUMIN 4.1 3.4 - 5.0 g/dL

METHOD: SPECTROPHOTOMETRY

GLOBULIN

GLOBULIN 3.8 2.0 - 4.1 g/dL

 ${\tt METHOD}: {\tt CALCULATED} \ {\tt PARAMETER}$

ELECTROLYTES (NA/K/CL), SERUM

SODIUM, SERUM 137 136 - 145 mmol/L

METHOD: ION SELECTIVE ELECTRODE TECHNOLOGY

POTASSIUM, SERUM 4.4 3.50 - 5.10

METHOD: ION SELECTIVE ELECTRODE TECHNOLOGY

CHLORIDE, SERUM

110 High

98 - 107 mmol/L

Dr. Itisha Dhiman

Pathologist



Page 9 Of 18

View Details

View Repor

PERFORMED AT :

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001

Rajasthan, India

Tel: 0291-2646000, 2644000, Fax: CIN - U74899PB1995PLC045956 Email: srl.jodhpur@gmail.com



mmol/L



REF. DOCTOR: SELF PATIENT NAME: SHIVANI DUBE 43384

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0061XB000624 PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

:09/02/2024 11:42:00 DRAWN

:33 Years

AGE/SEX

RECEIVED: 09/02/2024 11:43:09 REPORTED :15/02/2024 14:16:24

Test Report Status Results **Biological Reference Interval** <u>Final</u> Units

METHOD: ION SELECTIVE ELECTRODE TECHNOLOGY

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake, prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome, osmotic diuresis (e.g., hyperglycemia), alkalosis, familial periodic paralysis, trauma (transient). Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics.
Increased in: Dehydration (excessivesweating, severe vomiting or diarrhea),diabetes mellitus, diabetesinsipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice,oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration,renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium- sparing diuretics,NSAIDs, beta-blockers, ACE inhibitors, highdose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in

 db-Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.
 db-Decreased in :Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease, malignancy (adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. LIVER FUNCTION PROFILE, SERUM-

 bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts.

Dr. Itisha Dhiman **Pathologist**



Page 10 Of 18





PERFORMED AT:

Agilus Diagnostics Ltd.

M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001





REF. DOCTOR: SELF PATIENT NAME: SHIVANI DUBE 43384

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000624

PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

AGE/SEX :33 Years Female :09/02/2024 11:42:00 DRAWN

RECEIVED: 09/02/2024 11:43:09 REPORTED :15/02/2024 14:16:24

Test Report Status Results **Biological Reference Interval Final** Units

Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

 measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver,liver cancer,kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

 obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

 has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that

increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection,including HIV and hepatitis B or C,Multiple myeloma,Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc
BLOOD UREA NITROGEN (BUN), SERUM-
b>Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

b>Causes of decreased level include Liver disease, SIADH. CREATININE, SERUM-
b>Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

<br

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin.

<
b>Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage),Burns,Glomerulonephritis, Liver disease, Malabsorption, Malnutrition,

Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Itisha Dhiman **Pathologist**





Page 11 Of 18

View Report

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

CIN - U74899PB1995PLC045956 Email: srl.jodhpur@gmail.com





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHÍ

NEW DELHI 110030

8800465156

ACCESSION NO: **0061XB000624**PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO : AGE/SEX :33 Years Female
DRAWN :09/02/2024 11:42:00
RECEIVED :09/02/2024 11:43:09
REPORTED :15/02/2024 14:16:24

Test Report Status <u>Final</u> Results Biological Reference Interval Units

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW
APPEARANCE SLIGHTLY HAZY

CHEMICAL EXAMINATION, URINE

PH	5.0	4.7 - 7.5
SPECIFIC GRAVITY	1.010	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	2-3	0-5	/HPF
EPITHELIAL CELLS	8-10	0-5	/HPF

CASTS NOT DETECTED
CRYSTALS NOT DETECTED

BACTERIA DETECTED NOT DETECTED

(OCCASIONAL)

 ${\tt METHOD}: {\tt MICROSCOPIC} \ {\tt EXAMINATION}$

YEAST NOT DETECTED NOT DETECTED

Dr. Itisha Dhiman Pathologist



Page 12 Of 18

View Details



PERFORMED AT :

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre,Ground Floor,C-22,Shastri Nagar,Near Central Academy School Jodhpur, 342001

Rajasthan, India





PATIENT NAME: SHIVANI DUBE 43384 REF. DOCTOR: SELF CODE/NAME & ADDRESS : C000138375 ACCESSION NO: 0061XB000624 AGE/SEX :33 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : SHIVF09029161 :09/02/2024 11:42:00 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 09/02/2024 11:43:09 DELHI REPORTED :15/02/2024 14:16:24 ABHA NO **NEW DELHI 110030** 8800465156

Test Report Status Final Results **Biological Reference Interval** Units

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions		
Proteins	Inflammation or immune illnesses		
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind		
·	of kidney impairment		
Glucose	Diabetes or kidney disease		
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst		
Urobilinogen	Liver disease such as hepatitis or cirrhosis		
Blood	Renal or genital disorders/trauma		
Bilirubin	Liver disease		
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases		
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions		
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time		
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein		
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases		
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice		
Uric acid	arthritis		
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.		
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis		

Dr. Itisha Dhiman **Pathologist**






Page 13 Of 18



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001





CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0061XB000624 PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

AGE/SEX :33 Years Female :09/02/2024 11:42:00 DRAWN RECEIVED: 09/02/2024 11:43:09 REPORTED :15/02/2024 14:16:24

Test Report Status Results Biological Reference Interval Units <u>Final</u>

CYTOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PAPANICOLAOU SMEAR

TEST METHOD CONVENTIONAL GYNEC CYTOLOGY

TWO UNSTAINED CERVICAL SMEARS RECEIVED SPECIMEN TYPE

2014 BETHESDA SYSTEM FOR REPORTING CERVICAL CYTOLOGY REPORTING SYSTEM

SATISFACTORY SPECIMEN ADEQUACY

SMEARS SHOW SUPERFICIAL AND INTERMEDIATE SQUAMOUS **MICROSCOPY**

CELLS SHOWING MILD REACTIVE CHANGES OF INFLAMMATION. BACKGROUND SHOW LACTOBACILLI AND NEUTROPHILS.

ENDOCERVICAL COMPONENT ABSENT.

METHOD: MANUAL

INTERPRETATION / RESULT NEGATIVE FOR INTRAEPITHELIAL LESION OR MALIGNANCY

Dr. Itisha Dhiman **Pathologist**





Page 14 Of 18



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





CODE/NAME & ADDRESS: C000138375

ARCOFEMI HEALTHCARE LTD (MEDIWHEEL
F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156 ACCESSION NO: **0061XB000624**PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO : AGE/SEX :33 Years Female
DRAWN :09/02/2024 11:42:00
RECEIVED :09/02/2024 11:43:09
REPORTED :15/02/2024 14:16:24

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE PHYSICAL EXAMINATION, STOOL

COLOUR SAMPLE NOT RECEIVED

grisha.

Dr. Itisha Dhiman Pathologist





Page 15 Of 18

View Details

View Report



Agilus Diagnostics Ltd.
M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001





PATIENT NAME: SHIVANI DUBE 43384 REF. DOCTOR: SELF

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030

8800465156

ACCESSION NO: 0061XB000624 PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

DRAWN :09/02/2024 11:42:00 RECEIVED: 09/02/2024 11:43:09

:33 Years

AGE/SEX

2nd Trimester 0.200 - 3.000 3rd Trimester 0.300 - 3.000

REPORTED :15/02/2024 14:16:24

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

THYROID PANEL, SERUM

98.74 ng/dL T3 Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester: 105.0 - 230.0 2nd Trimester: 129.0 - 262.0 3rd Trimester: 135.0 - 262.0 T4 7.41 Non-Pregnant Women µg/dL 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70 17.430 High TSH (ULTRASENSITIVE) Non Pregnant Women µIU/mL 0.27 - 4.20Pregnant Women (As per American Thyroid Association) 1st Trimester 0.100 - 2.500

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyporthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Total T4 FT4 Total T3 **Possible Conditions**

Dr. Itisha Dhiman **Pathologist**



Page 16 Of 18

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001

Rajasthan, India





PATIENT NAME: SHIVANI DUBE 43384 REF. DOCTOR: SELF CODE/NAME & ADDRESS: C000138375 ACCESSION NO: 0061XB000624 AGE/SEX :33 Years Female ARCOFEMI HEALTHCARE LTD (MEDIWHEEL PATIENT ID : SHIVF09029161 :09/02/2024 11:42:00 F-703, LADO SARAI, MEHRAULISOUTH WEST CLIENT PATIENT ID: RECEIVED: 09/02/2024 11:43:09 DELHI ABHA NO REPORTED :15/02/2024 14:16:24 **NEW DELHI 110030** 8800465156

Test Report Status Final Results Biological Reference Interval Units

1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

Dr. Itisha Dhiman


Pathologist



Page 17 Of 18

PERFORMED AT:

Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School Jodhpur, 342001







REF. DOCTOR: SELF PATIENT NAME: SHIVANI DUBE 43384

CODE/NAME & ADDRESS: C000138375 ARCOFEMI HEALTHCARE LTD (MEDIWHEEL F-703, LADO SARAI, MEHRAULISOUTH WEST

DELHI

NEW DELHI 110030 8800465156

ACCESSION NO: 0061XB000624 PATIENT ID : SHIVF09029161

CLIENT PATIENT ID: ABHA NO

AGE/SEX :33 Years Female :09/02/2024 11:42:00 DRAWN RECEIVED: 09/02/2024 11:43:09

REPORTED :15/02/2024 14:16:24

Test Report Status Results Biological Reference Interval Units **Final**

> **End Of Report** Please visit www.agilusdiagnostics.com for related Test Information for this accession

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the AGILUS Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- AGILUS Diagnostics confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- Test results cannot be used for Medico legal purposes.
- In case of queries please call customer care (91115 91115) within 48 hours of the report.

Agilus Diagnostics Ltd

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062

Dr. Itisha Dhiman **Pathologist**





Page 18 Of 18



Agilus Diagnostics Ltd. M/S S.S. Wellness Centre, Ground Floor, C-22, Shastri Nagar, Near Central Academy School

Jodhpur, 342001 Rajasthan, India

