

CERTIFICATE OF MEDICAL FITNESS

NAME: Padmaja

AGE/ GENDER: 24y / Female

HEIGHT: 153cm

WEIGHT: 63kg

IDENTIFICATION MARK: -

BLOOD PRESSURE: 110/70 mmHg

PULSE: 102 bpm

CVS: Normal

RS:P

ANY OTHER DISEASE DIAGNOSED IN THE PAST: Nil

ALLERGIES, IF ANY: Nil

LIST OF PRESCRIBED MEDICINES: Nil

ANY OTHER REMARKS: No

I Certify that I have carefully examined Mr/Mrs. padmaja. son/daughter of Mr Havinath who has signed in my presence. He/ she has no physical disease and is fit for employment.

P. Padmaja

Signature of candidate

Dr. BINDU P. J. R

MBBS MD

Signature of Medical Officer

Reg. No. 62200

Place: Spectrum Diagnostics & Health Care

Date: 24/2/24

Disclaimer: The patient has not been checked for COVID. This certificate does not relate to the covid status of the patient examined





SPECTRUM DIAGNOSTICS

Bangalore

Patient ID : 0167

Name : MRS PADMAJA



CR Number : 20240224123151

Registration Date : 24-Feb-2024

Age : 24

Gender : Female

Operator : spectrum diagnostics



125 Hz 250 Hz 500 Hz 750 Hz 1000 H 1500 H 2000 H 3000 H 4000 H 6000 H 8000 H

X - Air Left	30	45	50	40	25	30	25	20	35	15	20
O - Air Right	45	45	45	40	35	35	30	25	35	25	15
> - Bone Left											
< - Bone Right											

	Average	High	Mid	Low
AIR Left	30.45 dB	22.50 dB	26.67 dB	41.25 dB
AIR Right	34.00 dB	25.00 dB	33.33 dB	43.75 dB

Clinical Notes :

Not Found




HR	: 114	bpm
P	: 93	ms
PR	: 124	ms
QRS	: 73	ms
QT/QTc	: 300/414	ms
PQRST	: 55/54/36	ms
RV5/SV1	: 1.528/0.718	mV

Diagnosis Information:

Sinus Tachycardia
Low T Wave(V3,V4,V5,V6)

Report Confirmed by:




NAME : MRS.PADMAJA	DATE : 24/02/2024
AGE/SEX : 24YEARS/FEMALE	REG NO: 2402240031
REF BY : APOLLO CLINIC	

CHEST PA VIEW

- Visualised lungs are clear .
- Bilateral hila appears normal .
- Cardia is normal in size
- No pleural effusion

IMPRESSION: No Significant Abnormality Detected



**DR PRAVEEN B,DMRD ,DNB
Consultant Radiologist**

SCAN FOR LOCATION



NAME AND LAB NO	MRS PADMAJA	REG -40031
AGE & SEX	24 YRS	FEMALE
DATE AND AREA OF INTEREST	24.02.2024	ABDOMEN & PELVIS
REF BY	C/O APOLO CLINIC	

USG ABDOMEN AND PELVIS

- LIVER:** Normal in size and echotexture.
No e/o IHBR dilatation. No evidence of focal lesion
Portal vein appears normal.
CBD appears normal.
- GALL BLADDER:** Well distended. Wall appears normal.No e/o calculus.
- SPLEEN:** Normal in size and echotexture. No focal lesion
- PANCREAS:** Head and body appears normal . Tail obscured by bowel gas shadows
- RETROPERITONEUM:** Appears normal to the extent visualized . No significant lymphadenopathy .
- RIGHT KIDNEY:** Measures 9.4 X1.3 cm Right kidney is normal in size & echotexture
No evidence of calculus/ hydronephrosis.
- LEFT KIDNEY:** Measures 8.9 X1.3 cm .Left kidney is normal in size & echotexture
No evidence of calculus/ hydronephrosis.
- URINARY BLADDER:** Well distended. No wall thickening/ calculi.
- UTERUS:** Anteverted, Normal in size and echotexture
Endometrium is normal.ET -9.4 mm.
- OVARIES:** B/L ovaries normal in size and echotexture.
No obvious adnexal mass lesions .

- Minimal free fluid in POD – likely reactive

IMPRESSION:

- No significant sonological abnormality detected



**DR PRAVEEN B , DMRD , DNB
CONSULTANT RADIOLOGIST**



NAME	AGE	GENDER
Ms. Pooja	24 Yrs	Female

DENTAL EXAMINATION REPORT:

8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8
8	7	6	5	4	3	2	1	1	2	3	4	5	6	7	8

C: CAVITY → None

M: MISSING → None

O: OTHERS → Gen. Plaque

ADVISED:

CLEANING / SCALING / ROOTS PLANNING / FLOSSING & POLISHING / OTHERS

REMARKS:

[Signature]
24/02/24

SIGNATURE OF THE DENTAL SURGEON

SEAL

DATE

Dr. SACHDEV NAGARKAR
B.D.S., F.A.G.E., F.P.F.A. (USA)
Reg. No : 2247/A

SCAN FOR LOCATION



Dr. Ashok S
Bsc., MBBS., D.O.M.S
Consultant Ophthalmologist
KMC No: 31827

DATE: 24-02-24.

EYE EXAMINATION

NAME: *Ms. padmaja.* AGE: *24y* GENDER: F / M

	RIGHT EYE	LEFT EYE
Vision	<i>6/6:06</i>	<i>6/6:06</i>
Vision With glass		
Color Vision	Normal	Normal
Anterior segment examination	Normal	Normal
Fundus Examination	Normal	Normal
Any other abnormality	Nil	Nil
Diagnosis/ Impression	Normal	Normal

DR. ASHOK SARODHE
B.Sc., M.B.B.S., D.O.M.S.
Consultant & Surgeon
KMC 31827

SCAN FOR LOCATION



PATIENT NAME	MRS PADMAJA	ID NO	2402240031
AGE	24YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	24.02.2024

2D ECHO CARDIOGRAHIC STUDY

M-MODE

AORTA	22mm
LEFT ATRIUM	31mm
RIGHT VENTRICLE	20mm
LEFT VENTRICLE (DIASTOLE)	40mm
LEFT VENTRICLE(SYSTOLE)	29mm
VENTRICULAR SEPTUM (DIASTOLE)	06mm
VENTRICULAR SEPTUM (SYSTOLE)	08mm
POSTERIOR WALL (DIASTOLE)	06mm
POSTERIOR WALL (SYSTOLE)	09mm
FRACTIONAL SHORTENING	30%
EJECTION FRACTION	60%

DOPPLER /COLOUR FLOW

Mitral Valve Velocity : MVE- 0.89m/s MVA – 0.40m/s E/A-2.23

Tissue Doppler : e' (Septal) -9cm/s E/e'(Septal) -9

Velocity/ Gradient across the Pulmonic valve : 0.83m/s 3mmHg

Max. Velocity / Gradient across the Aortic valve : 1.19m/s 4mmHg

Velocity / Gradient across the Tricuspid valve : 2.14m/s 15mmHg

SCAN FOR LOCATION



PATIENT NAME	MRS PADMAJA	ID NO	2402240031
AGE	24YEARS	SEX	FEMALE
REF BY	DR.APOLO CLINIC	DATE	24.02.2024

2D ECHO CARDIOGRAHIC STUDY

LEFT VENTRICLE	SIZE & THICKNESS	NORMAL
CONTRACTILITY	REGIONAL GLOBAL	NO RWMA

RIGHT VENTRICLE	: NORMAL
LEFT ATRIUM	: NORMAL
RIGHT ATRIUM	: NORMAL
MITRAL VALVE	: NORMAL
AORTIC VALVE	: NORMAL
PULMONARY VALVE	: NORMAL
TRICUSPID VALVE	: NORMAL
INTER ATRIAL SEPTUM	: INTACT
INTER VENTRICULAR SEPTUM	: INTACT
PERICARDIUM	: NORMAL
OTHERS	: - NIL

IMPRESSION

- NO REGIONAL WALL MOTION ABNORMALITY PRESENT
- NORMAL VALVES AND DIMENSIONS
- NORMAL LV FUNCTION, LVEF- 60%
- TRIVIAL TR / NO PAH
- NO CLOT / VEGETATION / EFFUSION



The science of radiology is based upon interpretation of shadows of normal and abnormal tissues. This is neither complete nor accurate; hence, findings should always be interpreted in to the light of clinico-pathological correlation.



Name	: MRS. PADMAJA	UHIP	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 10:56 AM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Glycosylated Haemoglobin (HbA1c)-Whole Blood EDTA	6.50	%	Non diabetic adults :<5.7	HPLC
Glycosylated Haemoglobin (HbA1c)			At risk (Prediabetes) : 5.7 - 6.4	
			Diagnosing Diabetes >= 6.5	
			Diabetes	
			Excellent Control : 6-7	
			Fair to good Control : 7-8	
			Unsatisfactory Control :8-10	
			Poor Control >10	
Estimated Average Glucose(eAG)	139.84	mg/dL		Calculated


Note: 1. Since HbA1c reflects long term fluctuations in the blood glucose concentration, a diabetic patient who is recently under good control may still have a high concentration of HbA1c. Converse is true for a diabetic previously under good control but now poorly controlled.

2. Target goals of < 7.0 % may be beneficial in patients with short duration of diabetes, long life expectancy and no significant cardiovascular disease. In patients with significant complications of diabetes, limited life expectancy or extensive co-morbid conditions, targeting a goal of < 7.0 % may not be appropriate.

Comments: HbA1c provides an index of average blood glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycaemic control as compared to blood and urinary glucose determinations.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 11:52 AM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Urine Routine Examination-Urine				
Physical Examination				
Colour	Pale Yellow		Pale Yellow	Visual
Appearance	Clear		Clear	Visual
Reaction (pH)	5.5		5.0-7.5	Dipstick
Specific Gravity	1.025		1.000-1.030	Dipstick
Biochemical Examination				
Albumin	Negative		Negative	Dipstick/Precipitation
Glucose	Negative		Negative	Dipstick/Benedicts
Bilirubin	Negative		Negative	Dipstick/Fouchets
Ketone Bodies	Negative		Negative	Dipstick/Rotheras
Urobilinogen	Normal		Normal	Dipstick/Ehrlichs
Nitrite	Negative		Negative	Dipstick
Microscopic Examination				
Pus Cells	2-4	hpf	0.0-5.0	Microscopy
Epithelial Cells	1-2	hpf	0.0-10.0	Microscopy
RBCs	Absent	hpf	Absent	Microscopy
Casts	Absent		Absent	Microscopy
Crystals	Absent		Absent	Microscopy
Others	Absent		Absent	Microscopy

Comments: The kidneys help filtration of the blood by eliminating waste out of the body through urine. They also regulate water in the body by conserving electrolytes, proteins, and other compounds. But due to some conditions and abnormalities in kidney function, the urine may encompass some abnormal constituents, which are not normally present. A complete urine examination helps in detecting such abnormal constituents in urine. Several disorders can be detected by identifying and measuring the levels of such substances. Blood cells, bilirubin, bacteria, pus cells, epithelial cells may be present in urine due to kidney disease or infection. Routine urine examination helps to diagnose kidney diseases, urinary tract infections, diabetes and other metabolic disorders.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist

Page 2 of 14



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 11:58 AM
Reg. No.	: 2402240031		2402240031	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Complete Haemogram-Whole Blood EDTA				
Haemoglobin (HB)	10.90	g/dL	Male: 14.0-17.0 Female: 12.0-15.0 Newborn: 16.50 - 19.50	Spectrophotometer
Red Blood Cell (RBC)	4.18	million/cumm	3.50 - 5.50	Volumetric Impedance
Packed Cell Volume (PCV)	32.70	%	Male: 42.0-51.0 Female: 36.0-45.0	Electronic Pulse
Mean corpuscular volume (MCV)	78.30	fL	78.0- 94.0	Calculated
Mean corpuscular hemoglobin (MCH)	26.10	pg	27.50-32.20	Calculated
Mean corpuscular hemoglobin concentration (MCHC)	33.40	%	33.00-35.50	Calculated
Red Blood Cell Distribution Width SD (RDW-SD)	47.00	fL	40.0-55.0	Volumetric Impedance
Red Blood Cell Distribution CV (RDW-CV)	18.40	%	Male: 11.80-14.50 Female: 12.20-16.10	Volumetric Impedance
Mean Platelet Volume (MPV)	8.10	fL	8.0-15.0	Volumetric Impedance
Platelet	3.34	lakh/cumm	1.50-4.50	Volumetric Impedance
Platelet Distribution Width (PDW)	9.00	%	8.30 - 56.60	Volumetric Impedance
White Blood cell Count (WBC)	9140.00	cells/cumm	Male: 4000-11000 Female: 4000-11000 Children: 6000-17500 Infants : 9000-30000	Volumetric Impedance
Neutrophils	58.60	%	40.0-75.0	Light scattering/Manual
Lymphocytes	35.50	%	20.0-40.0	Light scattering/Manual
Eosinophils	0.80	%	0.0-8.0	Light scattering/Manual



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 11:58 AM
Reg. No.	: 2402240031	2402240031		Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Monocytes	5.10	%	0.0-10.0	Light scattering/Manual
Basophils	0.00	%	0.0-1.0	Light scattering/Manual
Absolute Neutrophil Count	5.35	10 ³ /uL	2.0- 7.0	Calculated
Absolute Lymphocyte Count	3.25	10 ³ /uL	1.0-3.0	Calculated
Absolute Monocyte Count	0.47	10 ³ /uL	0.20-1.00	Calculated
Absolute Eosinophil Count	70.00	cells/cumm	40-440	Calculated
Absolute Basophil Count	0.00	10 ³ /uL	0.0-0.10	Calculated
Erythrocyte Sedimentation Rate (ESR)	37	mm/hr	Female : 0.0-20.0 Male : 0.0-10.0	Westergren

Peripheral Smear Examination-Whole Blood EDTA

Method: (Microscopy-Manual)

RBC'S : are microcytic hypochromic. Poikilocytes like tear drop cells and pencil shaped cells are seen.
WBC'S : Are normal in total number, morphology and distribution.
Platelets : Adequate in number and normal in morphology.
No abnormal cells or hemoparasites are present.

Impression : Mild degree of microcytic hypochromic anaemia.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist

Page 4 of 14



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 12:49 PM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
LFT-Liver Function Test -Serum				
Bilirubin Total-Serum	0.49	mg/dL	0.2-1.0	Caffeine Benzoate
Bilirubin Direct-Serum	0.11	mg/dL	0.0-0.2	Diazotised Sulphanilic Acid
Bilirubin Indirect-Serum	0.38	mg/dL	Female: 0.0 - 1.10	Direct Measure
Aspartate Aminotransferase (AST/SGOT)-Serum	15.00	U/L	Female: 15.0 - 37.0	UV with Pyridoxal - 5 - Phosphate
Alanine Aminotransferase (ALT/SGPT)-Serum	16.00	U/L	Female: 14.0 - 59.0	UV with Pyridoxal - 5 - Phosphate
Alkaline Phosphatase (ALP)-Serum	81.00	U/L	Female: 45.0 - 117.0	PNPP,AMP-Buffer
Protein, Total-Serum	6.70	g/dL	6.40-8.20	Biuret/Endpoint-With Blank
Albumin-Serum	3.94	g/dL	Female: 3.40 - 5.50	Bromocresol Purple
Globulin-Serum	2.76	g/dL	2.0-3.50	Calculated
Albumin/Globulin Ratio-Serum	1.43	Ratio	0.80-2.0	Calculated
Gamma-Glutamyl Transferase (GGT)-Serum	12.00	U/L	Male: 15.0-85.0 Female: 5.0-55.0	Other g-Glut-3-carboxy-4 nitro



Name	: MRS. PADMAJA	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female	UHD	: 2402240031
Ref. By Dr.	: Dr. APOLO CLINIC		Sample Col. Date : 24-Feb-2024 08:46 AM
Reg. No.	: 2402240031	2402240031	Result Date : 24-Feb-2024 12:49 PM
C/o	: Apollo Clinic		Report Status : Final

Test Name	Result	Unit	Reference Value	Method
-----------	--------	------	-----------------	--------

Comments: Gamma-glutamyltransferase (GGT) is primarily present in kidney, liver, and pancreatic cells. Small amounts are present in other tissues. Even though renal tissue has the highest level of GGT, the enzyme present in the serum appears to originate primarily from the hepatobiliary system, and GGT activity is elevated in any and all forms of liver disease. It is highest in cases of intra- or posthepatic biliary obstruction, reaching levels some 5 to 30 times normal. GGT is more sensitive than alkaline phosphatase (ALP), leucine aminopeptidase, aspartate transaminase, and alanine aminotransferase in detecting obstructive jaundice, cholangitis, and cholecystitis; its rise occurs earlier than with these other enzymes and persists longer. Only modest elevations (2-5 times normal) occur in infectious hepatitis, and in this condition, GGT determinations are less useful diagnostically than are measurements of the transaminases. High elevations of GGT are also observed in patients with either primary or secondary (metastatic) neoplasms. Elevated levels of GGT are noted not only in the sera of patients with alcoholic cirrhosis but also in the majority of sera from persons who are heavy drinkers. Studies have emphasized the value of serum GGT levels in detecting alcohol-induced liver disease. Elevated serum values are also seen in patients receiving drugs such as phenytoin and phenobarbital, and this is thought to reflect induction of new enzyme activity.



Printed By : spectrum
 Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 12:49 PM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Lipid Profile-Serum				
Cholesterol Total-Serum	166.00	mg/dL	Female: 0.0 - 200	Cholesterol Oxidase/Peroxidase
Triglycerides-Serum	92.00	mg/dL	Female: 0.0 - 150	Lipase/Glycerol Dehydrogenase
High-density lipoprotein (HDL) Cholesterol-Serum	47.00	mg/dL	Female: 40.0 - 60.0	Accelerator/Selective Detergent
Non-HDL cholesterol-Serum	119	mg/dL	Female: 0.0 - 130	Calculated
Low-density lipoprotein (LDL) Cholesterol-Serum	106.00	mg/dL	Female: 0.0 - 100.0	Cholesterol esterase and cholesterol oxidase
Very-low-density lipoprotein (VLDL) cholesterol-Serum	18	mg/dL	Female: 0.0 - 40	Calculated
Cholesterol/HDL Ratio-Serum	3.53	Ratio	Female: 0.0 - 5.0	Calculated

Interpretation:

Parameter	Desirable	Borderline High	High	Very High
Total Cholesterol	<200	200-239	>240	
Triglycerides	<150	150-199	200-499	>500
Non-HDL cholesterol	<130	160-189	190-219	>220
Low-density lipoprotein (LDL) Cholesterol	<100	100-129	160-189	>190

Comments: As per Lipid Association of India (LAI), for routine screening, overnight fasting preferred but not mandatory. Indians are at very high risk of developing Atherosclerotic Cardiovascular (ASCVD). Among the various risk factors for ASCVD such as dyslipidemia, Diabetes Mellitus, sedentary lifestyle, Hypertension, smoking etc., dyslipidemia has the highest population attributable risk for MI both because of direct association with disease pathogenesis and very high prevalence in Indian population. Hence monitoring lipid profile regularly for effective management of dyslipidemia remains one of the most important healthcare targets for prevention of ASCVD. In addition, estimation of ASCVD risk is an essential, initial step in the management of individuals requiring primary prevention of ASCVD. In the context of lipid management, such a risk estimate forms the basis for several key therapeutic decisions, such as the need for and aggressiveness of statin therapy.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithin Reddy C,MD,Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 12:49 PM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
KFT (Kidney Function Test) :				
Blood Urea Nitrogen (BUN)-Serum	7.00	mg/dL	7.0-18.0	GLDH,Kinetic Assay
Creatinine-Serum	0.58	mg/dL	Male: 0.70-1.30 Female: 0.55-1.02	Modified kinetic Jaffe
Uric Acid-Serum	5.01	mg/dL	Male: 3.50-7.20 Female: 2.60-6.00	Uricase PAP
Sodium (Na+)-Serum	140.6	mmol/L	135.0-145.0	Ion-Selective Electrodes (ISE)
Potassium (K+)-Serum	4.81	mmol/L	3.5 to 5.5	Ion-Selective Electrodes (ISE)
Chloride(Cl-)-Serum	100.00	mmol/L	96.0-108.0	Ion-Selective Electrodes (ISE)
Calcium,Total- Serum	9.20	mg/dL	8.50-10.10	Spectrophotometry (O-Cresolphthalein complexone)



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nitin Reddy C,MD,Consultant Pathologist



SCAN FOR LOCATION

Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 12:49 PM
Reg. No.	: 2402240031		2402240031	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Fasting Blood Sugar (FBS)- Plasma	81	mg/dL	60.0-110.0	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nitrum Reddy C.MD, Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 12:49 PM
Reg. No.	: 2402240031		2402240031	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Thyroid function tests (TFT)- Serum				
Tri-Iodo Thyronine (T3)-Serum	1.45	ng/mL	Female: 0.60 - 1.81	Chemiluminescence Immunoassay (CLIA)
Thyroxine (T4)-Serum	9.4	µg/dL	Female: 5.50 - 12.10	Chemiluminescence Immunoassay (CLIA)
Thyroid Stimulating Hormone (TSH)-Serum	5.87	µIU/mL	Female: 0.35 - 5.50	Chemiluminescence Immunoassay (CLIA)

Comments:Triiodothyronine (T3) assay is a useful test for hyperthyroidism in patients with low TSH and normal T4 levels. It is also used for the diagnosis of T3 toxicosis. It is not a reliable marker for Hypothyroidism. This test is not recommended for general screening of the population without a clinical suspicion of hyperthyroidism.

Reference range: Cord: (37 Weeks): 0.5-1.41, Children:1-3 Days: 1.0-7.40,1-11 Months: 1.05-2.45,1-5 Years: 1.05-2.69,6-10 Years: 0.94-2.41,11-15 Years: 0.82-2.13,Adolescents (16-20 Years): 0.80-2.10

Reference range: Adults: 20-50 Years: 0.70-2.04, 50-90 Years: 0.40-1.81,

Reference range in Pregnancy: First Trimester : 0.81-1.90,Second Trimester : 1.0-2.60

Increased Levels: Pregnancy, Graves disease, T3 thyrotoxicosis, TSH dependent Hyperthyroidism, increased Thyroid-binding globulin (TBG).

Decreased Levels: Nonthyroidal illness, hypothyroidism, nutritional deficiency, systemic illness, decreased Thyroid-binding globulin (TBG).

Comments:Total T4 levels offer a good index of thyroid function when TBG is normal and non-thyroidal illness is not present. This assay is useful for monitoring treatment with synthetic hormones (synthetic T3 will cause low total T4).It also helps to monitor treatment of Hyperthyroidism with Thiouracil or other anti-thyroid drugs.

Reference Range: Males : 4.6-10.5,Females : 5.5-11.0,> 60 Years: 5.0-10.70,Cord :7.40-13.10,Children:1-3 Days :11.80-22.60,1-2 Weeks : 9.90-16.60,1-4 Months: 7.20-14.40,1-5 Years : 7.30-15.0,5-10 Years: 6.4-13.3

1-15 Years: 5.60-11.70,Newborn Screen:1-5 Days: >7.5,6 Days : >6.5

Increased Levels: Hyperthyroidism, increased TBG, familial dysalbuminemic hyperthyroxinemia,Increased transthyretin, estrogen therapy, pregnancy.

Decreased Levels: Primary hypothyroidism, pituitary TSH deficiency, hypothalamic TRH deficiency, non thyroidal illness, decreased TBG.

Comments:TSH is a glycoprotein hormone secreted by the anterior pituitary. TSH is a labile hormone & is secreted in a pulsatile manner throughout the day and is subject to several non-thyroidal pituitary influences. Significant variations in TSH can occur with circadian rhythm, hormonal status, stress, sleep deprivation, caloric intake, medication & circulating antibodies. It is important to confirm any TSH abnormality in a fresh specimen drawn after ~ 3 weeks before assigning a diagnosis, as the cause of an isolated TSH abnormality.

Reference range in Pregnancy: I- trimester:0.1-2.5; II -trimester:0.2-3.0; III- trimester:0.3-3.0


Reference range in Newborns: 0-4 days: 1.0-39.0; 2-20 Weeks:1.7-9.1

Increased Levels: Primary hypothyroidism, Subclinical hypothyroidism, TSH dependent Hyperthyroidism and Thyroid hormone resistance.

Decreased Levels: Graves disease, Autonomous thyroid hormone secretion, TSH defici



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithun Reddy C,MD,Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 01:21 PM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Fasting Urine Glucose-Urine	Negative		Negative	Dipstick/Benedicts (Manual)



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nitium Reddy C, MD, Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 02:39 PM
Reg. No.	: 2402240031			Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Blood Group & Rh Typing-Whole Blood EDTA				
Blood Group	O			Slide/Tube agglutination
Rh Type	Positive			Slide/Tube agglutination

Note: Confirm by tube or gel method.

Comments: ABO blood group system, the classification of human blood based on the inherited properties of red blood cells (erythrocytes) as determined by the presence or absence of the antigens A and B, which are carried on the surface of the red cells. Persons may thus have type A, type B, type O, or type AB blood.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithum Reddy C.MD, Consultant Pathologist



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 03:37 PM
Reg. No.	: 2402240031	2402240031		Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Post Prandial Urine Sugar	Negative		Negative	Dipstick/Benedicts(Manual)



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nitin Reddy C, MD, Consultant Pathologist

Page 13 of 14



Name	: MRS. PADMAJA	UHID	: 2402240031	Bill Date	: 24-Feb-2024 08:46 AM
Age / Gender	: 24 years / Female			Sample Col. Date	: 24-Feb-2024 08:46 AM
Ref. By Dr.	: Dr. APOLO CLINIC			Result Date	: 24-Feb-2024 03:45 PM
Reg. No.	: 2402240031		2402240031	Report Status	: Final
C/o	: Apollo Clinic				

Test Name	Result	Unit	Reference Value	Method
Post prandial Blood Glucose (PPBS)-Plasma	88	mg/dL	70-140	Hexo Kinase

Comments: Glucose, also called dextrose, one of a group of carbohydrates known as simple sugars (monosaccharides). Glucose has the molecular formula $C_6H_{12}O_6$. It is found in fruits and honey and is the major free sugar circulating in the blood of higher animals. It is the source of energy in cell function, and the regulation of its metabolism is of great importance (fermentation; gluconeogenesis). Molecules of starch, the major energy-reserve carbohydrate of plants, consist of thousands of linear glucose units. Another major compound composed of glucose is cellulose, which is also linear. Dextrose is the molecule D-glucose. Blood sugar, or glucose, is the main sugar found in the blood. It comes from the food you eat, and it is body's main source of energy. The blood carries glucose to all of the body's cells to use for energy. Diabetes is a disease in which your blood sugar levels are too high. Usage: Glucose determinations are useful in the detection and management of Diabetes mellitus.

Note: Additional tests available for Diabetic control are Glycated Hemoglobin (HbA1c), Fructosamine & Microalbumin urine

Comments: Conditions which can lead to lower postprandial glucose levels as compared to fasting glucose are excessive insulin release, rapid gastric emptying & brisk glucose absorption.

Probable causes : Early Type II Diabetes / Glucose intolerance, Drugs like Salicylates, Beta blockers, Pentamidine etc., Alcohol ,Dietary – Intake of excessive carbohydrates and foods with high glycemic index ? Exercise in between samples ? Family history of Diabetes, Idiopathic, Partial / Total Gastrectomy.



Printed By : spectrum
Printed On : 24 Feb, 2024 04:18 pm



Dr. Nithun Reddy C, MD, Consultant Pathologist



SCAN FOR LOCATION