PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 11/03/2023 12:34 PM

Printed On : 17/03/2023 6:16 PM

Ref. Dr : MediWheel

: OP

Type



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING (Blood/Agglutination)	'B' 'Positive'		
Complete Blood Count With - ESR			
Haemoglobin (Blood/Spectrophotometry)	15.5	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (Blood/Numeric Integration of MCV)	46.9	%	42 - 52
RBC Count (Blood/Electrical Impedance)	5.19	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (Blood/Calculated)	90.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (Blood/Calculated)	29.9	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (Blood/Calculated)	33.1	g/dL	32 - 36
RDW-CV (Calculated)	13.9	%	11.5 - 16.0
RDW-SD (Calculated)	43.98	fL	39 - 46
Total Leukocyte Count (TC) (Blood/Electrical Impedance)	7210	cells/cu.mm	4000 - 11000
Neutrophils (Blood/Impedance and absorbance)	55.19	%	40 - 75
Lymphocytes (Blood/Impedance and absorbance)	32.85	%	20 - 45
Eosinophils (Blood/Impedance and absorbance)	2.70	%	01 - 06







APPROVED BY

The results pertain to sample tested.

Page 1 of 9

 PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 11/03/2023 12:34 PM

Printed On

medall DIAGNOSTICS

Ref. Dr : MediWheel

: OP

Type

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
Monocytes (Blood/Impedance and absorbance)	8.91	%	01 - 10
Basophils (Blood/Impedance and absorbance)	0.36	%	00 - 02
INTERPRETATION: Tests done on Automated F	ive Part cell counte	er. All abnormal results are review	ved and confirmed microscopically.
Absolute Neutrophil count (Blood/Impedance and absorbance)	3.98	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (Blood/Impedance)	2.37	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (Blood/Impedance)	0.19	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (Blood/Impedance)	0.64	10^3 / μl	< 1.0
Absolute Basophil count (Blood/Impedance)	0.03	10^3 / μl	< 0.2
Platelet Count (Blood/Impedance)	3.6	lakh/cu.mm	1.4 - 4.5
INTERPRETATION: Platelet count less than 1.5	akhs will be confir	med microscopically.	
MPV (Blood/Derived from Impedance)	8.26	fL	7.9 - 13.7
PCT (Calculated)	0.30	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated ESR analyser)	18	mm/hr	< 15
BUN / Creatinine Ratio	9.7		
Glucose Fasting (FBS) (Plasma - F/Glucose oxidase/Peroxidase)	95	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

: 17/03/2023 6:16 PM







APPROVED BY

 PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 11/03/2023 12:34 PM

Ref. Dr : MediWheel



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
INTERPRETATION: Factors such as type, blood glucose level.	quantity and time of food	d intake, Physical activ	ity, Psychological stress, and drugs can influence
Glucose, Fasting (Urine) (Urine - F)	Negative		Negative
Glucose Postprandial (PPBS)	111	mg/dL	70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Calculated)	8.8	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe ~Alkaline Picrate)	1	mg/dL	0.9 - 1.3
Uric Acid (Serum/Uricase/Peroxidase)	5.9	mg/dL	3.5 - 7.2
Liver Function Test			
Bilirubin(Total) (Serum/Diazotized Sulphanilic acid)	0.9	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulphanilic acid)	0.3	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Calculated)	0.60	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/IFCC without P-5-P)	20	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/IFCC without P-5-P)	36	U/L	5 - 41







APPROVED BY

The results pertain to sample tested.

Page 3 of 9

 PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 11/03/2023 12:34 PM

Type : OP Printed On : 17/03/2023 6:16 PM

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Alkaline Phosphatase (SAP) (Serum/IFCC AMP Buffer)	114	U/L	53 - 128
Total Protein (Serum/Biuret)	7.1	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.7	gm/dl	3.5 - 5.2
Globulin (Serum/Calculated)	3.40	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Calculated) INTERPRETATION: Enclosure : Graph	1.09		1.1 - 2.2
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	29	U/L	< 55
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	213	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol-phosphate oxidase/Peroxidase)	144	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.









APPROVED BY

PID No. : MED111534820 Register On : 11/03/2023 9:09 AM : 80025395 SID No. Collection On : 11/03/2023 9:59 AM

: 11/03/2023 12:34 PM **Type** : OP **Printed On** : 17/03/2023 6:16 PM

Report On

Ref. Dr : MediWheel

Age / Sex : 38 Year(s) / Male



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
HDL Cholesterol (Serum/Immunoinhibition)	52	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	132.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	28.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	161.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

4.1 Optimal: < 3.3 Total Cholesterol/HDL Cholesterol Low Risk: 3.4 - 4.4 Ratio Average Risk: 4.5 - 7.1 (Serum/Calculated) Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 2.8 Optimal: < 2.5Mild to moderate risk: 2.5 - 5.0 (TG/HDL) High Risk: > 5.0(Serum/Calculated)







APPROVED BY

 PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

Age / Sex: 38 Year(s) / Male **Report On**: 11/03/2023 12:34 PM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.5		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/ <i>HPLC-Ion exchange</i>)	5.4	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Mean Blood Glucose 108.28 mg/dl

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.59 ng/ml 0.7 - 2.04

 $(Serum/{\it Chemiluminescent\ Immunometric\ Assay}$

(CLIA))

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 11.09 µg/dl 4.2 - 12.0

(Serum/Chemiluminescent Immunometric Assay

Lab Manager

VERIFIED BY

(CLIA))





APPROVED BY

The results pertain to sample tested.

Page 6 of 9

 PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

Age / Sex: 38 Year(s) / Male **Report On**: 11/03/2023 12:34 PM

Type : OP Printed On : 17/03/2023 6:16 PM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.97 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescence)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.

2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.

3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

Urine Analysis - Routine

Others

(Urine/Microscopy)

INTERPRETATION: Note: Done with Automated Urine Analyser & microscopy

Physical Examination(Urine Routine)

Colour PALE YELLOW Yellow to Amber

(Urine/Physical examination)

Appearance Clear Clear

(Urine/Physical examination)

<u>Chemical Examination(Urine Routine)</u>

Protein Negative Negative

(Urine/Dipstick-Error of indicator/ Sulphosalicylic acid method)









APPROVED BY

The results pertain to sample tested.

Page 7 of 9

 PID No.
 : MED111534820
 Register On
 : 11/03/2023 9:09 AM

 SID No.
 : 80025395
 Collection On
 : 11/03/2023 9:59 AM

 Age / Sex
 : 38 Year(s) / Male
 Report On
 : 11/03/2023 12:34 PM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine/Dip Stick Method / Glucose Oxidase - Peroxidase / Benedict š semi quantitative method.)	Negative		Negative
<u>Microscopic Examination(Urine</u> Routine)			
Pus Cells (Urine/Microscopy exam of urine sediment)	3-4	/hpf	0 - 5
Epithelial Cells (Urine/Microscopy exam of urine sediment)	1-2	/hpf	0 - 5
RBCs (Urine/Microscopy exam of urine sediment)	NIL	/hpf	0 - 5
PHYSICAL EXAMINATION			
PHYSICAL EXAMINATION Colour (Stool/Physical examination)	Brown		Brown
Colour (Stool/Physical examination) Consistency	Brown SEMI SOFT		Brown Well Formed
Colour			
Colour (Stool/Physical examination) Consistency (Stool/Physical examination) Mucus	SEMI SOFT		Well Formed



Reaction

(Stool)



Acidic



Acidic

APPROVED BY

The results pertain to sample tested.

Page 8 of 9

PID No. : MED111534820 **Register On** : 11/03/2023 9:09 AM : 80025395 SID No. Collection On : 11/03/2023 9:59 AM Age / Sex : 38 Year(s) / Male

Report On : 11/03/2023 12:34 PM

Type : OP **Printed On** : 17/03/2023 6:16 PM

Ref. Dr : MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
MICROSCOPIC EXAMINATION (STOOL COMPLETE)			
Ova (Stool)	NIL		
Cysts (Stool)	NIL		
Trophozoites (Stool)	NIL		
Pus Cells (Stool)	0-2	/hpf	
RBCs (Stool)	NIL	/hpf	
Others (Stool)	NIL		







APPROVED BY

-- End of Report --



Name	PEDADA VENKATA RAO	ID	MED111534820
Age & Gender	38Y/M	Visit Date	Mar 11 2023 9:09AM
Ref Doctor	MediWheel		

ULTRASOUND WHOLE ABDOMEN

Liver : Normal in size (14.2 cm) with regular outlines and normal

echopattern.

There is no evidence of IHBR / EHBR dilatation seen.

No focal space occupying lesions seen.

CBD is normal. PV normal.

Gall Bladder : Normal in volume and wall thickness.

No e/o intraluminal calculi seen.

Pancreas : Head, body and tail are identified with normal echopattern and

smooth outlines.

Spleen : Measured 9.1 cm, in size with normal echotexture.

Right kidney : Measured 9.6 x 5.1 cm in size.

Left kidney : Measured 9.5 x 4.8 cm in size.

Both kidneys are normal in size, position, with well preserved cortico medullary differentiation and normal pelvicalyceal

anatomy.

No e/o calculi / space occupying lesion seen. No e/o suprarenal / retroperitoneal masses noted.

Urinary : Normal in volume and wall thickness.
bladder : No e/o intraluminal calculi / masses seen.

Prostate : Measured 2.9 x 3.1 x 2.8 cm in size (Vol : 13.4 cc) with normal

echotexture.

No e/o ascites / pleural effusion seen. No e/o detectable bowel pathology seen.

IMPRESSION:

• Essentially normal study.



Name	PEDADA VENKATA RAO	ID	MED111534820
Age & Gender	38Y/M	Visit Date	Mar 11 2023 9:09AM
Ref Doctor	MediWheel		

- For clinical correlation.

Dr.Jahn av i Barla ,MD (RD)

Consultant Radiologist