

CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251 CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00

RECEIVED : 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status Results Biological Reference Interval Units <u>Final</u>

н	AEMATOLOGY - CBC		
MEDI WHEEL FULL BODY HEALTH CHECKUP BE	LOW 40FEMALE		
BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB) METHOD: CYANIDE FREE DETERMINATION	12.7	12.0 - 15.0	g/dL
RED BLOOD CELL (RBC) COUNT METHOD: ELECTRICAL IMPEDANCE	4.49	3.8 - 4.8	mi l /μL
WHITE BLOOD CELL (WBC) COUNT METHOD: ELECTRICAL IMPEDANCE	6.90	4.0 - 10.0	thou/μL
PLATELET COUNT METHOD: ELECTRONIC IMPEDANCE	319	150 - 410	thou/μL
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV) METHOD: CALCULATED PARAMETER	39.1	36 - 46	%
MEAN CORPUSCULAR VOLUME (MCV) METHOD: CALCULATED PARAMETER	87.0	83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH) METHOD: CALCULATED PARAMETER	28.2	27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) METHOD: CALCULATED PARAMETER	32.4	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW) METHOD: CALCULATED PARAMETER	12.5	11.6 - 14.0	%
MENTZER INDEX	19.4		
MEAN PLATELET VOLUME (MPV) METHOD: CALCULATED PARAMETER	8.6	6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT			
NEUTROPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	57	40 - 80	%
LYMPHOCYTES METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	37	20 - 40	%
MONOCYTES METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	04	2 - 10	%
EOSINOPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	02	1 - 6	%



Dr. Akansha Jain **Consultant Pathologist**





Page 1 Of 15









CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00 RECEIVED : 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
PAGORUNG	00	0 0	0/
BASOPHILS METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY	00	0 - 2	%
ABSOLUTE NEUTROPHIL COUNT METHOD: CALCULATED PARAMETER	3.93	2.0 - 7.0	thou/µL
ABSOLUTE LYMPHOCYTE COUNT METHOD: CALCULATED PARAMETER	2.55	1.0 - 3.0	thou/µL
ABSOLUTE MONOCYTE COUNT METHOD: CALCULATED PARAMETER	0.28	0.2 - 1.0	thou/µL
ABSOLUTE EOSINOPHIL COUNT METHOD: CALCULATED PARAMETER	0.14	0.02 - 0.50	thou/µL
ABSOLUTE BASOPHIL COUNT	0 Low	0.02 - 0.10	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	1.5		

from Beta thalassaemia trait

Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

Dr. Akansha Jain **Consultant Pathologist** Page 2 Of 15







View Report



CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00 RECEIVED: 05/02/2023 10:41:06

REPORTED :05/02/2023 16:14:42

Biological Reference Interval Units Test Report Status Results <u>Final</u>

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD

E.S.R 0 - 20mm at 1 hr

METHOD: AUTOMATED (PHOTOMETRICAL CAPILLARY STOPPED FLOW KINETIC ANALYSIS)"

Interpretation(s)

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays' fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging,

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia False Decreased: Poikilocytosis,(SickleCells,spherocytes),Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, salicylates)

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

Dr. Akansha Jain **Consultant Pathologist**



View Report

Page 3 Of 15



CODE/NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00 RECEIVED : 05/02/2023 10:41:06

REPORTED :05/02/2023 16:14:42

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

ABHA NO

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE A

METHOD: TUBE AGGLUTINATION

POSITIVE RH TYPE

METHOD: TUBE AGGLUTINATION

Interpretation(s)

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.'

The test is performed by both forward as well as reverse grouping methods.

Dr. Akansha Jain **Consultant Pathologist**



Page 4 Of 15





CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251 CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00 RECEIVED: 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR) 108 High 74 - 99 mg/dL

METHOD: GLUCOSE OXIDASE

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE

BLOOD

HBA1C 6.0 High Non-diabetic: < 5.7 %

> Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5Therapeutic goals: < 7.0 Action suggested: > 8.0 (ADA Guideline 2021)

METHOD: HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)

ESTIMATED AVERAGE GLUCOSE(EAG) 125.5 High < 116.0 mg/dL

METHOD: CALCULATED PARAMETER

GLUCOSE, POST-PRANDIAL, PLASMA

184 High PPBS(POST PRANDIAL BLOOD SUGAR) 70 - 140mg/dL

 ${\tt METHOD}: {\tt GLUCOSE} \ {\tt OXIDASE}$ LIPID PROFILE, SERUM

208 High

< 200 Desirable mg/dL CHOLESTEROL, TOTAL

200 - 239 Borderline High >/= 240 High

METHOD: CHOLESTEROL OXIDASE

162 High TRIGLYCERIDES < 150 Normal mg/dL

150 - 199 Borderline High

200 - 499 High >/=500 Very High

METHOD: LIPASE/GPO-PAP NO CORRECTION

HDL CHOLESTEROL mg/dL 46 < 40 Low

>/=60 High

METHOD: DIRECT CLEARANCE METHOD

Dr. Akansha Jain **Consultant Pathologist**



Page 5 Of 15







CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: **0251WB000330**PATIENT ID: SWATF050293251
CLIENT PATIENT ID: 012302050010

ABHA NO :

AGE/SEX :30 Years Female
DRAWN :05/02/2023 08:57:00
RECEIVED :05/02/2023 10:41:06
REPORTED :05/02/2023 16:14:42

Test Report Status <u>Final</u>	Results	Biological Reference Interva	l Units
CHOLESTEROL LDL	130 High	< 100 Optimal 100 - 129 Near optimal/ above optima 130 - 159 Borderline High 160 - 189 High >/= 190 Very High	mg/dL il
NON HDL CHOLESTEROL	162 High	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
METHOD: CALCULATED PARAMETER		, 3	
VERY LOW DENSITY LIPOPROTEIN	32 . 4 High	= 30.0</td <td>mg/dL</td>	mg/dL
CHOL/HDL RATIO	4.5 High	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk	
LDL/HDL RATIO Interpretation(s)	2.8	0.5 - 3.0 Desirable/Low Risl 3.1 - 6.0 Borderline/Modera Risk >6.0 High Risk	
LIVER FUNCTION PROFILE, SERUM			
BILIRUBIN, TOTAL METHOD: DIAZO WITH SULPHANILIC ACID	0.36	0 - 1	mg/dL
BILIRUBIN, DIRECT METHOD: DIAZO WITH SULPHANILIC ACID	0.10	0.00 - 0.25	mg/dL
BILIRUBIN, INDIRECT METHOD: CALCULATED PARAMETER	0.26	0.1 - 1.0	mg/dL
TOTAL PROTEIN METHOD: BIURET REACTION, END POINT	7.6	6.4 - 8.2	g/dL
ALBUMIN	4.3	3.8 - 4.4	g/dL







Page 6 Of 15













CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: **0251WB000330**PATIENT ID: SWATF050293251
CLIENT PATIENT ID: 012302050010

ABHA NO :

AGE/SEX :30 Years Female
DRAWN :05/02/2023 08:57:00
RECEIVED :05/02/2023 10:41:06
REPORTED :05/02/2023 16:14:42

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units		
METHOD: BROMOCRESOL GREEN				
GLOBULIN	3.3	2.0 - 4.1	g/dL	
METHOD : CALCULATED PARAMETER	1.2	10.21	DATIO	
ALBUMIN/GLOBULIN RATIO METHOD: CALCULATED PARAMETER	1.3	1.0 - 2.1	RATIO	
ASPARTATE AMINOTRANSFERASE (AST/SGOT) METHOD: TRIS BUFFER NO P5P IFCC / SFBC 37° C	23	0 - 31	U/L	
ALANINE AMINOTRANSFERASE (ALT/SGPT)	30	0 - 31	U/L	
METHOD : TRIS BUFFER NO P5P IFCC / SFBC 37° C				
ALKALINE PHOSPHATASE METHOD: AMP OPTIMISED TO IFCC 37° C	64	39 - 117	U/L	
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: GAMMA GLUTAMYL-3 CARBOXY-4 NITROANILIDE (IFO	31 CC) 37° C	7 - 32	U/L	
LACTATE DEHYDROGENASE	366	230 - 460	U/L	
BLOOD UREA NITROGEN (BUN), SERUM				
BLOOD UREA NITROGEN	12	5.0 - 18.0	mg/dL	
METHOD : UREASE KINETIC			-	
CREATININE, SERUM				
CREATININE	0.89	0.6 - 1.2	mg/dL	
METHOD : ALKALINE PICRATE NO DEPROTEINIZATION				
BUN/CREAT RATIO				
BUN/CREAT RATIO	13.48			
METHOD: CALCULATED PARAMETER				
URIC ACID, SERUM				
URIC ACID	6.2 High	2.4 - 5.7	mg/dL	
METHOD: URICASE PEROXIDASE WITH ASCORBATE OXIDASE				
TOTAL PROTEIN, SERUM				
TOTAL PROTEIN	7.6	6.4 - 8.3	g/dL	
METHOD: BIURET REACTION, END POINT				
ALBUMIN, SERUM	4.0	20 44	7.11	
ALBUMIN	4.3	3.8 - 4.4	g/dL	
METHOD : BROMOCRESOL GREEN GLOBULIN				
	2.2	2.0 4.1	a/dl	
GLOBULIN	3.3	2.0 - 4.1	g/dL	



Dr. Akansha Jain Consultant Pathologist





Page 7 Of 15

View Details











REF. DOCTOR: SELF **PATIENT NAME: SWATI SINGH**

CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251 CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00 RECEIVED: 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status <u>Final</u>	Results	Biological Reference	e Interval Units
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM, SERUM METHOD: ION-SELECTIVE ELECTRODE	139.8	137 - 145	mmo l /L
POTASSIUM, SERUM METHOD: ION-SELECTIVE ELECTRODE	4.16	3.6 - 5.0	mmo l /L
CHLORIDE, SERUM METHOD: ION-SELECTIVE ELECTRODE	103.3	98 - 107	mmol/L

Interpretation(s)

Interpretation(s)

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the urine.

Increased in

Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides. **Decreased in**

Pancreatic islet cell disease with increased insulin,insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia), Drugs- insulin, ethanol, propranolol; sulfonylureas, tolbutamide, and other oral hypoglycemic agents.

while random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- Diagnosing diabetes.
- 3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- 1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
 2. eAG gives an evaluation of blood glucose levels for the last couple of months.
 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to:

I.Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss,hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II.Vitamin C & E are reported to falsely lower test results (possibly by inhibiting glycation of hemoglobin.

III.Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.

IV.Interference of hemoglobinopathies in HbA1c estimation is seen in a.Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.

b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-LIVER FUNCTION PROFILE

Dr. Akansha Jain **Consultant Pathologist**





Page 8 Of 15

View Report







CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00 RECEIVED: 05/02/2023 10:41:06

REPORTED :05/02/2023 16:14:42

Results **Test Report Status** Biological Reference Interval Units <u>Fina</u>

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin wiral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that

attaches sugar molecules to bilirubin.
AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys,heart,muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget''''s disease,Rickets,Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia,Malnutrition,Protein deficiency,Wilson'''s disease.GGT is an enzyme found in cell membranes of many tissues mainly in the liver,kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom'''s disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.Human serum albumin is the most abundant protein in human blood plasma.It is produced in the liver Albumin constitutes about half of the blood serum protein.Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc
BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol,

Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism) Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

- Blockage in the urinary tract
 Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia GravisMuscular dystrophy

URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome

Causes of decreased levels-Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM-Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom""""""" disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum

protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

Dr. Akansha Jain Consultant Pathologist



Page 9 Of 15









CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: **0251WB000330**PATIENT ID: SWATF050293251

CLIENT PATIENT ID: 012302050010

AGE/SEX :30 Years Female
DRAWN :05/02/2023 08:57:00
RECEIVED :05/02/2023 10:41:06
REPORTED :05/02/2023 16:14:42

Test Report Status <u>Final</u> Results Biological Reference Interval Units

ABHA NO

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

METHOD: GROSS EXAMINATION

APPEARANCE HAZY

METHOD: GROSS EXAMINATION

CHEMICAL EXAMINATION, URINE

PH 5.5 4.7 - 7.5

METHOD: DOUBLE INDICATOR PRINCIPLE

SPECIFIC GRAVITY 1.025 1.003 - 1.035

METHOD: IONIC CONCENTRATION METHOD

PROTEIN NOT DETECTED NOT DETECTED

METHOD: PROTEIN ERROR OF INDICATORS WITH REFLECTANCE

GLUCOSE NOT DETECTED NOT DETECTED

METHOD: GLUCOSE OXIDASE PEROXIDASE / BENEDICTS

NOT DETECTED

METHOD: GLUCUSE UXIDASE PERUXIDASE / BENEDICTS

KETONES NOT DETECTED NOT DETECTED

 ${\tt METHOD: SODIUM\ NITROPRUSSIDE\ REACTION}$

BLOOD NOT DETECTED NOT DETECTED

METHOD: PEROCIDASE ANTI PEROXIDASE

BILIRUBIN NOT DETECTED NOT DETECTED

METHOD : DIPSTICK

UROBILINOGEN NORMAL NORMAL

METHOD: EHRLICH REACTION REFLECTANCE

NITRITE NOT DETECTED NOT DETECTED

METHOD: NITRATE TO NITRITE CONVERSION METHOD

LEUKOCYTE ESTERASE NOT DETECTED NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS NOT DETECTED NOT DETECTED /HPF

METHOD: MICROSCOPIC EXAMINATION

METHOD: MICROSCOPIC EXAMINATION

PUS CELL (WBC'S) 2-3 0-5 /HPF

METHOD: DIPSTICK, MICROSCOPY

EPITHELIAL CELLS **8-10** 0-5 /HPF

CASTS NOT DETECTED

Dr. Akansha Jain Consultant Pathologist Page 10 Of 15







View Report





CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

<u>Final</u>

JAIPUR 302017 9314660100

Test Report Status

ACCESSION NO: **0251WB000330**PATIENT ID: SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO :

AGE/SEX :30 Years Female
DRAWN :05/02/2023 08:57:00
RECEIVED :05/02/2023 10:41:06
REPORTED :05/02/2023 16:14:42

Results Biological Reference Interval Units

METHOD: MICROSCOPIC EXAMINATION

CRYSTALS NOT DETECTED

METHOD: MICROSCOPIC EXAMINATION

METHOD: MICROSCOPIC EXAMINATION

BACTERIA NOT DETECTED NOT DETECTED

YEAST NOT DETECTED NOT DETECTED

Interpretation(s)

Dr. Akansha Jain Consultant Pathologist





View Details





Page 11 Of 15



CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO : **0251WB000330**PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO :

AGE/SEX : 30 Years Female
DRAWN : 05/02/2023 08:57:00

RECEIVED : 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status Final Results Biological Reference Interval Units

CYTOLOGY

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PAPANICOLAOU SMEAR

TEST METHOD SAMPLE NOT RECEIVED

Dr. Akansha Jain Consultant Pathologist



Page 12 Of 15

view Details

View Report









PATIENT NAME: SWATI SINGH

CODE/NAME & ADDRESS: C000049066

SRL JAIPUR WELLNESS CORPORATE WALK IN
AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100 REF. DOCTOR: SELF

ACCESSION NO: **0251WB000330**PATIENT ID: SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00

RECEIVED : 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

PHYSICAL EXAMINATION, STOOL

COLOUR

METHOD: GROSS EXAMINATION

SAMPLE NOT RECEIVED

Dr. Abhishek Sharma Consultant Microbiologist



view Details

View Report





Page 13 Of 15



CODE/NAME & ADDRESS: C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female DRAWN :05/02/2023 08:57:00

RECEIVED: 05/02/2023 10:41:06 REPORTED :05/02/2023 16:14:42

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

THYROID PANEL, SERUM

T3 METHOD: CHEMILUMINESCENCE	95.76	60.0 - 181.0	ng/dL
T4	8.60	4.5 - 10.9	μg/dL
METHOD: CHEMILUMINESCENCE TSH (ULTRASENSITIVE) METHOD: CHEMILUMINESCENCE	4.282	0.550 - 4.780	μIU/mL

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. owidetlparowidetlparBelow mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism

Dr. Akansha Jain **Consultant Pathologist**



Page 14 Of 15











CODE/NAME & ADDRESS : C000049066 SRL JAIPUR WELLNESS CORPORATE WALK IN AAKRITI LABS PVT LTD. A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251WB000330 PATIENT ID : SWATF050293251

CLIENT PATIENT ID: 012302050010

ABHA NO

AGE/SEX :30 Years Female :05/02/2023 08:57:00 DRAWN RECEIVED : 05/02/2023 10:41:06

REPORTED :05/02/2023 16:14:42

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2 Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. NOTE: It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4. TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

> **End Of Report** Please visit www.srlworld.com for related Test Information for this accession

Dr. Akansha Jain **Consultant Pathologist**





Page 15 Of 15












Heart Rate: 87 bpm / Tested On: 05-Feb-23 12:35:56 / HF 0.05 Hz - LF 100 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s





Aakriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563

Scoadi, Shrith.

21/P/ 9988947273.

05/02/23

A) >.

616.

Colous. Blindness-Nosmer.

Dr. RAKESHASHARWA



akriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com CIN NO.: U85195RJ2004PTC019563

PATIENT NAME: MS SWATI SINGH	AGE & SEX: 30 Y/Female
REF. BY DR: MEDIWHEEL	DATE: 05.02.2023
•	

USG: WHOLE ABDOMEN (Female)

LIVER

: Is normal in size with bright echogenecity. The IHBR and hepatic radicals are not dilated.

No evidence of focal echopoor/echorich lesion seen. Portal vein diameter and Common bile duct normal in size

GALL

: Is normal in size, shape and echotexture. Walls are smooth and

BLADDER regular with normal thickness. There is no evidence of cholelithiasis.

PANCREAS: Is normal in size, shape and echotexture. Pancreatic duct is not dilated. SPLEEN: Is normal in size, shape and echogenecity. Spleenic hilum is not dilated.

KIDNEYS: Right Kidney:-Size: 95x38 mm, Left Kidney:-Size: 97x44 mm. Bilateral Kidneys are normal in size, shape and echotexture. corticomedullary differentiation is fair and ratio appears normal.

Pelvi calyceal system is normal. No evidence of hydronephrosis/ nephrolithiasis.

URINARY: Bladder walls are smooth regular and normal thickness.

BLADDER: No evidence of mass or stone in bladder lumen.

UTERUS: Uterus is anteverted with normal in size shape & echotexture.

Uterine muscular shadows normal echopattern.

Endometrium is normal and centrally placed with size: 7 mm. No evidence of mass lesion is seen. Size of uterus: 68x49x37 mm.

ADNEXA: Both the ovaries are normal in size shape and echotexture.

No mass lesion/ polycystic ovarian cyst is seen.

SPECIFIC: - Mild free fluid seen in pouch of dougls.

: No evidence of retroperitoneal mass or free fluid seen in peritoneal cavity.

: NO evidence of lymphadenopathy or mass lesion in retroperitoneum.

: Visualized bowel loop appear normal. Great vessels appear normal.

IMPRESSION: - Fatty liver

:- Free fluid in POD ? PID

DR NEERA MEHTA MBBS, DMRD RMCNO.005807/14853



Aakriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563



Name

: Mrs. SWATI SINGH

Age/Gender: 30 Y/Female
Patient ID : 012302050010

BarcodeNo: 10075363

Referred By: Self

Registration No: 51370

Registered

: 05/Feb/2023 08:57AM

Analysed

: 05/Feb/2023 10:27AM

Reported

: 05/Feb/2023 10:27AM

Panel

: Medi Wheel (ArcoFemi

Healthcare Ltd)

DIGITAL X-RAY CHEST PA VIEW

Soft tissue shadow and bony cages are normal.

Trachea is central.

Bilateral lung field and both CP angle are clear.

Domes of diaphragm are normally placed.

Transverse diameter of heart appears with normal limits.

IMPRESSION:- NO OBVIOUS ABNORMALITY DETECTED.

wenness partner

*** End Of Report ***

Page 1 of 1

Dr. Nee/ Mehta M.B.B.S., D.M.R.D. RMCNO.005807/14853



Aakriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563

NAME		MRS SWATI SINGH MEDIWHEEL			AGE	GE 30Y		SEX	FEMALE
REF BY	MED			DATE	DATE 05/02/2023		REG NO	LIVIALE	
			ECH	OCARDIOG	RAM RE				200
WINDO	N- POO	R/ADEQ	UATE/GO	ODVALVE					
MITRAL	12 125		NORMA		TRICU	SPID		NORMAI	<u>.</u>
AORTIC			NORMAI		THE RESERVE CO.	ONARY		NORMAL	
2D/M-M	OD							TO THE T	
IVSD mm		7.1		IVSS mm	12.9		AORTA	mm	24.4
LVID mm		34.2		LVIS mm	21.3		LA mm		25.0
LVPWD n	nm	8.5		LVPWS mm	11.8		EF%	0	60%
CHAMBE	RS						21.70		0070
LA			NC	RMAL	RA			NOR	ΜΔΙ
LV			NO	NORMAL		RV		NORMAL	
PERICARDIUM		NO	RMAL				IVOIN	VIAL	
DOPPLER	STUDY	MITRA	L						
PEAK VEL			1.4	1.42/1.04		GRADIANT	MmHg		
MEAN VE	LOCITY	m/s				MEAN GRADIANT MmHg		-	J
MVA cm2	(PLAN	ITMETER	RY)			MVA cm2 (PHT)			
MR				and the second		1000	- 1		
AORTIC					140	N. T.			
PEAK VEL			1.8	6	PEAK	GRADIANT	MmHg		No.
MEAN VE	LOCITY	m/s				MEAN GRADIANT MmHg			
AR				THE WORLD	CO. T. CO.				
TRICUSPI					and the second		Ware		
PEAK VELO		A Property Control	0.79		PEAK	GRADIANT	MmHg		
MEAN VEI	OCITY	m/s		A CONTRACTOR OF THE PARTY OF TH		MEAN GRADIANT MmHg			
TR				mmHg					
PULMONA	und later.		P					A) Fee	
PEAK VELO	OCITY n	n/s	1.61		PEAK	GRADIANT	MmHø		
MEAN VEL	OCITY	m/s		1000		GRADIANT			
PR				111		PmmHg			

IMPRESSION

- NORMAL LV SYSTOLIC & DIASTOLIC FUNCTION
- NO RWMA LVEF 60%
- NORMAL RV FUNCTION
- NORMAL CHAMBER DIMENSIONS
- NORMAL VALVULAR ECHO
- INTACT IAS / IVS
- NO THROMBUS, NO VEGETATION, NORMAL PERICARDIUM.
- IVC NORMAL

CONCLUSION: FAIR LV FUNCTION.

Cardiologist