



CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

PATIENT NAME: SHARAD THEKEDA

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

> PATIENT ID: SHARM020187290

ACCESSION NO: 0290VL001162 AGE: 35 Years SEX: Male ABHA NO:

RECEIVED: 10/12/2022 10:49 13/12/2022 13:57 DRAWN: REPORTED:

REFERRING DOCTOR: DR. BANK OF BARODA -CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

BLOOD COUNTS,EDTA WHOLE BLOOD				
HEMOGLOBIN (HB)	14.1		13.0 - 17.0	g/dL
RED BLOOD CELL (RBC) COUNT	5.09		4.5 - 5.5	mil/μL
WHITE BLOOD CELL (WBC) COUNT	7.40		4.0 - 10.0	thou/µL
PLATELET COUNT	328		150 - 410	thou/µL
RBC AND PLATELET INDICES				
HEMATOCRIT (PCV)	43.0		40 - 50	%
MEAN CORPUSCULAR VOLUME (MCV)	84.0		83 - 101	fL
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	27.7		27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	32.8		31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW)	12.9		11.6 - 14.0	%
MENTZER INDEX	16.5			
MEAN PLATELET VOLUME (MPV)	7.6		6.8 - 10.9	fL
WBC DIFFERENTIAL COUNT				
NEUTROPHILS	58		40 - 80	%
LYMPHOCYTES	33		20 - 40	%
MONOCYTES	04		2 - 10	%
EOSINOPHILS	05		1 - 6	%
BASOPHILS	00		0 - 2	%
ERYTHROCYTE SEDIMENTATION RATE (ESR) BLOOD),WHOLE			
E.S.R	15	High	0 - 14	mm at 1 hr
GLUCOSE FASTING, FLUORIDE PLASMA				
FBS (FASTING BLOOD SUGAR)	106	High	74 - 99	mg/dL
GLYCOSYLATED HEMOGLOBIN(HBA1C), EDT. BLOOD	A WHOLE			
HBA1C	5.6		Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 Therapeutic goals: < 7.0 Action suggested: > 8.0 (ADA Guideline 2021)	%
ESTIMATED AVERAGE GLUCOSE(EAG)	114.0		< 116.0	mg/dL



GLUCOSE, POST-PRANDIAL, PLASMA



Scan to View Report





SHARM020187290

CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA

8800465156

Opposite St Raphael's Higher Secondary School , Old Seshore Road,

PATIENT ID:

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

RECEIVED: 10/12/2022 10:49 13/12/2022 13:57 DRAWN: REPORTED:

CLIENT PATIENT ID: REFERRING DOCTOR: DR. BANK OF BARODA -

Test Report Status <u>Final</u>	Results		Biological Reference Interv	al Units
PPBS(POST PRANDIAL BLOOD SUGAR)	108		Normal: < 140, Impaired Glucose Tolerance:14 199 Diabetic > or = 200	mg/dL 40-
LIPID PROFILE, SERUM				
CHOLESTEROL, TOTAL	263	High	Desirable: <200 BorderlineHigh : 200-239 High : > or = 240	mg/dL
TRIGLYCERIDES	153	High	Desirable: < 150 Borderline High: 150 - 199 High: 200 - 499 Very High: > or = 500	mg/dL
HDL CHOLESTEROL	43		< 40 Low > or = 60 High	mg/dL
CHOLESTEROL LDL	189	High	Adult levels: Optimal < 100 Near optimal/above optimal: 1 129 Borderline high: 130-159 High: 160-189 Very high: = 190	mg/dL L00-
NON HDL CHOLESTEROL	220	High	Desirable: Less than 130 Above Desirable: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very high: > or = 220	mg/dL
CHOL/HDL RATIO	6.1		, J	
LDL/HDL RATIO	-1.7	Low	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN	30.6		2 or ornight work	mg/dL
LIVER FUNCTION PROFILE, SERUM				
BILIRUBIN, TOTAL	0.24		0.0 - 1.2	mg/dL
BILIRUBIN, DIRECT	0.11		0.0 - 0.2	mg/dL
BILIRUBIN, INDIRECT	0.13		0.00 - 1.00	mg/dL
TOTAL PROTEIN	7.9		6.4 - 8.3	g/dL
ALBUMIN	5.0		3.50 - 5.20	g/dL
GLOBULIN	2.9		2.0 - 4.1	g/dL
ALBUMIN/GLOBULIN RATIO	1.7		1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	17		UPTO 40	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT)	18		UP TO 45	U/L
ALKALINE PHOSPHATASE	65		40 - 129	U/L









CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Opposite St Raphael's Higher Secondary School , Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA - CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units	
Tillar	Results	Diological Reference Interve	011115
GAMMA GLUTAMYL TRANSFERASE (GGT)	27	8 - 61	U/L
LACTATE DEHYDROGENASE	190	135 - 225	U/L
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	12	6 - 20	mg/dL
CREATININE, SERUM			
CREATININE	0.69 Lov	v 0.70 - 1.20	mg/dL
BUN/CREAT RATIO			
BUN/CREAT RATIO	17.39 Hig	h 5.0 - 15.0	
URIC ACID, SERUM			
URIC ACID	7.1	3.5 - 7.2	mg/dL
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	7.9	6.4 - 8.3	g/dL
ALBUMIN, SERUM			
ALBUMIN	5.0	3.5 - 5.2	g/dL
GLOBULIN			
GLOBULIN	2.9	2.0 - 4.1	g/dL
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM, SERUM	143.6	136.0 - 146.0	mmol/L
POTASSIUM, SERUM	5.47 Hig	h 3.50 - 5.10	mmol/L
CHLORIDE, SERUM	104.1	98.0 - 106.0	mmol/L
PHYSICAL EXAMINATION, URINE			
COLOR	PALE YELLOW		
APPEARANCE	CLEAR		
CHEMICAL EXAMINATION, URINE			
PH	5.5	4.7 - 7.5	
SPECIFIC GRAVITY	1.015	1.003 - 1.035	
PROTEIN	NOT DETECTED	NOT DETECTED	
GLUCOSE	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BLOOD	NOT DETECTED	NOT DETECTED	
BILIRUBIN	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
NITRITE	NOT DETECTED	NOT DETECTED	
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED	









CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA

8800465156

Opposite St Raphael's Higher Secondary School , Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

RECEIVED: 10/12/2022 10:49 13/12/2022 13:57 DRAWN: REPORTED:

CLIENT PATIENT ID: REFERRING DOCTOR: DR. BANK OF BARODA -

Test Report Status <u>Final</u>	Results	Biological Reference Interva	l Units
MICROSCOPIC EXAMINATION, URINE			
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	2-3	0-5	/HPF
EPITHELIAL CELLS	2-3	0-5	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	
REMARKS	Please note that all the urin	nary findings are confirmed manu	ually as well.
THYROID PANEL, SERUM			
ТЗ	99.10	80.00 - 200.00	ng/dL
T4	6.59	5.10 - 14.10	μg/dL
TSH (ULTRASENSITIVE)	1.200	0.270 - 4.200	μIU/mL









CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: 0290VL001162 AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA - CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Biological Reference Interval Units

Interpretation(s)

Triiodothyronine T3, **Thyroxine T4**, and **Thyroid Stimulating Hormone TSH** are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. owidctlparowidctlparBelow mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hypothyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE A
RH TYPE POSITIVE

XRAY-CHEST

»» BOTH THE LUNG FIELDS ARE CLEAR

»» BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR





Scan to View Report





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA - CLIENT PATIENT ID:

Test Report Status Final Results Biological Reference Interval Units

»» BOTH THE HILA ARE NORMAL

»»CARDIAC AND AORTIC SHADOWS APPEAR NORMAL»»BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL

»» VISUALIZED BONY THORAX IS NORMAL

IMPRESSION NO ABNORMALITY DETECTED

TMT OR ECHO

TMT OR ECHO TMT TEST IS NEGATIVE

ECG

ECG SINUS RHYTHM, NORMAL ECG

MEDICAL HISTORY

RELEVANT PRESENT HISTORY

RELEVANT PAST HISTORY

RELEVANT PERSONAL HISTORY

RELEVANT FAMILY HISTORY

OCCUPATIONAL HISTORY

HISTORY OF MEDICATIONS

NOT SIGNIFICANT

NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.60 mts WEIGHT IN KGS. 67 Kgs

BMI 8 Weight Status as follows: kg/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

NORMAL MENTAL / EMOTIONAL STATE PHYSICAL ATTITUDE NORMAL GENERAL APPEARANCE / NUTRITIONAL STATUS **OVERWEIGHT BUILT / SKELETAL FRAMEWORK AVERAGE** FACIAL APPEARANCE **NORMAL NORMAL** SKIN UPPER LIMB NORMAL LOWER LIMB **NORMAL NECK** NORMAL

NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER

THYROID GLAND NOT ENLARGED



Page 6 Of 12





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA - CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Biological Reference Interval Units

CAROTID PULSATION NORMAL TEMPERATURE AFEBRILE

PULSE 61/MIN, REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

BRUIT

RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM

BP 130/80 MM HG mm/Hg

(SITTING) NORMAL

APEX BEAT NORMAL HEART SOUNDS NORMAL MURMURS ABSENT

RESPIRATORY SYSTEM

SIZE AND SHAPE OF CHEST

MOVEMENTS OF CHEST

BREATH SOUNDS INTENSITY

NORMAL

NORMAL

BREATH SOUNDS QUALITY VESICULAR (NORMAL)

ADDED SOUNDS ABSENT

PER ABDOMEN

PERICARDIUM

APPEARANCE NORMAL VENOUS PROMINENCE ABSENT

LIVER NOT PALPABLE
SPLEEN NOT PALPABLE

HERNIA ABSENT

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS NORMAL
CRANIAL NERVES NORMAL
CEREBELLAR FUNCTIONS NORMAL
SENSORY SYSTEM NORMAL
MOTOR SYSTEM NORMAL
REFLEXES NORMAL

MUSCULOSKELETAL SYSTEM

SPINE NORMAL JOINTS NORMAL

BASIC EYE EXAMINATION



Page 7 Of 12





CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: 0290VL001162 AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA -CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

CONJUNCTIVA NORMAL **EYELIDS** NORMAL EYE MOVEMENTS NORMAL CORNEA **NORMAL**

DISTANT VISION RIGHT EYE WITHOUT GLASSES 6/6, WITHIN NORMAL LIMIT DISTANT VISION LEFT EYE WITHOUT GLASSES 6/6, WITHIN NORMAL LIMIT NEAR VISION RIGHT EYE WITHOUT GLASSES N/6, WITHIN NORMAL LIMIT NEAR VISION LEFT EYE WITHOUT GLASSES N/6, WITHIN NORMAL LIMIT COLOUR VISION **NORMAL**

BASIC ENT EXAMINATION

EXTERNAL EAR CANAL **NORMAL** TYMPANIC MEMBRANE NORMAL

NOSE NO ABNORMALITY DETECTED

SINUSES NORMAL

THROAT NO ABNORMALITY DETECTED

TONSILS NOT ENLARGED

SUMMARY

RELEVANT HISTORY NOT SIGNIFICANT RELEVANT GP EXAMINATION FINDINGS **OVERWEIGHT**

REMARKS / RECOMMENDATIONS CLINICAL FINDINGS- OVERWEIGHT STATUS SLIGHTLY DYSLIPIDEMIA

ADVICE- LOW FAT DIET AND REGULAR PHYSICAL EXERCISE FOR

OVERWEIGHT STATUS AND DYSLIPIDEMIA - NEED PHYSICIAN CONSULTATION FOR PROPER EVALUATION

FITNESS STATUS

FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS) FITNESS STATUS

Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13)

from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.4 (20.1%) covid-19 patients with mild disease might become severe. 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.
ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION:



Page 8 Of 12 Scan to View Report

Scan to View Details





CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHT **NEW DELHI 110030 DELHI INDIA** 8800465156

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

0290VL001162 AGE: 35 Years SEX: Male ACCESSION NO: ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA -CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change. **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy,

Estrogen medication, Aging.
Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, salicylates)

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition.

GLUCOSE FASTING,FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in

Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency, hypopituitarism,diffuse liver disease, malignancy (adrenocortical, stomach,fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia),Drugs- insulin, ethanol, propranolol sulfonylureas,tolbutamide, and other oral hypoglycemic agents

NOTE:

Hypoglycemia is defined as a glucoseof < 50 mg/dL in men and < 40 mg/dL in women.

While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2.Diagnosing diabetes.
- 3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range. 1.eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.

- 2. eAG gives an evaluation of blood glucose levels for the last couple of months. 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :

I.Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic

anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

III. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results. IV.Interference of hemoglobinopathies in HbA1c estimation is seen in

a. Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c. b.Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c.HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give

yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin



Page 9 Of 12 Scan to View Report





CLIENT CODE: C000138355

CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

SRL LTD

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

0290VL001162 AGE: 35 Years ACCESSION NO: SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA -CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal tevels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Mephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)
Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
 Loss of body fluid (dehydration)
 Muscle problems, such as breakdown of muscle fibers

- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis
- Muscular dystrophy

URIC ACID, ŚERUM-Causes of Increased levels: -Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome

Causes of decreased levels-Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM-Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom""""""""" disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB. Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for

The test is performed by both forward as well as reverse grouping methods.

HISTORY-** THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

FITNESS STATUS-

availability of the same.

Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for .These are then further correlated with details of the job under consideration to eventually fit the right man to the right job.

- Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:
 Fit (As per requested panel of tests) SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the
- specific test panel requested for. • Fit (with medical advice) (As per requested panel of tests) - This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been





Scan to View Details





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030

NEW DELHI 11003 DELHI INDIA 8800465156 SRL LTD

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA - CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Biological Reference Interval Units

detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician's consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.

consultation and counseling in order to bring back to normal the mildly deranged parameters. For all purposes the individual is FIT to join the job.
• Fitness on Hold (Temporary Unfit) (As per requested panel of tests) - Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.

• Unfit (As per requested panel of tests) - An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.







CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Opposite St Raphael's Higher Secondary School, Old Seshore Road,

Residency Area INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: SHARAD THEKEDA PATIENT ID: SHARM020187290

ACCESSION NO: **0290VL001162** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 10/12/2022 10:49 REPORTED: 13/12/2022 13:57

REFERRING DOCTOR: DR. BANK OF BARODA - CLIENT PATIENT ID:

Test Report Status Final Results Units

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ULTRASOUND ABDOMEN
ULTRASOUND ABDOMEN

NO ABNORMALITIES DETECTED

End Of Report
Please visit www.srlworld.com for related Test Information for this accession

Proite

Dr.Arpita Pasari, MD Consultant Pathologist

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062





Scan to View Report