



SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

RECEIVED: 25/03/2023 09:25 03/04/2023 12:53 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ΒI	LOOD	COUNT	S,EDTA	WHOLE	BLOOD
----	------	-------	--------	-------	-------

HEMOGLOBIN (HB)	11.9	Low	12.0 - 15.0	g/dL
METHOD: SLS- HEMOGLOBIN DETECTION METHOD				
RED BLOOD CELL (RBC) COUNT	4.66		3.8 - 4.8	mil/µL
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION				
WHITE BLOOD CELL (WBC) COUNT	6.90		4.0 - 10.0	thou/µL
METHOD: FLUORESCENCE FLOW CYTOMETRY				
PLATELET COUNT	447	High	150 - 410	thou/µL
METHOD: HYDRODYNAMIC FOCUSING BY DC DETECTION				
RBC AND PLATELET INDICES				
HEMATOCRIT (PCV)	39.0		36.0 - 46.0	%
METHOD : CUMULATIVE PULSE HEIGHT DETECTION METHOD				
MEAN CORPUSCULAR VOLUME (MCV)	83.7		83.0 - 101.0	fL
METHOD: CALCULATED FROM RBC & HCT				
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	25.5	Low	27.0 - 32.0	pg
METHOD: CALCULATED FROM THE RBC & HGB				
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) METHOD: CALCULATED FROM THE HGB & HCT	30.5	Low	31.5 - 34.5	g/dL
RED CELL DISTRIBUTION WIDTH (RDW)	13.2		11.6 - 14.0	%
METHOD: CALCULATED FROM RBC SIZE DISTRIBUTION CURVE				
MENTZER INDEX	18.0			
MEAN PLATELET VOLUME (MPV)	11.2	High	6.8 - 10.9	fL
METHOD: CALCULATED FROM PLATELET COUNT & PLATELET HEMATO	CRIT			
WBC DIFFERENTIAL COUNT				
NEUTROPHILS	51		40 - 80	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
LYMPHOCYTES	41	High	20 - 40	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
MONOCYTES	6		2 - 10	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
EOSINOPHILS	2		1 - 6	%
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE NEUTROPHIL COUNT	3.52		2.0 - 7.0	thou/µL
METHOD: FLOW CYTOMETRY WITH LIGHT SCATTERING				



Page 1 Of 12 Scan to View Report





SRL Ltd S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results	Biological Reference Inter	rval Units
ABSOLUTE LYMPHOCYT	E COUNT	2.86	1.0 - 3.0	thou/µL
METHOD : FLOW CYTOMETRY	WITH LIGHT SCATTERING			
ABSOLUTE MONOCYTE METHOD: FLOW CYTOMETRY		0.41	0.2 - 1.0	thou/μL
ABSOLUTE EOSINOPHI METHOD: FLOW CYTOMETRY		0.14	0.02 - 0.50	thou/µL
NEUTROPHIL LYMPHOC		1.2		
MORPHOLOGY	- ()			
RBC		NORMOCYTIC NOR	MOCHROMIC	
WBC		NORMAL MORPHOL		
METHOD : MICROSCOPIC EX	ΔΜΙΝΑΤΙΟΝ	NORMAL MORETION	.001	
PLATELETS		INCREASED		
	MENTATION RATE (ES			
E.S.R		14	< 20	mm at 1 hr
METHOD : MODIFIED WESTE	RGREN			
GLUCOSE FASTING,F	LUORIDE PLASMA			
FBS (FASTING BLOOD	SUGAR)	81	Normal 75 - 99 Pre-diabetics: 100 - 125 Diabetic: > or = 126	mg/dL
METHOD : ENZYMATIC REFER	RENCE METHOD WITH HEXOKINA	ASE		
GLYCOSYLATED HEM BLOOD	OGLOBIN(HBA1C), ED	OTA WHOLE		
НВА1С		5.4	Non-diabetic Adult < 5.7 Pre-diabetes 5.7 - 6.4 Diabetes diagnosis: > or = 6 Therapeutic goals: < 7.0 Action suggested : > 8.0 (ADA Guideline 2021)	% 5.5
METHOD: HPLC				
ESTIMATED AVERAGE (` ,	108.3	< 116.0	mg/dL
METHOD : CALCULATED PAR				
GLUCOSE, POST-PRA				
PPBS(POST PRANDIAL METHOD : ENZYMATIC REFER	BLOOD SUGAR) RENCE METHOD WITH HEXOKINA	75 ASE	70 - 139	mg/dL

LIPID PROFILE, SERUM



Page 2 Of 12





SRL Ltd S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results		Biological Reference Interva	l Units
CHOLESTEROL, TOTAL	220	High	Desirable cholesterol level < 200 Borderline high cholesterol 200 - 239 High cholesterol > / = 240	mg/dL
METHOD: ENZYMATIC COLORIMETRIC ASSAY TRIGLYCERIDES	90		Normal: < 150 Borderline high: 150 - 199 High: 200 - 499 Very High: >/= 500	mg/dL
METHOD : ENZYMATIC COLORIMETRIC ASSAY HDL CHOLESTEROL	38	Low	Low HDL Cholesterol <40	mg/dL
METHOD : ENZYMATIC, COLORIMETRIC CHOLESTEROL LDL	164	High	High HDL Cholesterol >/= 60 Adult levels: Optimal < 100	mg/dL
METHOD: ENZYMATIC COLORIMETRIC ASSAY			Near optimal/above optimal: 10 129 Borderline high: 130-159 High: 160-189 Very high: = 190	00-
NON HDL CHOLESTEROL	182	High	Desirable : < 130 Above Desirable : 130 -159 Borderline High : 160 - 189 High : 190 - 219 Very high : > / = 220	mg/dL
VERY LOW DENSITY LIPOPROTEIN	18.0		< OR = 30.0	mg/dL
CHOL/HDL RATIO	5.8	High	Low Risk: 3.3 - 4.4 Average Risk: 4.5 - 7.0 Moderate Risk: 7.1 - 11.0 High Risk: > 11.0	
LDL/HDL RATIO	4.3	High	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate R >6.0 High Risk	Risk
LIVER FUNCTION PROFILE, SERUM				
BILIRUBIN, TOTAL METHOD: COLORIMETRIC DIAZO	0.64		Upto 1.2	mg/dL
BILIRUBIN, DIRECT	0.27		< 0.30	mg/dL
BILIRUBIN, INDIRECT	0.37		0.1 - 1.0	mg/dL
TOTAL PROTEIN METHOD: COLORIMETRIC	7.2		6.0 - 8.0	g/dL



Page 3 Of 12





SRL Ltd S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
ALBUMIN	4.1	3.97 - 4.94	a/dl
METHOD : COLORIMETRIC	4.1	3.97 - 4.94	g/dL
GLOBULIN	3.1	2.0 - 3.5	g/dL
ALBUMIN/GLOBULIN RATIO	1.3	1.0 - 2.1	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	19	< OR = 35	U/L
METHOD : UV ABSORBANCE	19	1 OK = 33	0, 2
ALANINE AMINOTRANSFERASE (ALT/SGPT)	20	< OR = 35	U/L
METHOD : UV ABSORBANCE			-,
ALKALINE PHOSPHATASE	61	35 - 104	U/L
METHOD: COLORIMETRIC			
GAMMA GLUTAMYL TRANSFERASE (GGT)	23	0 - 40	U/L
METHOD: ENZYMATIC, COLORIMETRIC			
LACTATE DEHYDROGENASE	211	125 - 220	U/L
METHOD: UV ABSORBANCE			
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	7	6 - 20	mg/dL
METHOD: ENZYMATIC ASSAY			
CREATININE, SERUM			
CREATININE	0.66	0.5 - 0.9	mg/dL
METHOD: COLORIMETRIC			
BUN/CREAT RATIO			
BUN/CREAT RATIO	10.61	8.0 - 15.0	
URIC ACID, SERUM			
URIC ACID	3.5	2.4 - 5.7	mg/dL
METHOD: ENZYMATIC COLORIMETRIC ASSAY			
TOTAL PROTEIN, SERUM			
TOTAL PROTEIN	7.2	6.0 - 8.0	g/dL
METHOD : COLORIMETRIC			
ALBUMIN, SERUM			
ALBUMIN	4.1	3.97 - 4.94	g/dL
METHOD : COLORIMETRIC			
GLOBULIN		2.2.5	
GLOBULIN	3.1	2.0 - 3.5	g/dL
ELECTROLYTES (NA/K/CL), SERUM			
SODIUM, SERUM	138	136 - 145	mmol/L
POTASSIUM, SERUM	4.29	3.5 - 5.1	mmol/L



Page 4 Of 12





SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

RECEIVED: 25/03/2023 09:25 03/04/2023 12:53 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
CHLORIDE, SERUM	103	98 - 107	mmol/L
PHYSICAL EXAMINATION, URINE	103	96 - 107	IIIIIOI/ L
COLOR	PALE YELLOW		
APPEARANCE	CLEAR		
CHEMICAL EXAMINATION, URINE	CLLAIN		
PH	6.0	5.00 - 7.50	
SPECIFIC GRAVITY	1.010	1.010 - 1.030	
METHOD: URINE ROUTINE & MICROSCOPY EXAMINATION BY		1.010 1.030	
PROTEIN	NOT DETECTED	NOT DETECTED	
GLUCOSE	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BLOOD	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
NITRITE	NOT DETECTED	NOT DETECTED	
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, URINE			
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	2-3	0-5	, /HPF
EPITHELIAL CELLS	2-3	0-5	, /HPF
CASTS	NOT DETECTED		•
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	
METHOD: URINE ROUTINE & MICROSCOPY EXAMINATION BY	INTEGRATED AUTOMATED SYSTEM		
THYROID PANEL, SERUM			
ТЗ	139.0	Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester:105.0 - 230.0 2nd Trimester:129.0 - 262.0 3rd Trimester:135.0 - 262.0	ng/dL
METHOD: ELECTROCHEMILUMINESCENCE			
T4	8.28	Non-Pregnant Women 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70	μg/dL



Page 5 Of 12

μIU/mL





CLIENT CODE: C000138394 CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

0.519

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u> METHOD: ELECTROCHEMILUMINESCENCE

> Non Pregnant Women 0.27 - 4.20Pregnant Women

> > 1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15

METHOD: ELECTROCHEMILUMINESCENCE

PAPANICOLAOU SMEAR

TSH (ULTRASENSITIVE)

TEST METHOD CONVENTIONAL GYNEC CYTOLOGY

METHOD: MICROSCOPIC EXAMINATION

SPECIMEN TYPE P-458/23

TWO UNSTAINED CERVICAL SMEARS RECEIVED

METHOD: MICROSCOPIC EXAMINATION

REPORTING SYSTEM 2014 BETHESDA SYSTEM FOR REPORTING CERVICAL CYTOLOGY

SATISFACTORY SPECIMEN ADEQUACY

METHOD: PAP STAIN & MICROSCOPIC EXAMINATION

THE SMEARS SHOW FEW SUPERFICIAL SQUAMOUS CELLS, FEW **MICROSCOPY**

INTERMEDIATE SQUAMOUS CELLS, OCCASIONAL SQUAMOUS

METAPLASTIC CELLS, AND FEW CLUSTERS OF ENDOCERVICAL CELLS IN

THE BACKGROUND OF FEW POLYMORPHS.

METHOD: PAP STAIN NEGATIVE FOR INTRAEPITHELIAL LESION OR MALIGNANCY INTERPRETATION / RESULT

METHOD: PAP STAIN & MICROSCOPIC EXAMINATION

Comments

PLEASE NOTE PAPANICOLAU SMEAR STUDY IS A SCREENING PROCEDURE FOR CERVICAL CANCER WITH INHERENT FALSE NEGATIVE RESULTS HENCE SHOULD BE INTERPRETED WITH CAUTION. NO CYTOLOGICAL EVIDENCE OF HPV INFECTION IN THE SMEARS STUDIED. SMEARS WILL BE PRESERVED FOR 5 YEARS ONLY.

MICROSCOPIC EXAMINATION, STOOL

SAMPLE NOT RECEIVED REMARK

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE B

METHOD: GEL COLUMN AGGLUTINATION METHOD.

RH TYPE **POSITIVE**

METHOD: GEL COLUMN AGGLUTINATION METHOD.

XRAY-CHEST

IMPRESSION NO ABNORMALITY DETECTED

TMT OR ECHO



Page 6 Of 12

Scan to View Report





SRL Ltd S.K. Tower,Hari Niwas, LBS Marg THANE, 400602

MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Biological Reference Interval Units

TMT OR ECHO NEGATIVE

ECG

ECG WITHIN NORMAL LIMITS

MEDICAL HISTORY

RELEVANT PRESENT HISTORY NOT SIGNIFICANT RELEVANT PAST HISTORY NOT SIGNIFICANT

RELEVANT PERSONAL HISTORY MARRIED / MIXED DIET / NO ALLERGIES / NO SMOKING / NO

ALCOHOL.

LMP (FOR FEMALES) 18/3/2023

RELEVANT FAMILY HISTORY NOT SIGNIFICANT HISTORY OF MEDICATIONS NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS1.49mtsWEIGHT IN KGS.54Kgs

BMI & Weight Status as follows: kg/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

NORMAL MENTAL / EMOTIONAL STATE PHYSICAL ATTITUDE NORMAL GENERAL APPEARANCE / NUTRITIONAL STATUS **HEALTHY BUILT / SKELETAL FRAMEWORK AVERAGE** FACIAL APPEARANCE **NORMAL** SKIN NORMAL UPPER LIMB **NORMAL** LOWER LIMB **NORMAL NECK NORMAL**

NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER

THYROID GLAND NOT ENLARGED CAROTID PULSATION NORMAL

TEMPERATURE NORMAL

PULSE REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID BRUIT

RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM



View Details Scan to View Report

Page 7 Of 12





SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

RECEIVED: 25/03/2023 09:25 03/04/2023 12:53 DRAWN: REPORTED:

REFERRING DOCTOR: SELF	CLIENT PATIENT ID:		
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units	
ВР	120/80 MM HG (SUPINE)	mm/Hg	
PERICARDIUM	NORMAL		
APEX BEAT	NORMAL		
HEART SOUNDS	NORMAL		
MURMURS	ABSENT		
RESPIRATORY SYSTEM			
SIZE AND SHAPE OF CHEST	NORMAL		
MOVEMENTS OF CHEST	SYMMETRICAL		
BREATH SOUNDS INTENSITY	NORMAL		
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)		
ADDED SOUNDS	ABSENT		
PER ABDOMEN			
APPEARANCE	NORMAL		
VENOUS PROMINENCE	ABOENT		

VENOUS PROMINENCE **ABSENT**

LIVER **NOT PALPABLE SPLEEN** NOT PALPABLE

HERNIA ABSENT

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS **NORMAL** CRANIAL NERVES **NORMAL** CEREBELLAR FUNCTIONS **NORMAL** SENSORY SYSTEM **NORMAL** MOTOR SYSTEM NORMAL **REFLEXES NORMAL**

MUSCULOSKELETAL SYSTEM

SPINE NORMAL JOINTS NORMAL

BASIC EYE EXAMINATION

CONJUNCTIVA NORMAL EYELIDS NORMAL EYE MOVEMENTS **NORMAL CORNEA NORMAL**

WITHIN NORMAL LIMIT DISTANT VISION RIGHT EYE WITHOUT GLASSES



Page 8 Of 12

Scan to View Report





SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: 0181WC001638 AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

DISTANT VISION LEFT EYE WITHOUT GLASSES WITHIN NORMAL LIMIT NEAR VISION RIGHT EYE WITHOUT GLASSES WITHIN NORMAL LIMIT NEAR VISION LEFT EYE WITHOUT GLASSES WITHIN NORMAL LIMIT COLOUR VISION NORMAL

SUMMARY

RELEVANT HISTORY NOT SIGNIFICANT RELEVANT GP EXAMINATION FINDINGS NOT SIGNIFICANT

REMARKS / RECOMMENDATIONS LOW FAT, LOW CARBOHYDRATE, HIGH FIBRE DIET.

REGULAR EXERCISE.REGULAR WALK FOR 30-40 MIN DAILY. REPEAT B.SUGAR, LIPID PROFILE AFTER 3 MONTHS OF DIET AND

EXERCISE.

Interpretation(s)
BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive

patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION:
Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy,

Estioger interaction, raging. Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum. Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Polkilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine,

salicylates)

REFERENCE:

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLUCOSE FASTING,FLUORIDE PLASMA-**TEST DESCRIPTION**

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the



Page 9 Of 12

Scan to View Details

Scan to View Report





SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATTENT ID: RUPAF110696181

ACCESSION NO: 0181WC001638 26 Years AGE: SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

Increased in:Diabetes mellitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides. Decreased in :Pancreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease, malignancy(adrenocortical,stomach,fibrosarcoma),infant of a diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol

sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values),there is wide fluctuation within

High fasting glucose levels correlate with higher glucose infoliationing results (weekly mean capillary glucose values), there is wide nuctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- Diagnosing diabetes.
 Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- 1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
- 2. eAG gives an evaluation of blood glucose levels for the last couple of months. 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

HbA1c Estimation can get affected due to :

- 1. Shortened Erythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss,hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.
- 2.Vitamin C & E are reported to falsely lower test results.(possibly by inhibiting glycation of hemoglobin.
- 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.
- 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
- b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
 c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin whe there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver,liver cancer,kidney failure,hemolytic anemia,pancreatitis,hemochromatosis. AST levels may also increase after a heart attack or strenuous activity.ALT test measures the amount of this enzyme in the blood.ALT is found mainly in the liver, but also in smaller amounts in the kidneys,heart,muscles, and pancreas.It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health.AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.

Total Protein also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and

globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular

permeability or decreased lymphatic clearance,malnutrition and wasting etc
BLOOD UREA NITROGEN (BUN), SERUM-**Causes of Increased** levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.
CREATININE, SERUM-Higher than normal level may be due to:

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such



Page 10 Of 12

Scan to View Report





SRL Ltd S.K. Tower, Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units <u>Final</u>

as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to: Myasthenia Gravis, Muscuophy
URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake,Prolonged Fasting,Rapid weight loss),Gout,Lesch nyhan syndrome,Type 2 DM,Metabolic

syndrome **Causes of decreased levels**-Low Zinc intake,OCP,Multiple Sclerosis
TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum.Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome,Protein-losing enteropathy etc.

ALBUMIN, SERUMHuman serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. **Low** blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface

of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.





Scan to View Report





SRL Ltd S.K. Tower,Hari Niwas, LBS Marg THANE, 400602 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

Email: customercare.thane@srl.in

PATIENT NAME: RUPALI LOHAKARE PATIENT ID: RUPAF110696181

ACCESSION NO: **0181WC001638** AGE: 26 Years SEX: Female

DRAWN: RECEIVED: 25/03/2023 09:25 REPORTED: 03/04/2023 12:53

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Units

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ULTRASOUND ABDOMEN
ULTRASOUND ABDOMEN
NO ABNORMALITIES DETECTED

End Of Report
Please visit www.srlworld.com for related Test Information for this accession

CONDITIONS OF LABORATORY TESTING & REPORTING

- $1. \ It is presumed that the test sample belongs to the patient named or identified in the test requisition form.$
- 2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062



Scan to View Details

Page 12 Of 12