



CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI **NEW DELHI 110030** DELHI INDIA 8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: 0290WD00043 AGE: 35 Years SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 03/04/2023 16:46 DRAWN: REPORTED:

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

BLOOD COUNTS,EDTA WHOLE BLOOD			
HEMOGLOBIN (HB)	15.5	13.0 - 17.0	g/dL
METHOD: SPECTROPHOTOMETRY			
RED BLOOD CELL (RBC) COUNT	5.56	High 4.5 - 5.5	mil/μL
METHOD: ELECTRICAL IMPEDANCE			
WHITE BLOOD CELL (WBC) COUNT	7.00	4.0 - 10.0	thou/µL
METHOD: ELECTRICAL IMPEDANCE			
PLATELET COUNT	389	150 - 410	thou/µL
METHOD : ELECTRICAL IMPEDANCE			
RBC AND PLATELET INDICES			
HEMATOCRIT (PCV)	46.7	40 - 50	%
METHOD: CALCULATED			
MEAN CORPUSCULAR VOLUME (MCV)	84.0	83 - 101	fL
METHOD: CALCULATED			
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	27.9	27.0 - 32.0	pg
METHOD : CALCULATED			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	33.3	31.5 - 34.5	g/dL
METHOD : CALCULATED			
RED CELL DISTRIBUTION WIDTH (RDW)	12.7	11.6 - 14.0	%
METHOD: CALCULATED			
MENTZER INDEX	15.1		
MEAN PLATELET VOLUME (MPV)	8.0	6.8 - 10.9	fL
METHOD: CALCULATED			
WBC DIFFERENTIAL COUNT			
NEUTROPHILS	56	40 - 80	%
METHOD: IMPEDANCE / MICROSCOPY			
LYMPHOCYTES	38	20 - 40	%
METHOD : IMPEDANCE / MICROSCOPY			
MONOCYTES	04	2 - 10	%
METHOD : IMPEDANCE / MICROSCOPY			
EOSINOPHILS	02	1 - 6	%
METHOD: IMPEDANCE / MICROSCOPY			
BASOPHILS	00	0 - 2	%
METHOD: IMPEDANCE / MICROSCOPY			

3.92

2.0 - 7.0



ABSOLUTE NEUTROPHIL COUNT

Page 1 Of 18

thou/µL





CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA

8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: 0290WD00043 AGE: 35 Years SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 03/04/2023 16:46 DRAWN: REPORTED:

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results		Biological Reference Interva	l Units
METHOD : CALCULATED					
ABSOLUTE LYMPHOCYTE METHOD: CALCULATED	COUNT	2.66		1.0 - 3.0	thou/µL
ABSOLUTE MONOCYTE (COUNT	0.28		0.2 - 1.0	thou/µL
ABSOLUTE EOSINOPHIL METHOD : CALCULATED	. COUNT	0.14		0.02 - 0.50	thou/µL
ERYTHROCYTE SEDIM	IENTATION RATE ((ESR),WHOLE			
E.S.R METHOD : MODIFIED WESTER		05		0 - 14	mm at 1 hr
GLUCOSE FASTING,FL					
FBS (FASTING BLOOD S	SUGAR)	89		74 - 99	mg/dL
METHOD: HEXOKINASE GLYCOSYLATED HEMO BLOOD	OGLOBIN(HBA1C),	EDTA WHOLE			
HBA1C		5.7		Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 Therapeutic goals: < 7.0 Action suggested: > 8.0 (ADA Guideline 2021)	%
METHOD : HPLC TECHNOLOGY		116.0	Wich	. 110.0	/ dl
ESTIMATED AVERAGE G	` ,	116.9	nigii	< 116.0	mg/dL
PPBS(POST PRANDIAL E		114		Normal: < 140, Impaired Glucose Tolerance:140 199 Diabetic > or = 200	mg/dL)-
METHOD: HEXOKINASE					
LIPID PROFILE, SERU	IM				
CHOLESTEROL, TOTAL		202	High	Desirable: <200 BorderlineHigh: 200-239 High: > or = 240	mg/dL
METHOD : OXIDASE, ESTERAS	SE, PEROXIDASE			111g11 : > 01 = 240	
TRIGLYCERIDES		187	High	Desirable: < 150 Borderline High: 150 - 199 High: 200 - 499 Very High: > or = 500	mg/dL
METHOD : ENZYMATIC ASSAY	,	46		4.40 Law	/ dl
HDL CHOLESTEROL METHOD : DIRECT- NON IMM	UNOLOGICAL	46		< 40 Low > or = 60 High	mg/dL











CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: 0290WD00043 AGE: 35 Years $\mathsf{SEX}:\ \mathsf{Male}$ ABHA NO:

RECEIVED: 03/04/2023 08:28 03/04/2023 16:46 DRAWN: REPORTED:

CLIENT PATIENT ID: REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

Test Report Status <u>Final</u>	Results		Biological Reference Interv	al Units
CHOLESTEROL LDL	119	High	Adult levels: Optimal < 100 Near optimal/above optimal: 129 Borderline high: 130-159 High: 160-189 Very high: = 190	mg/dL 100-
NON HDL CHOLESTEROL	156	High	, ,	mg/dL
METHOD: CALCULATED			, 3	
VERY LOW DENSITY LIPOPROTEIN METHOD: CALCULATED	37.4			mg/dL
CHOL/HDL RATIO	4.4			
LDL/HDL RATIO	2.6		0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate >6.0 High Risk	Risk









CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA**

8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

0290WD00043 AGE: 35 Years ACCESSION NO: SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 DRAWN: REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units <u>Final</u>

Interpretation(s)

- 1) Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.
- 2) Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.
- 3)HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL
- 4) LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.
- 5)Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category					
Extreme risk group	A.CAD with > 1 feature of high risk group				
	B. CAD with > 1 feature of Very high risk g	group or recurrent ACS (within 1 year) despite LDL-C			
	< or = 50 mg/dl or polyvascular disease				
Very High Risk	1. Established ASCVD 2. Diabetes with 2 1	major risk factors or evidence of end organ damage 3.			
	Familial Homozygous Hypercholesterolemi	a			
High Risk		abetes with 1 major risk factor or no evidence of end			
	organ damage. 3. CKD stage 3B or 4. 4. LDL > 190 mg/dl 5. Extreme of a single risk factor. 6.				
	Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid				
	plaque				
Moderate Risk	2 major ASCVD risk factors				
Low Risk	0-1 major ASCVD risk factors				
Major ASCVD (Ath	Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors				
1. Age > or = 45 years in males and > or = 55 years in females 3. Current Cigarette smoking or tobacco use					
2. Family history of p	remature ASCVD	4. High blood pressure			
5. Low HDL					

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
Category A	$\langle OR = 30 \rangle$	$\langle OR = 60 \rangle$		





Scan to View Details





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030

DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND

PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results	Biological Reference Interval	Units
-				

Extreme Risk Group	< OR = 30	< OR = 60	> 30	>60
Category B				
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

^{*}After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.46	0.0 - 1.2	mg/dL
METHOD : JENDRASSIK AND GROFF			
BILIRUBIN, DIRECT	0.14	0.0 - 0.2	mg/dL
METHOD : DIAZOTIZATION			
BILIRUBIN, INDIRECT	0.32	0.00 - 1.00	mg/dL
METHOD: CALCULATED			
TOTAL PROTEIN	7.8	6.4 - 8.3	g/dL
METHOD : BIURET			
ALBUMIN	4.8	3.50 - 5.20	g/dL
METHOD: BROMOCRESOL GREEN			
GLOBULIN	3.0	2.0 - 4.1	g/dL
METHOD: CALCULATED			
ALBUMIN/GLOBULIN RATIO	1.6	1.0 - 2.0	RATIO
METHOD: CALCULATED			
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	27	UPTO 40	U/L
METHOD : UV WITH P5P			
ALANINE AMINOTRANSFERASE (ALT/SGPT)	45	UP TO 45	U/L
METHOD : UV WITH P5P			
ALKALINE PHOSPHATASE	105	40 - 129	U/L
METHOD: PNPP			
GAMMA GLUTAMYL TRANSFERASE (GGT)	32	8 - 61	U/L
METHOD: G-GLUTAMYL-CARBOXY-NITROANILIDE			
LACTATE DEHYDROGENASE	186	135 - 225	U/L
METHOD : ENZYMATIC LACTATE - PYRUVATE(IFCC)			
BLOOD UREA NITROGEN (BUN), SERUM			
BLOOD UREA NITROGEN	6	6 - 20	mg/dL

METHOD : UREASE KINETIC

CREATININE, SERUM



Page 5 Of 18







CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units			
CREATININE	0.90	0.70 - 1.20	mg/dL			
METHOD: ALKALINE PICRATE KINETIC JAFFES						
BUN/CREAT RATIO						
BUN/CREAT RATIO	6.67	5.0 - 15.0				
METHOD: CALCULATED						
URIC ACID, SERUM						
URIC ACID	5.5	3.5 - 7.2	mg/dL			
METHOD: URICASE/CATALASE UV						
TOTAL PROTEIN, SERUM						
TOTAL PROTEIN	7.8	6.4 - 8.3	g/dL			
METHOD : BIURET						
ALBUMIN, SERUM						
ALBUMIN	4.8	3.5 - 5.2	g/dL			
METHOD: BROMOCRESOL GREEN						
GLOBULIN						
GLOBULIN	3.0	2.0 - 4.1	g/dL			
ELECTROLYTES (NA/K/CL), SERUM						
SODIUM, SERUM	142.0	136.0 - 146.0	mmol/L			
METHOD: DIRECT ION SELECTIVE ELECTRODE						
POTASSIUM, SERUM	4.34	3.50 - 5.10	mmol/L			
METHOD: DIRECT ION SELECTIVE ELECTRODE						
CHLORIDE, SERUM	103.3	98.0 - 106.0	mmol/L			
METHOD: DIRECT ION SELECTIVE ELECTRODE						







CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA

8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: 0290WD00043 AGE: 35 Years SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 03/04/2023 16:46 DRAWN: REPORTED:

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval Units** <u>Final</u>

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in: CCF, cirrhosis,	Decreased in: Low potassium	Decreased in: Vomiting, diarrhea,
vomiting, diarrhea, excessive	intake,prolonged vomiting or diarrhea,	renal failure combined with salt
sweating, salt-losing	RTA types I and II,	deprivation, over-treatment with
nephropathy, adrenal insufficiency,	hyperaldosteronism, Cushing's	diuretics, chronic respiratory acidosis,
nephrotic syndrome, water	syndrome,osmotic diuresis (e.g.,	diabetic ketoacidosis, excessive
intoxication, SIADH. Drugs:	hyperglycemia),alkalosis, familial	sweating, SIADH, salt-losing
thiazides, diuretics, ACE inhibitors,	periodic paralysis,trauma	nephropathy, porphyria, expansion of
chlorpropamide,carbamazepine,anti	(transient).Drugs: Adrenergic agents,	extracellular fluid volume,
depressants (SSRI), antipsychotics.	diuretics.	adrenalinsufficiency,
		hyperaldosteronism, metabolic
		alkalosis. Drugs: chronic
		laxative,corticosteroids, diuretics.
Increased in: Dehydration	Increased in: Massive hemolysis,	Increased in: Renal failure, nephrotic
(excessivesweating, severe	severe tissue damage, rhabdomyolysis,	syndrome, RTA,dehydration,
vomiting or diarrhea),diabetes	acidosis, dehydration,renal failure,	overtreatment with
mellitus, diabetesinsipidus,	Addison's disease, RTA type IV,	saline,hyperparathyroidism, diabetes
hyperaldosteronism, inadequate	hyperkalemic familial periodic	insipidus, metabolic acidosis from
water intake. Drugs: steroids,	paralysis. Drugs: potassium salts,	diarrhea (Loss of HCO3-), respiratory
licorice,oral contraceptives.	potassium- sparing diuretics,NSAIDs,	alkalosis, hyperadrenocorticism.
	beta-blockers, ACE inhibitors, high-	Drugs: acetazolamide,androgens,
	dose trimethoprim-sulfamethoxazole.	hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or	Interferences: Hemolysis of sample,	Interferences:Test is helpful in
hyperproteinemi, if sodium analysis	delayed separation of serum,	assessing normal and increased anion
involves a dilution step can cause	prolonged fist clenching during blood	gap metabolic acidosis and in
spurious results. The serum sodium	drawing, and prolonged tourniquet	distinguishing hypercalcemia due to
falls about 1.6 mEq/L for each 100	placement. Very high WBC/PLT counts	hyperparathyroidism (high serum
mg/dL increase in blood glucose.	may cause spurious. Plasma potassium	chloride) from that due to malignancy
	levels are normal.	(Normal serum chloride)

PHYSICAL EXAMINATION, URINE

COLOR	PALE YELLOW	
APPEARANCE	CLEAR	
CHEMICAL EXAMINATION, URINE		
PH	5.5	4.7 - 7.5
SPECIFIC GRAVITY	<=1.005	1.003 - 1.035
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
BILIRUBIN	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED
MICROSCOPIC EXAMINATION, URINE		
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED



Scan to View Report

/HPF

Page 7 Of 18





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA

8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND

PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results	Biological Reference Interva	l Units
PUS CELL (WBC'S)		2-3	0-5	/HPF
EPITHELIAL CELLS		2-3	0-5	/HPF
CASTS		NOT DETECTED		
CRYSTALS		NOT DETECTED		
BACTERIA		NOT DETECTED	NOT DETECTED	
YEAST		NOT DETECTED	NOT DETECTED	
REMARKS		Please note that all the uri	nary findings are confirmed manu	ually as well.

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions			
Proteins	Inflammation or immune illnesses			
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind			
	of kidney impairment			
Glucose	Diabetes or kidney disease			
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst			
Urobilinogen	Liver disease such as hepatitis or cirrhosis			
Blood	Renal or genital disorders/trauma			
Bilirubin	Liver disease			
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary			
	tract infection and glomerular diseases			
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either			
	acute or chronic, polycystic kidney disease, urolithiasis, contamination by			
	genital secretions			
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or			
	bladder catheters for prolonged periods of time			
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein			
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal			
	diseases			
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous			
	infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl			
	oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of			
	ethylene glycol or of star fruit (Averrhoa carambola) or its juice			
Uric acid	arthritis			
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.			
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis			

THYROID PANEL, SERUM









CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA

8800465156

SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference	Biological Reference Interval Units	
T3	111.40	80.0 - 200.0	ng/dL	
METHOD: CHEMILUMINESCENCE TECHNOLOGY	111.40	00.0 - 200.0	lig/uL	
T4	8.60	5.10 - 14.10	μg/dL	
METHOD: CHEMILUMINESCENCE TECHNOLOGY				
TSH (ULTRASENSITIVE)	1.730	0.270 - 4.200	μIU/mL	
METHOD · CHEMILLIMINESCENCE TECHNOLOGY				

METHOD: CHEMILUMINESCENCE TECHNOLOGY

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions	
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)	
					Post Thyroidectomy (4) Post Radio-Iodine treatment	
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid	
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto	
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical	
					inflammation, drugs like amphetamines, Iodine containing drug and	
					dopamine antagonist e.g. domperidone and other physiological reasons.	
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism	
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre	
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid	
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4	
					replacement therapy (7) First trimester of Pregnancy	
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism	
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor	
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent	
					treatment for Hyperthyroidism	
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness	
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies	

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011. **NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.**TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

PHYSICAL EXAMINATION, STOOL



Page 9 Of 18

Scan to View Details







CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Ir	nterval Units
COLOUR	BROWN		
CONSISTENCY	WELL FORMED		
MUCUS	ABSENT	NOT DETECTED	
VISIBLE BLOOD	ABSENT	ABSENT	
ADULT PARASITE	NOT DETECTED		
CHEMICAL EXAMINATION, STOOL			
OCCULT BLOOD	NOT DETECTED	NOT DETECTED	
MICROSCOPIC EXAMINATION, STOOL			
PUS CELLS	1-2		/hpf
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CYSTS	NOT DETECTED	NOT DETECTED	
OVA	NOT DETECTED		
LARVAE	NOT DETECTED	NOT DETECTED	
TROPHOZOITES	NOT DETECTED	NOT DETECTED	









CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

0290WD00043 AGE: 35 Years ACCESSION NO: SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 DRAWN: REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units <u>Final</u>

Interpretation(s)

Stool routine analysis is only a screening test for disorders of gastrointentestinal tract like infection, malabsorption, etc. The following table describes the probable conditions, in which the analytes are present in stool.

PRESENCE OF	CONDITION
Pus cells	Pus in the stool is an indication of infection
Red Blood cells	Parasitic or bacterial infection or an inflammatory bowel condition such as ulcerative colitis
Parasites	Infection of the digestive system. Stool examination for ova and parasite detects presence of parasitic infestation of gastrointestinal tract. Various forms of parasite that can be detected include cyst, trophozoite and larvae. One negative result does not rule out the possibility of parasitic infestation. Intermittent shedding of parasites warrants examinations of multiple specimens tested on consecutive days. Stool specimens for parasitic examination should be collected before initiation of antidiarrheal therapy or antiparasitic therapy. This test does not detect presence of opportunistic parasites like Cyclospora, Cryptosporidia and Isospora species. Examination of Ova and Parasite has been carried out by direct and concentration techniques.
Mucus	Mucus is a protective layer that lubricates, protects& reduces damage due to bacteria or viruses.
Charcot-Leyden crystal	Parasitic diseases.
Ova & cyst	Ova & cyst indicate parasitic infestation of intestine.
Frank blood	Bleeding in the rectum or colon.
Occult blood	Occult blood indicates upper GI bleeding.
Macrophages	Macrophages in stool are an indication of infection as they are protective cells.
Epithelial cells	Epithelial cells that normally line the body surface and internal organs show up in stool when there is inflammation or infection.
Fat	Increased fat in stool maybe seen in conditions like diarrhoea or malabsorption.
pH	Normal stool pH is slightly acidic to neutral. Breast-fed babies generally have an acidic stool.

ADDITIONAL STOOL TESTS:

- 1. Stool Culture:- This test is done to find cause of GI infection, make decision about best treatment for GI infection & to find out if treatment for GI infection worked.
- Fecal Calprotectin: It is a marker of intestinal inflammation. This test is done to differentiate Inflammatory Bowel Disease (IBD) 2. from Irritable Bowel Syndrome (IBS).
- Fecal Occult Blood Test(FOBT): This test is done to screen for colon cancer & to evaluate possible cause of unexplained anaemia. 3.
- Clostridium Difficile Toxin Assay: This test is strongly recommended in healthcare associated bloody or waterydiarrhoea, due to 4. overuse of broad spectrum antibiotics which alter the normal GI flora.
- Biofire (Film Array) GI PANEL: In patients of Diarrhoea, Dysentry, Rice watery Stool, FDA approved, Biofire Film Array Test, (Real Time Multiplex PCR) is strongly recommended as it identifies organisms, bacteria, fungi, virus, parasite and other opportunistic pathogens, Vibrio cholera infections only in 3 hours. Sensitivity 96% & Specificity 99%.
- Rota Virus Immunoassay: This test is recommended in severe gastroenteritis in infants & children associated with watery diarrhoea, vomitting& abdominal cramps. Adults are also affected. It is highly contagious in nature.









CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Biological Reference Interval Units

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE O

METHOD: TUBE AGGLUTINATION

RH TYPE POSITIVE

METHOD: TUBE AGGLUTINATION

XRAY-CHEST

»» BOTH THE LUNG FIELDS ARE CLEAR

»» BOTH THE COSTOPHRENIC AND CARIOPHRENIC ANGELS ARE CLEAR

»» BOTH THE HILA ARE NORMAL

»» CARDIAC AND AORTIC SHADOWS APPEAR NORMAL»» BOTH THE DOMES OF THE DIAPHRAM ARE NORMAL

»» VISUALIZED BONY THORAX IS NORMAL

IMPRESSION NO ABNORMALITY DETECTED

TMT OR ECHO

TMT OR ECHO TMT TEST IS NEGATIVE.

ECG

ECG NORMAL SINUS RHYTHM

MEDICAL HISTORY

RELEVANT PRESENT HISTORY

RELEVANT PAST HISTORY

RELEVANT PERSONAL HISTORY

RELEVANT FAMILY HISTORY

OCCUPATIONAL HISTORY

HISTORY

HISTORY

NOT SIGNIFICANT

NOT SIGNIFICANT

NOT SIGNIFICANT

NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS 1.66 mts WEIGHT IN KGS. 71 Kgs

BMI & Weight Status as follows: kg/sqmts

Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese

GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE NORMAL PHYSICAL ATTITUDE NORMAL









CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030

DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Final Results Biological Reference Interval Units

GENERAL APPEARANCE / NUTRITIONAL STATUS OVERWEIGHT
BUILT / SKELETAL FRAMEWORK AVERAGE
FACIAL APPEARANCE NORMAL
SKIN NORMAL
UPPER LIMB NORMAL
LOWER LIMB NORMAL
NECK NORMAL

NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER

THYROID GLAND NOT ENLARGED

CAROTID PULSATION NORMAL TEMPERATURE AFEBRILE

PULSE 63/MIN, REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

BRUIT

RESPIRATORY RATE NORMAL

CARDIOVASCULAR SYSTEM

BP 124/80 MM HG mm/Hg

(SITTING) NORMAL NORMAL NORMAL ABSENT

RESPIRATORY SYSTEM

SIZE AND SHAPE OF CHEST

MOVEMENTS OF CHEST

BREATH SOUNDS INTENSITY

NORMAL

BREATH SOUNDS QUALITY VESICULAR (NORMAL)

ADDED SOUNDS ABSENT

PER ABDOMEN

PERICARDIUM

HEART SOUNDS

APEX BEAT

MURMURS

APPEARANCE NORMAL VENOUS PROMINENCE ABSENT

LIVER NOT PALPABLE SPLEEN NOT PALPABLE HERNIA NORMAL

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS NORMAL



Page 13 Of 18





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Biological Reference Interval Test Report Status Results Units <u>Final</u> CRANTAL NERVES NORMAL CEREBELLAR FUNCTIONS **NORMAL** SENSORY SYSTEM NORMAL MOTOR SYSTEM **NORMAL REFLEXES** NORMAL **MUSCULOSKELETAL SYSTEM SPINE NORMAL**

JOINTS NORMAL

BASIC EYE EXAMINATION
CONJUNCTIVA

CONJUNCTIVA NORMAL EYELIDS NORMAL EYE MOVEMENTS NORMAL CORNEA NORMAL

DISTANT VISION RIGHT EYE WITH GLASSES

6/6, WITH GLASSES NORMAL

DISTANT VISION LEFT EYE WITH GLASSES

6/6, WITH GLASSES NORMAL

NEAR VISION RIGHT EYE WITH GLASSES

N/6, WITHIN NORMAL LIMIT

NEAR VISION LEFT EYE WITH GLASSES

N/6, WITHIN NORMAL LIMIT

COLOUR VISION NORMAL

BASIC ENT EXAMINATION

EXTERNAL EAR CANAL

TYMPANIC MEMBRANE

NORMAL

NOSE NO ABNORMALITY DETECTED

SINUSES NORMAL

THROAT NO ABNORMALITY DETECTED

TONSILS NOT ENLARGED

SUMMARY

RELEVANT HISTORY NOT SIGNIFICANT RELEVANT GP EXAMINATION FINDINGS OVERWEIGHT

REMARKS / RECOMMENDATIONS NONE

FITNESS STATUS

FITNESS STATUS FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)









CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

0290WD00043 AGE: 35 Years ACCESSION NO: SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

Comments

CLINICAL FINDINGS:-

ESTIMATED AVERAG GLUCOSE (EAG)

DYSLIPIDEMIA.

OVER WEIGHT STATUS.

FITNESS STATUS :-

FITNESS STATUS: FIT (WITH MEDICAL ADVICE) (AS PER REQUESTED PANEL OF TESTS)

ADVICE: WEIGHT REDUCTION, LOW FAT& CARBOHYDRATE DIET AND REGULAR PHYSICAL EXERCISE FOR OVERWEIGHT STATUS AND

DYSLIPIDEMIA

NEED PHYSICIAN CONSULTATION FOR LIFE STYLE MODIFICATION.

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504

This ratio element is a calculated parameter and out of NABL scope. ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-**TEST DESCRIPTION**:-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change **TEST INTERPRETATION**

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging,

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias,

Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis). In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

LIMITATIONS


False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased: Poikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs (Quinine,

salicylates)

REFERENCE :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis,10th edition. GLUCOSE FASTING,FLUORIDE PLASMA-**TEST DESCRIPTION**











CLIENT CODE: C000138355

CLIENT'S NAME AND ADDRESS: ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

0290WD00043 AGE: 35 Years ACCESSION NO: SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 DRAWN: REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

Increased in:Diabetes mellitus, Cushing's syndrome (10 – 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in: Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases (e.g. galactosemia), Drugs-insulin, ethanol, propranolol sulfonylureas,tolbutamide,and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals.Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment,Renal Glyosuria,Glycaemic index & response to food consumed,Alimentary Hypoglycemia,Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-**Used For**:

- 1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- 2. Diagnosing diabetes.
- 3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

- 1. eAG (Estimated average glucose) converts percentage HbA1c to md/dl, to compare blood glucose levels.
- 2. eAG gives an evaluation of blood glucose levels for the last couple of months. 3. eAG is calculated as eAG (mg/dl) = 28.7 * HbA1c 46.7

- **HbA1c Estimation can get affected due to :**1. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic
- anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days. 2. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.
- 3. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addiction are reported to interfere with some assay methods, falsely increasing results.
- 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.
- b) Heterozygous state detected (D10 is corrected for HbS & HbC trait.)
 c) HbF > 25% on alternate paltform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is

c) HDF > 25% on alternate pailtorm (Boronate arrinity chromatography) is recommended for detecting a hemoglobin electropnoresis (HPLC method) is recommended for detecting a hemoglobinopathy GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glyosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.Additional test HbA1c LIVER FUNCTION PROFILE, SERUM
Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, hemolysis and ineffective erythropoiesis).

obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic

hepatitis, obstruction of bile ducts, cirrhosis. **ALP** is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilsons disease.

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, billiary system and pancreas. Conditions that increase serum GGT are obstructive

liver disease,high alcohol consumption and use of enzyme-inducing drugs etc. **Total Protein** also known as total protein,is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma.It is produced in the liver.Albumin constitutes about half of the blood serum protein.Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.
CREATININE, SERUM-Higher than normal level may be due to:









CLIENT CODE: C000138355 **CLIENT'S NAME AND ADDRESS:**

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI

SOUTH WEST DELHI **NEW DELHI 110030 DELHI INDIA** 8800465156

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND PATIENT ID: ADITM2212877

0290WD00043 AGE: 35 Years ACCESSION NO: SEX: Male ABHA NO:

RECEIVED: 03/04/2023 08:28 DRAWN: REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status Results **Biological Reference Interval** Units <u>Final</u>

• Blockage in the urinary tract, Kidney problems, such as kidney damage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as breakdown of muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia) Lower than normal level may be due to:

• Myasthenia Gravis, Muscuophy
URIC ACID, SERUM-Causes of Increased levels:-Dietary(High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome Causes of decreased levels-Low Zinc intake,OCP,Multiple Sclerosis

TOTAL PROTEIN, SERUM-is a biochemical test for measuring the total amount of protein in serum.Protein in the plasma is made up of albumin and globulin.

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc.

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface

of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.

FITNESS STATUS-Conclusion on an individual's Fitness, which is commented upon mainly for Pre employment cases, is based on multi factorial findings and does not depend on any one single parameter. The final Fitness assigned to a candidate will depend on the Physician's findings and overall judgement on a case to case basis, details of the candidate's past and personal history as well as the comprehensiveness of the diagnostic panel which has been requested for .These are then further correlated with details

of the job under consideration to eventually fit the right man to the right job.

Basis the above, SRL classifies a candidate's Fitness Status into one of the following categories:

- Fit (As per requested panel of tests) SRL Limited gives the individual a clean chit to join the organization, on the basis of the General Physical Examination and the specific test panel requested for.
- Fit (with medical advice) (As per requested panel of tests) This indicates that although the candidate can be declared as FIT to join the job, minimal problems have been detected during the Pre- employment examination. Examples of conditions which could fall in this category could be cases of mild reversible medical abnormalities such as height weight disproportions, borderline raised Blood Pressure readings, mildly raised Blood sugar and Blood Lipid levels, Hematuria, etc. Most of these relate to sedentary lifestyles and come under the broad category of life style disorders. The idea is to caution an individual to bring about certain lifestyle changes as well as seek a Physician'''s
- Fitness on Hold (Temporary Unfit) (As per requested panel of tests) Candidate's reports are kept on hold when either the diagnostic tests or the physical findings reveal the presence of a medical condition which warrants further tests, counseling and/or specialist opinion, on the basis of which a candidate can either be placed into Fit, Fit (With Medical Advice), or Unfit category. Conditions which may fall into this category could be high blood pressure, abnormal ECG, heart murmurs, abnormal vision, grossly elevated blood sugars, etc.
- Unfit (As per requested panel of tests) An unfit report by SRL Limited clearly indicates that the individual is not suitable for the respective job profile e.g. total color blindness in color related jobs.



Scan to View Report

Page 17 Of 18





CLIENT CODE: C000138355
CLIENT'S NAME AND ADDRESS:

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)

F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI

NEW DELHI 110030 DELHI INDIA 8800465156 SRL LTD

Gate no 2, Residency Area, OPP. ST. Raphaels School,

INDORE, 452001 Madhya Pradesh, India Tel: 0731 2490008

PATIENT NAME: ADITYA BHAND

PATIENT ID: ADITM2212877

ACCESSION NO: **0290WD00043** AGE: 35 Years SEX: Male ABHA NO:

DRAWN: RECEIVED: 03/04/2023 08:28 REPORTED: 03/04/2023 16:46

REFERRING DOCTOR: DR. ACROFEMI HEALTHCARE LTD (MEDIWHEEL) CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results Units

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ULTRASOUND ABDOMEN ULTRASOUND ABDOMEN

.

Comments

USG-

IMPRESSION- EARLY FATTY INFILTRATION OF LIVER.

End Of Report
Please visit www.srlworld.com for related Test Information for this accession

Proite

Dr.Arpita Pasari, MD Consultant Pathologist



Dr.Meena Jinwah ,MBBS . MD Consultant Microbiologist

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
- 3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
- 4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form

- 5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
- 7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
- 8. Test results cannot be used for Medico legal purposes.
- 9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062



