Name	JAIN AMRIT KUMAR	ID	MED121762391
Age & Gender	37Year(s)/MALE	Visit Date	3/25/2023 12:00:00 AM
Ref Doctor Name	MediWheel	-	-

# ECHO CARDIOGRAPHIC STUDY

M-Mode						
AORTA			30		mm	
LEFT ARTIUM				27		mm
RIGHT VENTRICLE				05		mm
LEFT VENTRICLE (Diastole	)			46		mm
LEFT VENTRICLE (Systole)				27		mm
VENTRICULAR SEPTUM (I	Diastole)			07		mm
POSTERIOR WALL (Diastol	e)			07		mm
END DIASTOLIC VOLUME				97		ml
END SYSTOLIC VOLUME				26		ml
STROKE VOLUME				71		ml
FRACTIONAL SHORTENIN	[G			42		%
EJECTION FRACTION				73		%
	DOP	PLER / C	COL	OU	R FLOW	
MITRAL VALVE	E- 0.9	A -0.6	m/s	se	NO MR	
			c			
AORTIC VALVE	1.0	-	m/s	se	NO AR	
c						
TRICUSPID VALVE	-	- m/s		se	NO TR	
			c			
PULMONARY VALVE	0.8	-	m/s	se	NO PR	
			c			

**FINDINGS** 

Name	JAIN AMRIT KUMAR	ID	MED121762391
Age & Gender	37Year(s)/MALE	Visit Date	3/25/2023 12:00:00 AM
Ref Doctor Name	MediWheel	•	•


Name	JAIN AMRIT KUMAR	ID	MED121762391
Age & Gender	37Year(s)/MALE	Visit Date	3/25/2023 12:00:00 AM
Ref Doctor Name	MediWheel	-	-

	THICKNESS	NORMAL	
LV FUNCTION	REGIONAL WALL MOTION ABNORMALITY	ABSENT	
LEFT ATRIUM	NORMAL		
RIGHT VENTRICLE	NORMAL		
RIGHT ATRIUM	NORMAL		
MITRAL VALVE	NORMAL		
AORTIC VALVE	NORMAL		
PULMONARY VALVE	NORMAL		
TRICUSPID VALVE	NORMAL		
INTER ATRIAL SEPTUM	INTACT		
INTER VENTRICULAR SEPTUM	INTACT		
PERICARDIUM	NORMAL, NO EFFUSION		
GREAT VESSELS	AORTA - NORMAL PULMONARY ARTERY - NORMAL		

# **CONCLUSION:**

- NORMAL CHAMBER DIMENSIONS.
- NORMAL VALVES
- NO REGIONAL WALL MOTION ABNORMALITIES
- NORMAL LV SYSTOLIC FUNCTION.LVEF- 73 %
- NO CLOTS / VEGETATION / PE.

DR NAGESH M B
CONSULTANT
CARDIOLOGIST
(PLEASE CORRELATE WITH ECG & CLINICAL FINDINGS)



Name	JAIN AMRIT KUMAR	ID	MED121762391
Age & Gender	37Year(s)/MALE	Visit Date	3/25/2023 12:00:00 AM
Ref Doctor Name	MediWheel		

### ABDOMINO-PELVIC ULTRASONOGRAPHY

**LIVER** is normal in shape, size (11.5cms) and has increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

**GALL BLADDER** is partially distended and shows clear contents. No evidence of calculus. CBD is of normal calibre.

**PANCREAS** has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

**SPLEEN** shows normal shape, size and echopattern.

#### **BOTH KIDNEYS**

**Right kidney:** Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

**Left kidney:** Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	9.7	1.4
Left Kidney	10.1	1.3

**URINARY BLADDER** shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

**PROSTATE** shows normal shape, size and echopattern.

No evidence of ascites.

## **IMPRESSION:**

- Grade I fatty infiltration of liver.
- No other significant sonological abnormality detected.

DR. HEMANANDINI V.N CONSULTANT RADIOLOGIST Hn/Lr

Name	JAIN AMRIT KUMAR	ID	MED121762391
Age & Gender	37Year(s)/MALE	Visit Date	3/25/2023 12:00:00 AM
Ref Doctor Name	MediWheel		-

Name	JAIN AMRIT KUMAR	Customer ID	MED121762391
Age & Gender	37Y/M	Visit Date	Mar 25 2023 10:03AM
Ref Doctor	MediWheel		

# X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

## **IMPRESSION**:

• No significant abnormality detected.

G, Kamel DR.G KAMESH CONSULTANT RADIOLOGIST

 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
<b>HAEMATOLOGY</b>			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	16.1	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	48.4	%	42 - 52
RBC Count (EDTA Blood)	5.31	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	91.1	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	30.4	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.4	g/dL	32 - 36
RDW-CV	12.9	%	11.5 - 16.0
RDW-SD	41.13	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	5600	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	55.4	%	40 - 75
Lymphocytes (Blood)	28.3	%	20 - 45
Eosinophils (Blood)	7.6	%	01 - 06
Monocytes (Blood)	7.6	%	01 - 10





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>				
Basophils (Blood)	1.1	%	00 - 02				
INTERPRETATION: Tests done on Automated Five Part cell counter. All abnormal results are reviewed and confirmed microscopically							
Absolute Neutrophil count (EDTA Blood)	3.10	10^3 / μl	1.5 - 6.6				
Absolute Lymphocyte Count (EDTA Blood)	1.58	10^3 / μl	1.5 - 3.5				
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.43	10^3 / μl	0.04 - 0.44				
Absolute Monocyte Count (EDTA Blood)	0.43	10^3 / μl	< 1.0				
Absolute Basophil count (EDTA Blood)	0.06	10^3 / μl	< 0.2				
Platelet Count (EDTA Blood)	255	10^3 / μl	150 - 450				
MPV (Blood)	7.2	fL	7.9 - 13.7				
PCT (Automated Blood cell Counter)	0.18	%	0.18 - 0.28				
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	3	mm/hr	< 15				





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<b>BIOCHEMISTRY</b>			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.45	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.17	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	0.28	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	20	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	23	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	10	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	79	U/L	53 - 128
Total Protein (Serum/Biuret)	6.37	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.07	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.30	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.77		1.1 - 2.2





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	138.64	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	99.23	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

**INTERPRETATION:** The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual\_circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	38.09	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	80.7	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	19.8	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	100.5	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Total Cholesterol/HDL Cholesterol Ratio (Serum/Calculated)	3.6		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/Calculated)	2.6		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.1		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0





APPROVED BY

 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	4.8	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 91.06 mg/dL

(Whole Blood)

#### **INTERPRETATION: Comments**

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

<u>Investigation</u>	Observed Unit	<u>Biological</u>
	Value	Reference Interval

## **IMMUNOASSAY**

#### THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.54 ng/ml 0.7 - 2.04

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total 6.18 μg/dl 4.2 - 12.0

(Serum/ECLIA)

#### INTERPRETATION:

#### **Comment:**

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.49 µIU/mL 0.35 - 5.50

(Serum/ECLIA)

#### INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

#### Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&amplt,0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	<u>Value</u>		Reference Interval

## **CLINICAL PATHOLOGY**

# <u>PHYSICAL EXAMINATION (URINE</u> <u>COMPLETE)</u>

Colour Yellow Yellow Yellow to Amber

(Urine)

Appearance Clear Clear

(Urine)

Volume(CLU) 30

(Urine)

## CHEMICAL EXAMINATION (URINE

**COMPLETE**)

pH 6 4.5 - 8.0

(Urine)

Specific Gravity 1.020 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)

Glucose Negative Negative

(Urine/GOD - POD)





APPROVED BY

 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
Leukocytes(CP) (Urine)	Negative		
MICROSCOPIC EXAMINATION (URINE COMPLETE)			
Pus Cells (Urine)	0-1	/hpf	NIL
Epithelial Cells (Urine)	0-1	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others (Urine)	NIL		

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.





 PID No.
 : MED121762391
 Register On
 : 25/03/2023 10:03 AM

 SID No.
 : 522304612
 Collection On
 : 25/03/2023 11:15 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 25/03/2023 6:05 PM

 Type
 : OP
 Printed On
 : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

InvestigationObservedUnitBiologicalValueReference Interval

## **IMMUNOHAEMATOLOGY**

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.





PID No. : MED121762391 Register On : 25/03/2023 10:03 AM : 522304612 SID No. Collection On : 25/03/2023 11:15 AM Age / Sex : 37 Year(s) / Male Report On 25/03/2023 6:05 PM **Type** : OP **Printed On** : 26/03/2023 11:37 AM

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<b>BIOCHEMISTRY</b>			
BUN / Creatinine Ratio	9.4		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	85.71	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine)	Negative		Negative
(Urine - F/GOD - POD)			
Glucose Postprandial (PPBS)	93.33	mg/dL	70 - 140

(Plasma - PP/GOD-PAP)

#### INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Blood Urea Nitrogen (BUN)	7.9	mg/dL	7.0 - 21
(Serum/Urease UV / derived)			
Creatinine	0.84	mg/dL	0.9 - 1.3
(Serum/Modified Jaffe)			

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists, N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 3.57 mg/dL 3.5 - 7.2

(Serum/Enzymatic)





**APPROVED BY** 

-- End of Report --