Name	Ms.SNIGDHA DAS P	ID	MED121139177
Age & Gender	43/FEMALE	Visit Date	18/06/2022
Ref Doctor Name	MediWheel		

X-ray mammogram (mediolateral oblique & craniocaudal views) followed by Sonomammography.

BILATERAL MAMMOGRAPHY

Breast composition Type B (These are scattered areas of fibroglandular density).

Macrocalcification is noted in the upper quadrant of left breast.

No evidence of cluster microcalcification.

Subcutaneous fat deposition is within normal limits.

BILATERAL SONOMAMMOGRAPHY

Right breast shows well defined hypoechoic wider than tall lesion at 10 o' clock position, measuring about 6.9 x 3.4 mm.

Both the breasts otherwise show normal echopattern.

No other focal solid / cystic areas.

No evidence of ductal dilatation.

No evidence of axillary lymphadenopathy on both sides.

IMPRESSION:

- **Hypoechoic lesion in right breast - likely fibroadenoma.**
- **No other significant abnormality detected.**

ASSESSMENT: BI-RADS CATEGORY - 3

BI-RADS CLASSIFICATION

CATEGORY RESULT

Name	Ms.SNIGDHA DAS P	ID	MED121139177
Age & Gender	43/FEMALE	Visit Date	18/06/2022
Ref Doctor Name	MediWheel		

3 **Probably benign finding. Short interval follow-up suggested.**

DR. HEMANANDINI V.N

CONSULTANT RADIOLOGIST

Hn/Ss

REPORT DISCLAIMER

1.This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.

2.The results reported here in are subject to interpretation by qualified medical professionals only.

3.Customer identities are accepted provided by the customer or their representative.

4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

6.Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

9.Liability is limited to the extend of amount billed.

10.Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

11.Disputes,if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/*Spectrophotometry*)	**12.30**	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	37.2	%	37 - 47
RBC Count (EDTA Blood)	4.51	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood)	82.4	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	27.3	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	33.1	g/dL	32 - 36
RDW-CV	14.0	%	11.5 - 16.0
RDW-SD	**40.38**	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	6920	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	71.98	%	40 - 75
Lymphocytes (Blood)	21.79	%	20 - 45
Eosinophils (Blood)	1.65	%	01 - 06

Dr.Arjun C.P
MBBS,MD Pathology
Reg No:KMC 89655

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
Monocytes (Blood)	4.38	%	01 - 10
Basophils (Blood)	0.20	%	00 - 02
INTERPRETATION: Tests done on Automated Five Part cell counter. All abnormal results are reviewed and confirmed microscopically.			
Absolute Neutrophil count (EDTA Blood)	4.98	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	1.51	10^3 / µl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.11	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.30	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood)	0.01	10^3 / µl	< 0.2
Platelet Count (EDTA Blood)	351.5	10^3 / µl	150 - 450
MPV (Blood)	8.22	fL	8.0 - 13.3
PCT (*Automated Blood cell Counter*)	**0.29**	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	**21**	mm/hr	< 20

Dr.Arjun C.P
MBBS,MD Pathology
Reg No:KMC 89655

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/*DCA with ATCS)*	0.37	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/*Diazotized Sulfanilic Acid)*	0.10	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/*Derived)*	0.27	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/*Modified IFCC)*	19.96	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/*Modified IFCC)*	10.76	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/*IFCC / Kinetic)*	12.07	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/*Modified IFCC)*	89.1	U/L	42 - 98
Total Protein (Serum/*Biuret)*	7.39	gm/dl	6.0 - 8.0
Albumin (Serum/*Bromocresol green)*	4.86	gm/dl	3.5 - 5.2
Globulin (Serum/*Derived)*	2.53	gm/dL	2.3 - 3.6
A : G RATIO (Serum/*Derived)*	1.92		1.1 - 2.2

Dr SURAJ JAIN
Consultant Pathologist
Reg No : 80423

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
Lipid Profile			
Cholesterol Total (Serum/*CHOD-PAP with ATCS)*	**205.71**	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/*GPO-PAP with ATCS)*	98.42	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_ circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/*Immunoinhibition)*	**49.33**	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/*Calculated)*	136.7	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/*Calculated)*	19.7	mg/dL	< 30
Non HDL Cholesterol (Serum/*Calculated)*	156.4	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220

Dr SURAJ JAIN
Consultant Pathologist
Reg No : 80423

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval

INTERPRETATION: 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol.
2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

Total Cholesterol/HDL Cholesterol Ratio (Serum/*Calculated*)	4.2		Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/*Calculated*)	2		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/*Calculated*)	2.8		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0

Dr SURAJ JAIN
Consultant Pathologist
Reg No : 80423

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/*HPLC)*	**5.8**	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control : 6.1 - 7.0 % , Fair control : 7.1 - 8.0 % , Poor control >= 8.1 %

Estimated Average Glucose (Whole Blood*)*	119.76	mg/dL	

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency, hypertriglyceridemia,hyperbilirubinemia,Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly,Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

Dr SURAJ JAIN
Consultant Pathologist
Reg No : 80423

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total (Serum/*ECLIA*)	1.24	ng/ml	0.7 - 2.04

INTERPRETATION:
Comment :
Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Tyroxine) - Total (Serum/*ECLIA*)	9.20	µg/dl	4.2 - 12.0

INTERPRETATION:
Comment :
Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) (Serum/*ECLIA*)	2.12	µIU/mL	0.35 - 5.50

INTERPRETATION:
Reference range for cord blood - upto 20
1 st trimester: 0.1-2.5
2 nd trimester 0.2-3.0
3 rd trimester : 0.3-3.0
(Indian Thyroid Society Guidelines)
Comment :
1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM.The variation can be of the order of 50%,hence time of the day has influence on the measured serum TSH concentrations.
3.Values&amplt;0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.

DR .VANITHA.R.SWAMY MD
Consultant Pathologist
Reg No : 99049
VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902
APPROVED BY

Investigation	Observed Value	Unit	Biological Reference Interval
CLINICAL PATHOLOGY			
PHYSICAL EXAMINATION (URINE COMPLETE)			
Colour (Urine)	Pale yellow		Yellow to Amber
Appearance (Urine)	Clear		Clear
Volume(CLU) (Urine)	20		
CHEMICAL EXAMINATION (URINE COMPLETE)			
pH (Urine)	7.0		4.5 - 8.0
Specific Gravity (Urine)	1.003		1.002 - 1.035
Ketone (Urine)	Negative		Negative
Urobilinogen (Urine)	Normal		Normal
Blood (Urine)	Negative		Negative
Nitrite (Urine)	Negative		Negative
Bilirubin (Urine)	Negative		Negative
Protein (Urine)	Negative		Negative

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
Glucose (Urine/*GOD - POD*)	Negative		Negative
Leukocytes(CP) (Urine)	Negative		
MICROSCOPIC EXAMINATION (URINE COMPLETE)			
Pus Cells (Urine)	**0-1**	/hpf	NIL
Epithelial Cells (Urine)	**0-1**	/hpf	NIL
RBCs (Urine)	NIL	/HPF	NIL
Others (Urine)	NIL		

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**				
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM		
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM		
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM		
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM		
Ref. Dr	: **MediWheel**				

Investigation	Observed Value	Unit	Biological Reference Interval

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING	'B' 'Positive'		
(EDTA Blood/*Agglutination)*			

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.

Dr.Arjun C.P
MBBS,MD Pathology
Reg No:KMC 89655

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

Name	: **Ms. SNIGDHA DAS P**		
PID No.	: MED121139177	**Register On**	: 18/06/2022 10:24 AM
SID No.	: 522219391	**Collection On**	: 18/06/2022 11:33 AM
Age / Sex	: 43 Year(s) / Female	**Report On**	: 18/06/2022 7:57 PM
Type	: OP	**Printed On**	: 25/06/2022 5:09 PM
Ref. Dr	: **MediWheel**		

MEDALL

Investigation	Observed Value	Unit	Biological Reference Interval
BIOCHEMISTRY			
BUN / Creatinine Ratio	10.69		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/*GOD-PAP)*	**105.28**	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/*GOD - POD)*	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/*GOD-PAP)*	111.19	mg/dL	70 - 140

INTERPRETATION:
Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP*)*	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/*Urease UV / derived)*	7.7	mg/dL	7.0 - 21
Creatinine (Serum/*Modified Jaffe)*	0.72	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/*Enzymatic)*	5.21	mg/dL	2.6 - 6.0

Dr SURAJ JAIN
Consultant Pathologist
Reg No : 80423

VERIFIED BY

DR SHAMIM JAVED
MD PATHOLOGY
KMC 88902

APPROVED BY

-- End of Report --