



LABORATORY REPORT



Name : **Mr. Vijaysingh M. Dudi** Registration on : 25-Dec-2021 09:23
 Lab ID : **122121202267** Ref. Id : Collected on :
 Sex/Age : **Male / 29 Years** Approved on : 25-Dec-2021 11:08
 Ref. By : Sample Type : EDTA Blood
 Location : Bob Health Checkup@Godhra Patient Source :

COMPLETE BOOD COUNT

Test	Result	Unit	Biological Ref. Interval
HB and Indices			
Hemoglobin	16.4	g/dL	13.0 - 16.5
RBC Count	H 5.87	million/cmm	4.5 - 5.5
Hematocrit	49.0	%	40 - 49
MCV	83.5	fL	83 - 101
MCH	27.9	pg	27.1 - 32.5
MCHC	33.5	g/dL	32.5 - 36.7
RDW CV	12.30	%	11.6 - 14
Total WBC and Differential Count			
WBC Count	8380	/cmm	4000 - 10000
Differential Count		Absolute Count	
Neutrophils	56.7	% 40 - 80	4751 /cmm 2000 - 6700
Lymphocytes	34.8	% 20 - 40	2916 /cmm 1000 - 3000
Eosinophils	2.3	% 1 - 6	193 /cmm 20 - 500
Monocytes	5.4	% 2 - 10	453 /cmm 200 - 1000
Basophils	0.8	% 0 - 2	67 /cmm 0 - 100
Platelet Count			
Platelet Count	311000	/cmm	150000 - 410000
Erythrocytes Sedimentation Rate			
ESR	03	mm/1hr	0 - 14

Dr. Harsh shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 1 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : Mr. Vijaysingh M. Dudi Registration on : 25-Dec-2021 09:23
Lab ID : 122121202267 Ref. Id : Collected on : 25-Dec-2021 16:00
Sex/Age : Male / 29 Years Approved on : 25-Dec-2021 10:59
Ref. By : Sample Type : Serum, Fluoride PP
Location : Bob Health Checkup@Godhra Patient Source :

Test	Result	Unit	Biological Ref. Interval
Fasting Blood Sugar	93.3		70 - 110
Fasting Urine Sugar	Absent		Absent
Post Prandial Blood Sugar <i>GOD-POD</i>	124.9	mg/dL	70 - 140
Postprandial Urine Sugar	Absent		Absent
Creatinine, Serum <i>Modified Jaffes method</i>	0.99	mg/dL	0.4 - 1.4
Urea <i>Urease Glutamate Dehydrogenase, UV method</i>	22.7	mg/dL	19 - 45
Blood Urea Nitrogen <i>Calculated</i>	10.61	mg/dL	9.0 - 20.0
Uric Acid <i>Uricase-Peroxidase method</i>	6.00	mg/dL	3.5 - 7.2
SGPT <i>IFCC method without pyridoxal phosphate activation</i>	19.7	U/L	0 - 45
SGOT <i>IFCC method without pyridoxal phosphate activation</i>	11.1	U/L	5 - 40
GGT <i>L-?-glutamyl-glycylglycine</i>	27.0	U/L	5 - 50
Alkaline Phosphatase <i>NPP-AMP Buffer</i>	L 49.0	U/L	53 - 128
Bilirubin			
Total Bilirubin <i>Diazo reaction</i>	0.23	mg/dL	0.2 - 1.3
Direct Bilirubin <i>Diazo reaction</i>	0.18	mg/dL	0.0 - 0.4
Indirect Bilirubin <i>Calculated</i>	L 0.05	mg/dL	0.1 - 1.1
Protein			
Total Protein <i>Biuret method</i>	8.12	g/dL	6.0 - 8.5
Albumin <i>BCG</i>	4.75	g/dL	3.5 - 5.2
Globulin	H 3.37	g/dL	2.2 - 3.0
A/G Ratio <i>Calculated</i>	1.41		1.3 - 1.7

Dr. Harsh Shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 2 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name	: Mr. Vijaysingh M. Dudi	Registration on	: 25-Dec-2021 09:23
Lab ID	: 122121202267 Ref. Id :	Collected on	: 25-Dec-2021 16:00
Sex/Age	: Male / 29 Years	Approved on	: 25-Dec-2021 10:59
Ref. By	:	Sample Type	: Serum, Fluoride PP
Location	: Bob Health Checkup@Godhra	Patient Source	:

Dr. Harsh Shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 3 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : Mr. Vijaysingh M. Dudi Registration on : 25-Dec-2021 09:23
Lab ID : 122121202267 Ref. Id : Collected on : 25-Dec-2021 16:00
Sex/Age : Male / 29 Years Approved on : 25-Dec-2021 10:59
Ref. By : Sample Type : Serum
Location : Bob Health Checkup@Godhra Patient Source :

Lipid Profile

Test	Result	Unit	Biological Ref. Interval
Cholesterol <i>Cholesterol oxidase, Esterase, Peroxidase</i>	158.0	mg/dL	Desirable : <200 Borderline High : 200-239 High : >240
Triglyceride <i>GPC-POD</i>	53.2	mg/dL	Normal : < 150 Borderline : 150-199 High : 200-499 Very High : > 500
HDL Cholesterol <i>PTA/MgCl2</i>	L 39.0	mg/dL	Low : <40.0 High : >60.0
Direct LDL <i>Direct measured</i>	95.70	mg/dL	Optimal : < 100 Near / above optimal : 100-129 Borderline High : 130-159 High : 160-189 Very High : >190
VLDL <i>Calculated</i>	L 10.64	mg/dL	15 - 35
CHOL/HDL Ratio <i>Calculated</i>	4.1		Up to 5.0
LDL/HDL Ratio <i>Calculated</i>	2.5		Up to 3.5

Dr. Harsh Shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 4 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : Mr. Vijaysingh M. Dudi Registration on : 25-Dec-2021 09:23
Lab ID : 122121202267 Ref. Id : Collected on : 25-Dec-2021 16:00
Sex/Age : Male / 29 Years Approved on : 25-Dec-2021 11:25
Ref. By : Sample Type : EDTA Sample
Location : Bob Health Checkup@Godhra Patient Source :

HbA1c (Glycosylated Hemoglobin)

Test	Result	Unit	Biological Ref. Interval
HbA1c	H 6.50	%	For Screening: Diabetes: >6.5% Pre-Diabetes: 5.7% - 6.4% Non-Diabetes: < 5.7% For Diabetic Patient: Poor Control : > 7.0 % Good Control : 6.0-7.0 %
Mean Blood Glucose <small>Calculated</small>	139.85	mg/dL	

Explanation:-

- Total haemoglobin A1 c is continuously synthesized in the red blood cell through its 120 days life span. The concentration of HBA1c in the cell reflects the average blood glucose concentration it encounters.
- The level of HBA1c increases proportionately in patients with uncontrolled diabetes. It reflects the average blood glucose concentration over an extended time period and remains unaffected by short-term fluctuations in blood glucose levels.
- The measurement of HbA1c can serve as a convenient test for evaluating the adequacy of diabetic control and in preventing various diabetic complications. Because the average half life of a red blood cell is sixty days, HbA1c has been accepted as a measurement which reflects the mean daily blood glucose concentration, better than fasting blood glucose determination, and the degree of carbohydrate imbalance over the preceding two months.
- It may also provide a better index of control of the diabetic patient without resorting to glucose loading procedures.

HbA1c assay Interferences:

Erroneous values might be obtained from samples with abnormally elevated quantities of other Haemoglobins as a result of either their simultaneous elution with HbA1c (HbF) or differences in their glycation from that of HbA (HbS).

Reference: ADA Guideline 2020

Dr. Harsh Shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 5 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : **Mr. Vijaysingh M. Dudi** Registration on : 25-Dec-2021 09:23
Lab ID : **122121202267** Ref. Id : Collected on : 25-Dec-2021 16:00
Sex/Age : **Male / 29 Years** Approved on : 25-Dec-2021 11:12
Ref. By : Sample Type : Serum
Location : Bob Health Checkup@Godhra Patient Source :

Thyroid Function Test

Test	Result	Unit	Biological Ref. Interval
T3 - Triiodothyronine <small>C/MIA</small>	1.22	ng/mL	0.58 - 1.59
T4 - Thyroxine <small>C/MIA</small>	7.33	micro g/dL	5.13 - 14.06
TSH - Thyroid Stimulating Hormone <small>C/MIA</small>	2.2300	microIU/mL	0.35 - 4.94

TSH	T3/FT3	T4/FT4	Suggested Interpretation for the Thyroid Function Tests Pattern
Within Range	Decreased	Within Range	- Isolated Low T3-offen seen in elderly & associated Non-Thyroidal illness. In elderly the drop in T3 level can be upto 25%
Raised	Within Range	Within Range	- Isolated High TSH especially in the range of 4.7 to 15 mIU/ml is commonly associated with physiological & Biological TSH Variability. - Subclinical Autoimmune Hypothyroidism - Intermittent T4 therapy for hypothyroidism - Recovery phase after Non-Thyroidal illness
Raised	Decreased	Decreased	- Chronic autoimmune Thyroiditis - Post thyroidectomy, Post radiiodine - Hypothyroid phase of transient thyroiditis
Raised or Within Range	Raised	Raised or Within range	- Interfering antibodies to thyroid hormones (anti-TPO antibodies) - intermittent T4 therapy or T4 overdose - Drug interference-Amiodarone, Heparin, Beta blockers, steroids, anti-epileptics
Decreased	Raised or within Range	Raised or within Range	- Isolated Low TSH - especially in the range of 0.1 to 0.4 often seen in elderly & associated with Non-Thyroidal illness - Subclinical Hyperthyroidism - Thyroxine ingestion
Decreased	Decreased	Decreased	- Central Hypothyroidism - Non-Thyroidal illness - Recent treatment for Hyperthyroidism (TSH remains suppressed)
Decreased	Raised	Raised	- Primary Hyperthyroidism (Graves disease), Multinodular goitre Toxic nodule - Transient thyroiditis:Postpartum, Silent (lymphocytic), Postviral (granulomatous, subacute, DeQuervain'a) Gestational thyrotoxicosis with hyperemesis gravidarum
Decreased or within range	Raised	Within Range	- T3 toxicosis - Non-Thyroidal illness

Dr. Harsh shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 6 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : Mr. Vijaysingh M. Dudi Registration on : 25-Dec-2021 09:23
Lab ID : 122121202267 Ref. Id : Collected on : 25-Dec-2021 16:00
Sex/Age : Male / 29 Years Approved on : 25-Dec-2021 11:12
Ref. By : Sample Type : Serum
Location : Bob Health Checkup@Godhra Patient Source :

Immunoassay

Test	Result	Unit	Biological Ref. Interval
HIV I & II, CMIA <small>CMIA</small>	0.120	S/Co	Non reactive : <1.0 Reactive : >1.0

Additional Information:

1. A NON REACTIVE result implies that no Anti HIV-1 or HIV -2 antibodies have been detected in the sample by this method. This means that either the patient has not been exposed to HIV-1 or HIV-2 infection or the sample has been tested during the "WINDOW PHASE" (before the development of detectable levels of antibodies).
 2. A PROVISIONALITY REACTIVE / BORDERLINE REACTIVE result suggests possibility of HIV-1 or/and HIV-2 infection. However these results must be verified by confirmatory WESTERN BLOT / HIV PCR method before declaring the patient positive for HIV-1 or HIV-2 infection.
 3. Very high levels of IgM Antibodies or Anti-HLA ABC and DR Antibodies can give false positive reaction.
- **Pre & Post test counselling for HIV testing is responsibility of referring Physician.

Dr. Harsh shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 7 of 9

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : Mr. Vijaysingh M. Dudi	Registration on : 25-Dec-2021 09:23
Lab ID : 122121202267 Ref. Id :	Collected on : 25-Dec-2021 16:00
Sex/Age : Male / 29 Years	Approved on : 25-Dec-2021 11:12
Ref. By :	Sample Type : Serum
Location : Bob Health Checkup@Godhra	Patient Source :

Immunoassay

Test	Result	Unit	Biological Ref. Interval
HBsAg <small>CMIA</small>	0.170	IU/mL	Non reactive: < 1.0 Reactive : >1.0

Interpretation:

- HBsAg is the earliest marker of acute HBV infection which typically becomes detectable 2-3 months (as early as 14 days) after infection. When symptoms of hepatitis are present, most patients have detectable HBsAg although few patients will have neither HBsAg nor anti-HBs and anti-HBc IgM is the only marker of acute HBV infection (Core Window). HBsAg typically persists for 12-20 weeks after onset of symptoms in uncomplicated HBV infection and disappears followed by a small but variable gap with onset of anti-HBs (Seroconversion).
- Detection of HBsAg beyond 06 months defines chronic HBV infection or a chronic carrier state. Chronic HBV infection is seen in 1-2% of adults and adolescents following acute HBV infection, 5-10% of immunocompromised individuals and upto 80% of neonates. The chronic carrier state of HBV shows only persistent HBsAg in the serum without any other HBV marker or evidence of liver injury.
- Hepatitis B vaccination does not cause a positive HBsAg result. Quantitation or Titer of HBsAg is of no clinical value.
- Presence of anti-HBs without detectable HBsAg indicates recovery from acute HBV infection, absence of infectivity and immunity against future HBV infection.
- HBsAg test is carried out with Chemiluminescent Microparticle immunoassay (CMIA) which uses microparticles coated with monoclonal anti-HBs for the detection of HBsAg. HBsAg assays are routinely used to aid in the diagnosis of suspected hepatitis B viral (HBV) infection and to monitor the status of infected individuals.
- All initial reactive specimens are subjected to further testing by one or two additional methods and final report is issued in accordance with the same. Repeat reactive specimens MUST be confirmed by any combination of the confirmatory tests (e.g. HBsAg neutralization test, Other HBV markers & LFT and HBV DNA by PCR method).

Limitations:

- If the ARCHITECT HBsAg Qualitative II results are inconsistent with clinical evidence, additional testing is suggested to confirm the result.
- For diagnostic purposes, results should be used in conjunction with patient history and other hepatitis markers for diagnosis of acute and chronic infection.
- Specimens from patients who have received preparations of mouse monoclonal antibodies for diagnosis or therapy may contain human anti-mouse antibodies (HAMA). Specimens containing HAMA may produce anomalous values when tested with assay kits such as ARCHITECT HBsAg Qualitative II that employ mouse monoclonal antibodies.
- Heterophilic antibodies in human serum can react with reagent immunoglobulins, interfering with in vitro immunoassays. Patients routinely exposed to animals or to animal serum products can be prone to this interference and anomalous results may be observed. Additional information may be required for diagnosis.

Dr. Harsh Shah

M.D., D.N.B.

Referred Test

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 8 of 9

Report Status: Revised

Printed on: 04-Jan-2022 16:05



LABORATORY REPORT



Name : Mr. Vijaysingh M. Dudi	Registration on : 25-Dec-2021 09:23
Lab ID : 122121202267 Ref. Id :	Collected on : 25-Dec-2021 16:00
Sex/Age : Male / 29 Years	Approved on : 25-Dec-2021 11:15
Ref. By :	Sample Type : Urine
Location : Bob Health Checkup@Godhra	Patient Source :

Urine Routine Examination

Test	Result	Unit	Biological Ref. Interval
Physical Examination			
Volume	10	ml	
Colour	Yellow		
Odour	Ammonical		
Transparency	Clear		
Chemical Examination (Dip Stick Method)			
Reaction	Acidic		
Specific Gravity	1.025		1.005 - 1.030
Albumin	Absent		Negative
Urine Glucose	Absent		Absent
Bile Salts	Absent		Absent
Bile Pigments	Absent		Absent
Urine Ketone	Absent		Absent
Nitrite	Negative		Negative
Microscopic Examination			
Pus Cells	0-1	/hpf	0 - 5
Red Cells	Absent	/hpf	0 - 2
Epithelial Cells	Occassional	/hpf	
Casts	Absent	/hpf	
Crystals	Absent	/hpf	
Amorphous Material	Absent		
Bacteria	Absent		Absent
Budding Yeast	Absent		Absent
Trichomonas	Absent		

----- End Of Report -----

Dr. Harsh shah

M.D., D.N.B.

Referred Test

Report Status: Revised

Dr. Farzana Vali

M.B., D.C.P.

This is an Electronically Authenticated Report.

Dr Nayan Modi

M.D.

Page 9 of 9

Printed on: 04-Jan-2022 16:05