Kshipra Scans & Labs

Name	:	DILPREET SINGH		Y	-
Thanks To	1	MEDIWHEEL	Age	:	34 Yrs. / M
		and and the second s	Date	:	22/07/2023

### X-RAY CHEST (PA VIEW)

Both lung fields appear normal. No e/o Koch's lesion or consolidation seen. Both CP angles appear clear. Both domes of diaphragm appear normal. Heart size and aorta are within normal limits. Bony thorax under vision appears normal. Both hila appear normal.


Consultant Radiologist

(This report is not valid for any Medico-legal purpose)

2-B Ground Floor, Court Choraha, Main Road Tehsil ke Samne, Udaipur 313001 (Raj.), Mob. : 7229961115, 7229970005, 7229901188 (24 x 7 Customer Service) Email : kshipralabsudaipur@gmail.com

Scanned with AnyScanner

9166046591





Mr. Dilfreet singh

34/M

22/07/23

Dr. Sharva Pandya

MBBS, M.S., (Ophthalmology)

RMC Reg. No. : 021537

वरिष्ठ नेत्र रोग विरोषज्ञ सर्जन

डॉ. शर्वा पण्डुया

Clo - For eye checkup

Draz 6/6 6/6

NYAZ NIG

Colour viscon - Normal

Sharvy

Dr. SHARVA PANDYA MBBS, MS (Ophth.) RMC No. : 021537 JAI DRISHTI EYE HOSPITAL UDAIPUR (Rai)

चित्रकूट नगर, उदयपुर में हमारी नैत्र चिकित्सा सेवाएं शीघ्र उपलब्ध होगी।

Jai Drishti Eye Hospital, 23A, Residency Road, Opp. Equitas Bank, Sardarpura, Udaipur जय दृष्टि आई हॉस्पिटल, 23-ए, रेजीडेन्सी रोड़, एक्विटास बैंक के सामने, सरदारपुरा, उदयपुर 🛚



9166046591

# shipra Scans & Labs

<b>v</b>		Mr. Dilaroot Singh	Age	:	34Yrs. / M
Name		Mr. Dilpreet Singh			22.07.2023
Thanks To	:	Mediweel wellness	Date	·	

### ULTRASOUND STUDY OF WHOLE ABDOMEN

### LIVER

Liver is normal in size, shape & bright echotexture. No focal mass lesion is seen. Intra hepatic biliary radicles are normal. Portal vein is normal in caliber.

### GALL BLADDER

Gall bladder is well distended. The wall thickness appears normal. No evidence of calculus or mass lesion is seen. C.B.D. appears normal.

#### PANCREAS

Pancreas is normal in size, shape & echotexture. No focal mass lesion is seen.

#### SPLEEN

Spleen is normal in size, shape & echotexture. No focal mass lesion is seen.

#### **BOTH KIDNEYS**

Both kidneys are normal in size, shape & echotexture. Renal parenchyma appears normal. No evidence of hydronephrosis, calculus or cortical scarring is seen in either kidney.

: 9.9 x 4.7 cms. Right kidney measures : 9.5 x 5.3 cms. Left kidney measures

.

#### URINARY BLADDER

Urinary bladder is well distended and appears normal in contour. The wall thickness appears normal.

#### PROSTATE

Prostate is normal in size, shape and echotexture. It measures 2.9 x 3.8 x 2.2 cms. Volume: 13 cc.

No obvious abdominal lymphadenopathy is seen. No free fluid is seen in peritoneal cavity.


#### **OPINION:**

Fatty liver grade II

Dr. Tapendra N. Tiwari MD (Radio-Diagnosis) Consultant Radiologist

(This report is not valid for any Medico-legal purpose) ENCL: - PCPNDT Registration Certificate is printed on the back side of this report.

2-B Ground Floor, Court Choraha, Main Road Tehsil ke Samne, Udaipur 313001 (Raj.), Mob.: 7229961115, 7229970005, 7229901188 (24 x 7 Customer Service) Email : kshipralabsudaipur@gmail.com



## 

**TEST REPORT** 

Reg. No:2307101704Name:Dilpreet SinghAge/Sex:34 Years / MaleRef. By:Client:MEDIWHEEL WELLNESS

 Reg. Date
 : 22-Jul-2023

 Collected On
 : 22-Jul-2023 08:47

 Approved On
 : 22-Jul-2023 10:20

 Printed On
 : 25-Jul-2023 12:22

Parameter	<u>Result</u>	<u>Unit</u>	Reference Interval			
COMPLETE BLOOD COUNT (CBC)						
	SPECIMEN:	EDTA BLOOD				
Hemoglobin	14.3	g/dL	13.0 - 17.0			
RBC Count	5.02	million/cmm	4.5 - 5.5			
Hematrocrit (PCV)	42.8	%	40 - 54			
MCH	28.5	Pg	27 - 32			
MCV	85.3	fL	83 - 101			
MCHC	33.4	%	31.5 - 34.5			
RDW	12.4	%	11.5 - 14.5			
WBC Count	6010	/cmm	4000 - 11000			
DIFFERENTIAL WBC COUNT (Flow	<u>cytometry)</u>					
Neutrophils (%)	56	%	38 - 70			
Lymphocytes (%)	33	%	20 - 40			
Monocytes (%)	07	%	2 - 8			
Eosinophils (%)	04	%	0 - 6			
Basophils (%)	0	%	0 - 2			
Neutrophils	3366	/cmm				
Lymphocytes	1983	/cmm				
Monocytes	421	/cmm				
Eosinophils	240	/cmm				
Basophils	0	/cmm				
Platelet Count (Flow cytometry)	285000	/cmm	150000 - 450000			
MPV	8.6	fL	7.5 - 11.5			
ERYTHROCYTE SEDIMENTATION F	RATE					
ESR (After 1 hour)	09	mm/hr	0 - 14			
Modified Westergren Method						

----- End Of Report ------

Page 1 of 11

This is an electronically authenticated report.



		TEST REP	PORT
Reg. No	: 2307101704		Reg. Date : 22-Jul-2023
Name	: Dilpreet Singh		Collected On : 22-Jul-2023 08:47
Age/Sex	: 34 Years / Male		Approved On : 22-Jul-2023 10:55
Ref. By	:		Printed On : 25-Jul-2023 12:22
Client	: MEDIWHEEL WEL	LNESS	
Paramet	ter	<u>Result</u>	
		BLOOD GROU Specimen: EDTA and Serum; Me	
ABO		'O'	
Rh (D)		Positive	

----- End Of Report ------

This is an electronically authenticated report.

Page 2 of 11

Approved by: DR F

DR PS RAO MD Pathologist

	TEST	REPORT	
Reg. No : 2307101704			Reg. Date : 22-Jul-2023
Name : Dilpreet Singh			Collected On : 22-Jul-2023 08:4
Age/Sex : 34 Years / Male			Approved On : 22-Jul-2023 11:0
Ref. By			Printed On : 25-Jul-2023 12:2
Client : MEDIWHEEL WELLNESS			
Parameter_	<u>Result</u>	<u>Unit</u>	Reference Interval
Fasting Blood Sugar (FBS)	77.0	mg/dL	70 - 110
		A GLUCOSE	
Hexokinase Method	11.0	ing/aL	70 110
Post Prandial Blood Sugar (PPBS) Hexokinase Method	114.0	mg/dL	70 - 140
Criteria for the diagnosis of diabetes1. HbA1c >/=	6.5 *		
Or 2. Fasting plasma glucose >126 gm/dL. Fasting is de	fined as no caloric intal	ke at least for 8 hrs.	
Or			

We note plasma glucose >/= 200 mg/dL during an oral glucose tolerence test by using a glucose load containing equivity dissolved in water.
 Or
 In a patient with classic symptoms of hyperglycemia or hyperglycemic crisis, a random plasma glucose >/= 200 mg/dL.
 \*In the absence of unequivocal hyperglycemia, criteria 1-3 should be confirmed by repeat testing.
 American diabetes association. Standards of medical care in diabetes 2011. Diabetes care 2011;34;S11.

----- End Of Report ------

Page 3 of 11

# 

**TEST REPORT** 

Reg. No:2307101704Name:Dilpreet SinghAge/Sex:34 Years / MaleRef. By:

 Reg. Date
 :
 22-Jul-2023

 Collected On
 :
 22-Jul-2023 08:47

 Approved On
 :
 22-Jul-2023 11:09

 Printed On
 :
 25-Jul-2023 12:22

Client : MEDIWHEEL WELLNESS

Printed On	: 25-Jul-202

<u>Parameter</u>	<u>Result</u>	<u>Unit</u>	Reference Interval
	L	IPID PROFILE	
Cholesterol (Enzymatic colorimetric)	149.0	mg/dL	Desirable : < 200.0 Borderline High : 200-239 High : > 240.0
<b>Triglyceride</b> (Enzymatic colorimetric)	87.8	mg/dL	Normal : < 150.0 Borderline : 150-199 High : 200-499 Very High : > 500.0
VLDL	17.56	mg/dL	15 - 35
Calculated			
LDL CHOLESTEROL	94.44	mg/dL	Optimal : < 100.0 Near / above optimal : 100-129 Borderline High : 130-159 High : 160-189 Very High : >190.0
HDL Cholesterol	37.0	mg/dL	30 - 70
Homogeneous enzymatic colorimetr	ic		
Cholesterol /HDL Ratio	4.03		0 - 5.0
LDL / HDL RATIO Calculated	2.55		0 - 3.5

Page 4 of 11



<b>Reg. No</b> : 2307101704		Reg. Date : 22-Jul-2023
Name : Dilpreet Singh		Collected On : 22-Jul-2023 08:47
Age/Sex : 34 Years / Male		Approved On : 22-Jul-2023 11:09
Ref. By		Printed On : 25-Jul-2023 12:22
Client : MEDIWHEEL WELLNESS		
Parameter Resu	ult <u>Unit</u>	Reference Interval

NEW ATP III GUIDELINES (MAY 2001), MODIFICATION OF NCEP<?xml:namespace prefix = "o" ns = "urn:schemasmicrosoft-com:office:office" />

LDL CHOLESTEROL CHOLESTEROL HDL CHOLESTEROL
TRIGLYCERIDES
Optimal<100
Desirable<200
Low<40
Normal<150
Near Optimal 100-129
Border Line 200-239
High >60
Border High 150-199
Borderline 130-159
High >240
-
High 200-499
High 160-189

• LDL Cholesterol level is primary goal for treatment and varies with risk category and assessment

For LDL Cholesterol level Please consider direct LDL value •

Risk assessment from HDL and Triglyceride has been revised. Also LDL goals have changed.

Detail test interpreation available from the lab

All tests are done according to NCEP guidelines and with FDA approved kits. •

• LDL Cholesterol level is primary goal for treatment and varies with risk category and assessment # For test performed on specimens received or collected from non-KSHIPRA locations, it is presumed that the specimen belongs to the patient named or identified as labeled on the container/test request and such verification has been carried out at the point generation of the said specimen by the sender.

KSHIPRA will be responsible Only for the analytical part of test carried out. All other responsibility will be of referring Laboratory. . All other responsibility will be of referring Laboratory.

----- End Of Report ------

Page 5 of 11



		TEST REPORT	
Reg. No : 2307101704 Name : Dilpreet Singh			Reg. Date         :         22-Jul-2023           Collected On         :         22-Jul-2023 08:47
Age/Sex : 34 Years / Male Ref. By : Client : MEDIWHEEL WELLNES	6		Approved On         :         22-Jul-2023 11:09           Printed On         :         25-Jul-2023 12:22
Parameter	<u>Result</u>	<u>Unit</u>	Reference Interval
	LIVER FU	INCTION TEST WIT	H GGT
Total Bilirubin Colorimetric diazo method	0.85	mg/dL	0.10 - 1.0
Conjugated Bilirubin Sulph acid dpl/caff-benz	0.27	mg/dL	0.0 - 0.3
Unconjugated Bilirubin Sulph acid dpl/caff-benz	0.58	mg/dL	0.0 - 1.1
SGOT (Enzymatic)	34.1	U/L	0 - 37
SGPT (Enzymatic)	39.7	U/L	0 - 40
GGT (Enzymatic colorimetric)	29.4	U/L	11 - 49
Alakaline Phosphatase (Colorimetric standardized method)	99.1	U/L	53 - 130
<u>Protien with ratio</u> Total Protein	7.1	g/dL	6.5 - 8.7
(Colorimetric standardized method) Albumin	4.6	mg/dL	3.5 - 5.3
(Colorimetric standardized method) Globulin Calculated	2.50	g/dL	2.3 - 3.5

----- End Of Report ------

1.84

0.8 - 2.0

Page 6 of 11

A/G Ratio

Calculated



	TEST	REPORT	
Reg. No : 2307101704			Reg. Date : 22-Jul-2023
Name : Dilpreet Singh			Collected On : 22-Jul-2023 08:47
Age/Sex : 34 Years / Male			Approved On : 22-Jul-2023 11:09
Ref. By			Printed On : 25-Jul-2023 12:22
Client : MEDIWHEEL WELLNESS			
Parameter	<b>Result</b>	<u>Unit</u>	Reference Interval
	KIDNEY FL	<b>JNCTION TEST</b>	
	KIDNEY FI	INCTION TEST	
	KIDNEY FL14.8	JNCTION TEST mg/dL	10 - 50
			10 - 50
UREA (Urease & glutamate dehydrogenase) Creatinine (Jaffe method)			10 - 50 0.5 - 1.4

----- End Of Report -----

Page 7 of 11

Approved by: DR P

DR PS RAO MD Pathologist

This is an electronically authenticated report.

	TEST R	EPORT	
Reg. No : 2307101704			Reg. Date : 22-Jul-2023
lame : Dilpreet Singh			Collected On : 22-Jul-2023 08:47
Age/Sex : 34 Years / Male			Approved On : 22-Jul-2023 10:56
Ref. By : Client : MEDIWHEEL WELLNES			Printed On : 25-Jul-2023 12:22
Parameter	<u>Result</u>	Unit	Reference Interval
	URINE ROUTIN	E EXAMINATIO	JN .
<u>PHYSICAL EXAMINATION</u> Quantity	20 cc		
Colour	Pale Yellow		
Appearance	Clear		
CHEMICAL EXAMINATION ( BY RE	FLECTANCE PHOTOMET	TRIC METHOD)	
pH	6.0	-	5.0 - 8.0
Sp. Gravity	1.015		1.002 - 1.03
Protein	Nil		
Glucose	Nil		
Ketone Bodies	Nil		
Urine Bile salt and Bile Pigment	Nil		
Urine Bilirubin	Nil		
Nitrite	Nil		
Leucocytes	Nil		
Blood	Nil		
MICROSCOPIC EXAMINATION (MA	NUAL BY MCIROSCOPY	)	
Leucocytes (Pus Cells)	Nil		
Erythrocytes (Red Cells)	Nil		
Epithelial Cells	1-2/hpf		
Amorphous Material	Nil		
Casts	Nil		
Crystals	Nil		
Bacteria	Nil		
Monilia	Nil		

----- End Of Report ------

Page 8 of 11



	TE	ST REPORT	
Reg. No         : 2307101704           Name         : Dilpreet Singh			Reg. Date         : 22-Jul-2023           Collected On         : 22-Jul-2023 08:47
Age/Sex : 34 Years / Male Ref. By : Client : MEDIWHEEL WELLN	ESS		Approved On         :         25-Jul-2023 09:59           Printed On         :         25-Jul-2023 12:22
Parameter	<u>Result</u>	Unit	Reference Interval
	STOO	L EXAMINATIO	Ν
Colour	Yellow		
Consistency	Solid		
CHEMICAL EXAMINATION			
Occult Blood	Negative		
Peroxidase Reaction with o- Dianisidine			
Reaction	Acidic		
pH Strip Method			
Reducing Substance	Absent		
Benedict's Method			
MICROSCOPIC EXAMINATION Mucus	Nil		
Pus Cells	1 - 2/hpf		
Red Cells	Nil		
Epithelial Cells	Nil		
Vegetable Cells	Nil		
Trophozoites	Nil		
Cysts	Nil		
Ova	Nil		
Neutral Fat	Nil		
Monilia	Nil		
OTHER	-		

Note: Stool occult blood test is highly sensitive to peroxidase like activity of free hemoglobin.

False negative: False negative occult blood test may be observed in case of excess (>250mg/day) Vitamin C intake and in case of occassinal unruptured RBCs.

**False positive:** False positive occult blood test may be observed in stool samples containing vegetable peroxidase (turnips, horseradish, cauliflower, brocoli, cantaloupe, parsnips) and myoglobin from food (meat diet) intake.

----- End Of Report ------

Page 9 of 11



Client Paramet	: MEDIWHEEL WELLNESS	<u>Result</u>	<u>Unit</u>	Reference Interval	
Ref. By				Printed On : 25-Jul-2023 12:22	
Age/Sex	: 34 Years / Male			Approved On : 22-Jul-2023 11:09	
Reg. No Name	: 2307101704 : Dilpreet Singh			Reg. Date         :         22-Jul-2023           Collected On         :         22-Jul-2023 08:47	
		TEST	REPORT		

Hb A1C Boronate Affinity with Fluorescent Quenching	5.8	% of Total Hb	Poor Control : > 7.0 % Good Control : 6.2-7.0 % Non-diabetic Level : 4.3-6.2 %
Mean Blood Glucose	129.18	mg/dL	

#### Degree of Glucose Control Normal Range:

Poor Control >7.0% \*

Good Control 6.0 - 7.0 %\*\*Non-diabetic level < 6.0 %

\* High risk of developing long term complication such as retinopathy, nephropathy, neuropathy, cardiopathy,etc.

\* Some danger of hypoglycemic reaction in Type I diabetics.

\* Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1c levels in this area.

#### EXPLANATION :-

\*Total haemoglobin A1 c is continuously symthesised in the red blood cell throught its 120 days life span. The concentration of HBA1c in the cell reflects the average blood glucose concentration it encounters.

\*The level of HBA1c increases proportionately in patients with uncontrolled diabetes. It reflects the average blood glucose oncentration over an extended time period and remains unaffected by short-term fluctuations in blood glucose levels. \*The measurement of HbA1c can serve as a convenient test for evaluating the adequacy of diabetic control and in preventing various diabetic complications. Because the average half life of a red blood cell is sixty days,HbA1c has been accepted as a measurnment which eflects the mean daily blood glucose concentration, better than fasting blood glucose determination, and the degree of carbohydrate imbalance over the preceding two months.

\*It may also provide a better index of control of the diabetic patient without resorting to glucose loading procedures.

#### HbA1c assay Interferences:

\*Errneous values might be obtained from samples with abnormally elevated quantities of other Haemoglobins as a result of either their simultaneous elution with HbA1c(HbF) or differences in their glycation from that of HbA(HbS)

----- End Of Report ------

This is an electronically authenticated report.

Page 10 of 11



DR PS RAO MD Pathologist

	Т	EST REPORT	
Reg. No : 2307101704			Reg. Date : 22-Jul-2023
Name : Dilpreet Singh			Collected On : 22-Jul-2023 08:47
Age/Sex : 34 Years / Male			Approved On : 22-Jul-2023 11:10
Ref. By : Client : MEDIWHEEL WEL	LNESS		Printed On : 25-Jul-2023 12:22
Parameter	<u>Result</u>	<u>Unit</u>	Reference Interval
	THYRC	DID FUNCTION T	EST
T3 (Triiodothyronine) Chemiluminescence	1.24	ng/mL	0.87 - 1.81
T4 (Thyroxine) Chemiluminescence	10.11	μg/dL	5.89 - 14.9
TSH ( ultra sensitive )	1.789	μIU/ml	0.34 - 5.6

Chemiluminescence

SUMMARY The hypophyseal release of TSH (thyrotropic hormone) is the central regulating mechanism for the biological action of thyroid hormones.TSH is a very sensitive and specific parameter for assessing thyroid function and is particularly suitable for early detection or exclusion of disorders in the central regulating circuit between the hypothalamus, pituitary and thyroid. LIMITATION Presence of autoantibodies may cause unexpected high value of TSH

----- End Of Report ------

Page 11 of 11

This is an electronically authenticated report.