

# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road,  
Sodala, Jaipur-302019

Tele : 0141-2293346, 4049787, 9887049787

### General Physical Examination

Website: www.drgoyaispathlab.com | E-mail: drgoyalpiyush@gmail.com

Date of Examination: 24-02-2024

Name: NARENDRA SINGH Age: 46 Sex: Male

DOB: 14-07-1977

Referred By: BOB (Mediwheel)

Photo ID: Wadhav ID #: attached

Ht: 165 (cm)

Wt: 66 (Kg)

Chest (Expiration): 92 (cm)

Abdomen Circumference: 90 (cm)

Blood Pressure: 122/73 mm Hg PR: 73 / min

BMI 24.2

Eye Examination: Dist Vision R.E. 6/6. L.E. 6/9 (with specs)

Near vision N/G Bil eyes. (with specs).

Other: Not significant

On examination he/she appears physically and mentally fit:  Yes / No

Signature Of Examinee : [Signature] Name of Examinee: \_\_\_\_\_

Signature Medical Examiner: [Signature] Name Medical Examiner \_\_\_\_\_

**Piyush Goyal**  
M.B.B.S., D.M.R.D.  
RMC Reg. No.-017996

भारत सरकार  
GOVERNMENT OF INDIA

भारत सरकार  
GOVERNMENT OF INDIA

Download Date: 24/12/2020

Issue Date: 06/07/2017


 नरेंद्र सिंह  
 Narendra Singh  
 जन्म तिथि/DOB: 14/07/1977  
 लिंग/ GENDER: MALE

9794 0973 0138

VID : 9170 9250 9092 5841

मेरा आधार, मेरी पहचान

DR

भारत सरकार  
GOVERNMENT OF INDIA

भारत सरकार  
GOVERNMENT OF INDIA

पता:  
 पद्मा सिंघ, 35, शिवपुरी योजना, रंकरि सोदला के पास,  
 रंकरि सोदला, जयपुर, राजस्थान - 302006

Address:  
 S/O. Bhawani Singh, 35, Shivpuri Yojana,  
 Near Nvn School, Rankari Sodala, Jaipur,  
 Rajasthan - 302006



9794 0973 0138

VID : 9170 9250 9092 5841


1947 | [help@uidai.gov.in](mailto:help@uidai.gov.in) | [www.uidai.gov.in](http://www.uidai.gov.in)

Dr. Piyush Goyal  
 H.B.S., D.M.R.D.  
 Rm. No. 10, 317236

DR. GOYAL PATH LAB

EKG

4397 / MR NARENDRA SINGH / 46 Yrs / M / Non Smoker  
Heart Rate : 70 bpm / Tested On : 24-Feb-24 11:15:16 / HF 0.05 Hz - LF 35 Hz / Notch 50 Hz / Sn 1.00 Cm/mV / Sw 25 mm/s  
/ Reled By: BOB



Vent Rate : 70 bpm  
PR Interval : 194 ms  
QRS Duration: 84 ms  
QT/QTc Int : 354/372 ms  
P-QRS-T axis: 50.00 • 18.00 • 12.00 •



*Narender Kumar Mishra*  
RMC No. 52703  
Miss. Dip. CARDIO (ESCORT)  
T. 22.00 (RCGP 50'00")  
R 18.00

Reported By:

*Praveen*

# Dr. Goyal's

## Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganeer Road, 5509  
Sodala, Jaipur-302019  
Tele : 0141-2293346, 4049787, 9887049787  
Website: www.drgoyalpathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09  
**NAME :- Mr. NARENDRA SINGH**  
Sex / Age :- Male 46 Yrs 7 Mon 13 Days  
Company :- MediWheel

Patient ID :-12235997  
Ref. By Dr:- BOB  
Lab/Hosp :-



Sample Type :- EDTA

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 13:24:04

### HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
-----------	-------	------	-------------------------

#### BOB PACKAGE ABOVE 40MALE

GLYCOSYLATED HEMOGLOBIN (HbA1C)  
Method:- HPLC

6.2 H %

Non-diabetic: < 5.7  
Pre-diabetics: 5.7-6.4  
Diabetics: = 6.5 or higher  
ADA Target: 7.0  
Action suggested: > 6.5

Instrument name: ARKRAY's ADAMS Lite HA 8380V, JAPAN.

#### Test Interpretation:

HbA1C is formed by the condensation of glucose with n-terminal valine residue of each beta chain of HbA to form an unstable schiff base. It is the major fraction, constituting approximately 80% of HbA1c. Formation of glycated hemoglobin (GHb) is essentially irreversible and the concentration in the blood depends on both the lifespan of the red blood cells (RBC) (120 days) and the blood glucose concentration. The GHb concentration represents the integrated values for glucose over the period of 6 to 8 weeks. GHb values are free of day to day glucose fluctuations and are unaffected by recent exercise or food ingestion. Concentration of plasma glucose concentration in GHb depends on the time interval, with more recent values providing a larger contribution than earlier values. The interpretation of GHb depends on RBC having a normal life span. Patients with hemolytic disease or other conditions with shortened RBC survival exhibit a substantial reduction of GHb. High GHb have been reported in iron deficiency anemia. GHb has been firmly established as an index of long term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. The absolute risk of retinopathy and nephropathy are directly proportional to the mean of HbA1C. Genetic variants (e.g. HbS trait, HbC trait), elevated HbF and chemically modified derivatives of hemoglobin can affect the accuracy of HbA1c measurements. The effects vary depending on the specific Hb variant or derivative and the specific HbA1c method.

Ref by ADA 2020

MEAN PLASMA GLUCOSE  
Method:- Calculated Parameter

131 H mg/dL

Non Diabetic < 100 mg/dL  
Prediabetic 100- 125 mg/dL  
Diabetic 126 mg/dL or Higher

MUKESH SINGH  
Technologist

Page No: 1 of 13



**Dr. Rashmi Bakshi**  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganeer Road, 5509  
Sodala, Jaipur-302019  
Tele : 0141-2293346, 4049787, 9887049787  
Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09  
**NAME :- Mr. NARENDRA SINGH**  
Sex / Age :- Male 46 Yrs 7 Mon 13 Days  
Company :- MediWheel

Patient ID :- 12235997  
Ref. By Dr:- BOB  
Lab/Hosp :-



Sample Type :- EDTA

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 13:24:04

### HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
<b>HAEMOGARAM</b>			
HAEMOGLOBIN (Hb)	12.8 L	g/dL	13.0 - 17.0
TOTAL LEUCOCYTE COUNT	4.33	/cumm	4.00 - 10.00
<b>DIFFERENTIAL LEUCOCYTE COUNT</b>			
NEUTROPHIL	46.4	%	40.0 - 80.0
LYMPHOCYTE	46.8 H	%	20.0 - 40.0
EOSINOPHIL	3.1	%	1.0 - 6.0
MONOCYTE	3.5	%	2.0 - 10.0
BASOPHIL	0.2	%	0.0 - 2.0
NEUT#	2.01	10 <sup>3</sup> /uL	1.50 - 7.00
LYMPH#	2.03	10 <sup>3</sup> /uL	1.00 - 3.70
EO#	0.13	10 <sup>3</sup> /uL	0.00 - 0.40
MONO#	0.15	10 <sup>3</sup> /uL	0.00 - 0.70
BASO#	0.01	10 <sup>3</sup> /uL	0.00 - 0.10
TOTAL RED BLOOD CELL COUNT (RBC)	5.36	x10 <sup>6</sup> /uL	4.50 - 5.50
HEMATOCRIT (HCT)	42.30	%	40.00 - 50.00
MEAN CORP VOLUME (MCV)	78.9 L	fL	83.0 - 101.0
MEAN CORP HB (MCH)	23.9 L	pg	27.0 - 32.0
MEAN CORP HB CONC (MCHC)	30.3 L	g/dL	31.5 - 34.5
<b>PLATELET COUNT</b>			
RDW-CV	16.0 H	%	11.6 - 14.0
MENTZER INDEX	14.72		

The Mentzer index is used to differentiate iron deficiency anemia from beta thalassemia trait. If a CBC indicates microcytic anemia, these are two of the most likely causes, making it necessary to distinguish between them.

If the quotient of the mean corpuscular volume divided by the red blood cell count is less than 13, thalassemia is more likely. If the result is greater than 13, then iron-deficiency anemia is more likely.

MUKESH SINGH  
Technologist

Page No: 2 of 13



Dr. Rashmi Bakshi  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road,  
Sodala, Jaipur-302019  
Tele : 0141-2293346, 4049787, 9887049787  
Website: www.drgoyalpathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09

Patient ID :-12235997



NAME :- Mr. NARENDRA SINGH

Ref. By Dr- BOB

Sex / Age :- Male 46 Yrs 7 Mon 13 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- EDTA

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 13:24:04

### HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
Erythrocyte Sedimentation Rate (ESR)	35 H	mm/hr.	00 - 13

(ESR) Methodology : Measurement of ESR by cells aggregation.

Instrument Name : Independent form Hematocrit value by Automated Analyzer (Roller-20)

Interpretation : ESR test is a non-specific indicator of inflammatory disease and abnormal protein states.

The test is used to detect, follow course of a certain disease (e.g-tuberculosis, rheumatic fever, myocardial infarction

Levels are higher in pregnancy due to hyperfibrinogenaemia.

The "3-figure ESR" >100 value nearly always indicates serious disease such as a serious infection, malignant paraproteinaemia or connective tissue disease.

(CBC), Methodology: TLC, DLC, Fluorescent Flow cytometry, HB SLS method, TRBC, PCV, PLT Hydrodynamically focused Impedance, and MCH, MCV, MCHC, MENTZER INDEX are calculated. Instrument Name: Sysmex 6 part fully automatic analyzer XN-L, Japan

MUKESH SINGH  
Technologist

Page No: 3 of 13



Dr. Rashmi Bakshi  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganeer Road, 5509  
 Sodala, Jaipur-302019  
 Tele : 0141-2293346, 4049787, 9887049787  
 Website: www.drgoyalpathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09 Patient ID :-12235997  
**NAME :- Mr. NARENDRA SINGH** Ref. By Dr:- BOB  
 Sex / Age :- Male 46 Yrs 7 Mon 13 Days Lab/Hosp :-  
 Company :- MediWheel



Sample Type :- PLAIN/SERUM Sample Collected Time 24/02/2024 09:21:51 Final Authentication : 24/02/2024 13:20:55

### BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
<b>LIPID PROFILE</b>			
TOTAL CHOLESTEROL Method:- Enzymatic Endpoint Method	190.65	mg/dl	Desirable <200 Borderline 200-239 High > 240
TRIGLYCERIDES Method:- GPO-PAP	141.40	mg/dl	Normal <150 Borderline high 150-199 High 200-499 Very high >500
DIRECT HDL CHOLESTEROL Method:- Direct clearance Method	38.90	mg/dl	Low < 40 High > 60
DIRECT LDL CHOLESTEROL Method:- Direct clearance Method	128.18	mg/dl	Optimal <100 Near Optimal/above optimal 100-129 Borderline High 130-159 High 160-189 Very High > 190
VLDL CHOLESTEROL Method:- Calculated	28.28	mg/dl	0.00 - 80.00
T.CHOLESTEROL/HDL CHOLESTEROL RATIO Method:- Calculated	4.90		0.00 - 4.90
LDL / HDL CHOLESTEROL RATIO Method:- Calculated	3.30		0.00 - 3.50
TOTAL LIPID Method:- CALCULATED	591.62	mg/dl	400.00 - 1000.00
<small>TOTAL CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: Cholesterol measurements are used in the diagnosis and treatment of lipid lipoprotein metabolism disorders.</small>			
<small>TRIGLYCERIDES InstrumentName:Randox Rx Imola Interpretation: Triglyceride measurements are used in the diagnosis and treatment of diseases involving lipid metabolism and various endocrine disorders e.g. diabetes mellitus, nephrosis and liver obstruction.</small>			
<small>DIRECT HDL CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: An inverse relationship between HDL-cholesterol (HDL-C) levels in serum and the incidence/prevalence of coronary heart disease (CHD) has been demonstrated in a number of epidemiological studies. Accurate measurement of HDL-C is of vital importance when assessing patient risk from CHD. Direct measurement gives improved accuracy and reproducibility when compared to precipitation methods.</small>			
<small>DIRECT LDL-CHOLESTEROL InstrumentName:Randox Rx Imola Interpretation: Accurate measurement of LDL-Cholesterol is of vital importance in therapies which focus on lipid reduction to prevent atherosclerosis or reduce its progress and to avoid plaque rupture.</small>			
<small>TOTAL LIPID AND VLDL ARE CALCULATED</small>			

SURENDRAKHANGA

Page No: 4 of 13



**Dr. Rashmi Bakshi**  
 MBBS, MD ( Path )  
 RMC No. 17975/008828



Date :- 24/02/2024 09:19:09

Patient ID :- 12235997



**NAME :- Mr. NARENDRA SINGH**

Ref. By Dr.- BOB

Sex / Age :- Male 46 Yrs 7 Mon 13 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 13:20:55

### BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
<b>LIVER PROFILE WITH GGT</b>			
SERUM BILIRUBIN (TOTAL) Method:- Colorimetric method	0.67	mg/dl	Up to - 1.0 Cord blood <2 Premature < 6 days <16 Full-term < 6 days= 12 1month - <12 months <2 1-19 years <1.5 Adult - Up to - 1.2 Ref-(ACCP 2020)
SERUM BILIRUBIN (DIRECT) Method:- Colorimetric Method	0.17	mg/dL	Adult - Up to 0.25 Newborn - <0.6 >- 1 month - <0.2
SERUM BILIRUBIN (INDIRECT) Method:- Calculated	0.50	mg/dl	0.30-0.70
SGOT Method:- IFCC	34.8	U/L	Men- Up to - 37.0 Women - Up to - 31.0
SGPT Method:- IFCC	39.9	U/L	Men- Up to - 40.0 Women - Up to - 31.0
SERUM ALKALINE PHOSPHATASE Method:- AMP Buffer	56.60	IU/L	30.00 - 120.00
SERUM GAMMA GT Method:- IFCC	19.90	U/L	11.00 - 50.00
SERUM TOTAL PROTEIN Method:- Buret Reagent	* 7.44	g/dl	6.40 - 8.30
SERUM ALBUMIN Method:- Bromocresol Green	4.62	g/dl	3.80 - 5.00
SERUM GLOBULIN Method:- CALCULATION	2.82	gm/dl	2.20 - 3.50
A/G RATIO	1.64		1.30 - 2.50

**Total Bilirubin** Methodology: Colorimetric method Instrument Name Randox Rx Inits Interpretation: An increase in bilirubin concentration in the serum occurs in toxic or infectious diseases of the liver e.g. hepatitis B or obstruction of the bile duct and in chronic incompatible babies. High levels of unconjugated bilirubin indicate that too much haemoglobin is being destroyed or that the liver is not actively treating the haemoglobin it is receiving.

**AST Aspartate Aminotransferase** Methodology: IFCC Instrument Name Randox Rx Inits Interpretation: Elevated levels of AST can signal myocardial infarction, hepatic disease, muscular dystrophy and organ damage. Although heart muscle is found to have the most activity of the enzyme, significant activity has also been seen in the brain, liver, gastric mucosa, adipose tissue and kidneys of humans.

**ALT Alanine Aminotransferase** Methodology: IFCC Instrument Name Randox Rx Inits Interpretation: The enzyme ALT has been found to be in highest concentrations in the liver, with decreasing concentrations found in kidney, heart, skeletal muscle, pancreas, spleen and lung tissue respectively. Elevated levels of the transaminase can indicate myocardial infarction, hepatic disease, muscular dystrophy and organ damage.

**Alkaline Phosphatase** Methodology: AMP Buffer Instrument Name Randox Rx Inits Interpretation: Measurements of alkaline phosphatase are of use in the diagnosis, treatment and investigation of hepatobiliary disease and in bone disease associated with increased osteoblastic activity. Alkaline phosphatase is also used in the diagnosis of parathyroid and renal disease.

**TOTAL PROTEIN** Methodology: Buret Reagent Instrument Name Randox Rx Inits Interpretation: Measurements obtained by this method are used in the diagnosis and treatment of a variety of diseases involving the liver, kidney and bone marrow as well as other metabolic or nutritional disorders.

**ALBUMIN (ALB)** Methodology: Bromocresol Green Instrument Name Randox Rx Inits Interpretation: Albumin measurements are used in the diagnosis and treatment of numerous diseases involving primarily the liver or kidneys. Globulin & A/G ratio is calculated.

**Instrument Name Randox Rx Inits Interpretation:** Elevations in GGT levels are seen earlier and more pronounced than those with other liver enzymes in cases of obstructive jaundice and metastatic neoplasms. It may reach 5 to 30 times normal levels in intra- or post-hepatic biliary obstruction. Only moderate elevations in the enzyme level (2 to 5 times normal).

SURENDRAKHANGA

**Dr. Rashmi Bakshi**  
MBBS, MD ( Path )  
RMC No. 17975/008828





# Dr. Goyal's

## Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sangarner Road, 5509  
Sodala, Jaipur-302019  
Tele : 0141-2293346, 4049787, 9887049787  
Website: www.drgoyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09

Patient ID :- 12235997

NAME :- Mr. NARENDRA SINGH

Ref. By Dr:- BOB

Sex / Age :- Male 46 Yrs 7 Mon 13 Days

Lab/Hosp :-

Company :- MediWheel



Sample Type :- PLAIN/SERUM

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 11:11:46

### IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
<b>TOTAL THYROID PROFILE</b>			
SERUM TOTAL T3 Method:- Chemiluminescence(Competitive immunoassay)	1.050	ng/ml	0.970 - 1.690
SERUM TOTAL T4 Method:- Chemiluminescence(Competitive immunoassay)	6.940	ug/dl	5.530 - 11.000
SERUM TSH ULTRA Method:- Enhanced Chemiluminescence Immunoassay	1.468	μIU/mL	0.350 - 5.500

**Interpretation:** Triiodothyronine (T3) contributes to the maintenance of the euthyroid state. A decrease in T3 concentration of up to 50% occurs in a variety of clinical situations, including acute and chronic disease. Although T3 results alone cannot be used to diagnose hypothyroidism, T3 concentration may be more sensitive than thyroxine (T4) for hyperthyroidism. Consequently, the total T3 assay can be used in conjunction with other assays to aid in the differential diagnosis of thyroid disease. T3 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, Free T3 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake, or T4 uptake can be used with the total T3 result to calculate the free T3 index and estimate the concentration of free T3.

**Interpretation:** The measurement of Total T4 aids in the differential diagnosis of thyroid disease. While >99.9% of T4 is protein-bound, primarily to thyroxine-binding globulin (TBG), it is the free fraction that is biologically active. In most patients, the total T4 concentration is a good indicator of thyroid status. T4 concentrations may be altered in some conditions, such as pregnancy, that affect the capacity of the thyroid hormone-binding proteins. Under such conditions, free T4 can provide the best estimate of the metabolically active hormone concentration. Alternatively, T3 uptake may be used with the total T4 result to calculate the free T4 index (FT4I) and estimate the concentration of free T4. Some drugs and some nonthyroidal patient conditions are known to alter TT4 concentrations in vivo.

**Interpretation:** TSH stimulates the production of thyroxine (T4) and triiodothyronine (T3) by the thyroid gland. The diagnosis of overt hypothyroidism by the finding of a low total T4 or free T4 concentration is readily confirmed by a raised TSH concentration. Measurement of low or undetectable TSH concentrations may assist the diagnosis of hyperthyroidism, where concentrations of T4 and T3 are elevated and TSH secretion is suppressed. These have the advantage of discriminating between the concentrations of TSH observed in thyrotoxicosis, compared with the low, but detectable, concentrations that occur in subclinical hyperthyroidism. The performance of this assay has not been established for neonatal specimens. Some drugs and some nonthyroidal patient conditions are known to alter TSH concentrations in vivo.

### INTERPRETATION

PREGNANCY	REFERENCE RANGE FOR TSH IN uIU/mL (As per American Thyroid Association)
1st Trimester	0.10-2.50
2nd Trimester	0.20-3.00
3rd Trimester	0.30-3.00

NARENDRAKUMAR  
Technologist

Page No: 6 of 13



Dr. Rashmi Bakshi  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road,  
Sodala, Jaipur-302019  
Tele : 0141-2293346, 4049787, 9887049787  
Website: www.drgoyalpathlab.com | E-mail: drgoyalpiyush@gmail.com

MC- 5509

Date :- 24/02/2024 09:19:09 Patient ID :-12235997  
**NAME :- Mr. NARENDRA SINGH** Ref. By Dr.- BOB  
 Sex / Age :- Male 46 Yrs 7 Mon 13 Days Lab/Hosp :-  
 Company :- MediWheel



Sample Type :- URINE Sample Collected Time 24/02/2024 09:21:51 Final Authentication : 24/02/2024 10:18:21

### CLINICAL PATHOLOGY

Test Name	Value	Unit	Biological Ref Interval
<b>Urine Routine</b>			
<b><u>PHYSICAL EXAMINATION</u></b>			
COLOUR	PALE YELLOW		PALE YELLOW
APPEARANCE	Clear		Clear
<b><u>CHEMICAL EXAMINATION</u></b>			
REACTION(PH) Method:- Reagent Strip(Double indication blue reaction)	6.0		5.0 - 7.5
SPECIFIC GRAVITY Method:- Reagent Strip(bromothymol blue)	1.025		1.010 - 1.030
PROTEIN Method:- Reagent Strip (Sulphosalicylic acid test)	NIL		NIL
GLUCOSE Method:- Reagent Strip (Gls.Oxidase Peroxidase Benedict)	NIL		NIL
BILIRUBIN Method:- Reagent Strip (Azo-coupling reaction)	NEGATIVE		NEGATIVE
UROBILINOGEN Method:- Reagent Strip (Modified ehrlich reaction)	NORMAL		NORMAL
KETONES Method:- Reagent Strip (Sodium Nitroprusside) Rothera's	NEGATIVE		NEGATIVE
NITRITE Method:- Reagent Strip (Diazotization reaction)	NEGATIVE		NEGATIVE
RBC Method:- Reagent Strip (Peroxidase like activity)	NIL		NIL
<b><u>MICROSCOPY EXAMINATION</u></b>			
RBC/HPF	NIL	/HPF	NIL
WBC/HPF	2-3	/HPF	2-3
EPITHELIAL CELLS	2-3	/HPF	2-3
CRYSTALS/HPF	ABSENT		ABSENT
CAST/HPF	ABSENT		ABSENT
AMORPHOUS SEDIMENT	ABSENT		ABSENT
BACTERIAL FLORA	ABSENT		ABSENT
YEAST CELL	ABSENT		ABSENT
OTHER	ABSENT		ABSENT

VIJENDRAMEENA  
Technologist

Page No: 7 of 13



**Dr. Rashmi Bakshi**  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre



B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganeer Road, Jaipur-302019  
 Tele : 0141-2293346, 4049787, 9887049787  
 Website: www.dr-goyalspathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09 Patient ID :- 12235997  
**NAME :- Mr. NARENDRA SINGH** Ref. By Dr.- BOB  
 Sex / Age :- Male 46 Yrs 7 Mon 13 Days Lab/Hosp :-  
 Company :- MediWheel



Sample Type :- KOx/Na FLUORIDE-F, KOx/Na Fluoride-F, KOx/Na Fluoride-F Date: 24/02/2024 13:16:13 Final Authentication : 24/02/2024 14:20:55

### BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
FASTING BLOOD SUGAR (Plasma) Method:- GOD PAP	99.6	mg/dl	75.0 - 115.0
Impaired glucose tolerance (IGT)	111 - 125 mg/dL		
Diabetes Mellitus (DM)	> 126 mg/dL		
<b>Instrument Name:</b> Randox Rx Imola <b>Interpretation:</b> Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.			
BLOOD SUGAR PP (Plasma) Method:- GOD PAP	103.3	mg/dl	70.0 - 140.0
<b>Instrument Name:</b> Randox Rx Imola <b>Interpretation:</b> Elevated glucose levels (hyperglycemia) may occur with diabetes, pancreatic neoplasm, hyperthyroidism and adrenal cortical hyper-function as well as other disorders. Decreased glucose levels (hypoglycemia) may result from excessive insulin therapy or various liver diseases.			
SERUM CREATININE Method:- Colorimetric Method	0.89	mg/dl	Men - 0.6-1.30 Women - 0.5-1.20
SERUM URIC ACID Method:- Enzymatic colorimetric	5.28	mg/dl	Men - 3.4-7.0 Women - 2.4-5.7

SURENDRAKHANGA

Page No: 9 of 13



**Dr. Rashmi Bakshi**  
 MBBS. MD ( Path )  
 RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road,  
Sodala, Jaipur-302019  
Tele : 0141-2293346, 4049787, 9887049787  
Website: www.drgoyalpathlab.com | E-mail: drgoyalplyush@gmail.com

Date :- 24/02/2024 09:19:09 Patient ID :-12235997  
NAME :- Mr. NARENDRA SINGH Ref. By Dr:- BOB  
Sex / Age :- Male 46 Yrs 7 Mon 13 Days Lab/Hosp :-  
Company :- Medi/Wheel



Sample Type -> EDTA, URINE

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 13:24:04

### HAEMATOLOGY

Test Name	Value	Unit	Biological Ref Interval
BLOOD GROUP ABO	"B" POSITIVE		
BLOOD GROUP ABO Methodology : Haemagglutination reaction Kit Name : Monoclonal agglutinating antibodies (Span clone).			
URINE SUGAR (FASTING) Collected Sample Received	Nil		Nil

MUKESH SINGH, VIJENDRAMEENA  
Technologist

Page No: 11 of 13



**Dr. Rashmi Bakshi**  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road,  
Sodala, Jaipur-302019

Tele : 0141-2293346, 4049787, 9887049787

Website: www.drgoyalpathlab.com | E-mail: drgoyalpiyush@gmail.com

Date :- 24/02/2024 09:19:09

Patient ID :-12235997



NAME :- Mr. NARENDRA SINGH

Ref. By Dr:- BOB

Sex / Age :- Male 46 Yrs 7 Mon 13 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLAIN/SERUM

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 13:20:55

### BIOCHEMISTRY

Test Name	Value	Unit	Biological Ref Interval
BLOOD UREA NITROGEN (BUN)	8.6	mg/dl	0.0 - 23.0

SURENDRAKHANGA

Page No: 12 of 13



**Dr. Rashmi Bakshi**  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Piller No. 109-110, New Sanganer Road,  
Sodala, Jaipur-302019

Tele : 0141-2293346, 4049787, 9887049787

Website: www.drgoyalspathlab.com | E-mail: drgoyalplyush@gmail.com

Date :- 24/02/2024 09:19:09

Patient ID :-12235997



NAME :- Mr. NARENDRA SINGH

Ref. By Dr:- BOB

Sex / Age :- Male 46 Yrs 7 Mon 13 Days

Lab/Hosp :-

Company :- MediWheel

Sample Type :- PLA/IN/SERUM

Sample Collected Time 24/02/2024 09:21:51

Final Authentication : 24/02/2024 11:11:46

### IMMUNOASSAY

Test Name	Value	Unit	Biological Ref Interval
TOTAL PSA Method:- Chemiluminescence	1.960	ng/ml	0.000 - 4.000

**InstrumentName:** VITROS ECI **Interpretation:** Elevated serum PSA concentrations are found in men with prostate cancer, benign prostatic hypertrophy (BHP) or inflammatory conditions of other adjacent genitourinary tissues, but not in apparently healthy men or in men with cancers other than prostate cancer. PSA has been demonstrated to be an accurate marker for monitoring advancing clinical stage in untreated patients and for monitoring response to therapy by radical prostatectomy, radiation therapy and anti-androgen therapy. PSA is also important in determining the potential and actual effectiveness of surgery or other therapies. Progressive disease is defined by an increase of at least 25%. Sampling should be repeated within two to four weeks for additional evidence. Different assay methods cannot be used interchangeably.

\*\*\* End of Report \*\*\*

NARENDRAKUMAR  
Technologist

Page No: 13 of 13



Dr. Rashmi Bakshi  
MBBS, MD ( Path )  
RMC No. 17975/008828

# Dr. Goyal's

Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur  
Tele : 0141-2293346, 4049787, 9887049787  
Website : www.drgoyalpathlab.com E-mail : drgoyalpiyush@gmail.com



Date :- 24/02/2024 09:19:09  
**NAME :- Mr. NARENDRA SINGH**  
Sex / Age :- Male 46 Yrs 7 Mon 13 Days  
Company :- MediWheel

Patient ID :-12235997  
Ref. By Doctor:-BOB  
Lab/Hosp :-

Final Authentication : 24/02/2024 15:38:18

BOB PACKAGE ABOVE 40MALE

## X RAY CHEST PA VIEW:

Both lung fields appears clear.

Bronchovascular markings appear normal.

Trachea is in midline.

Both the hilar shadows are normal.

Both the C.P.angles is clear.

Both the domes of diaphragm are normally placed.

Bony cage and soft tissue shadows are normal.

Heart shadows appear normal.

**Impression :- Normal Study**

(Please correlate clinically and with relevant further investigations)



Dr. NAVNEET AGARWAL (MD, DNB RADIO-DIAGNOSIS, MNAMS)  
EX-SR NEURO-RADIOLOGY AIIMS NEW DELHI  
(RMC No. 33613 / 14911)

\*\*\* End of Report \*\*\*

**Dr. Piyush Goyal**  
(D.M.R.D.) BILAL  
Transcript by.

Page No: 1 of 1

Dr. Piyush Goyal  
M.B.B.S., D.M.R.D.  
RMC Reg No. 017998

Dr. Ashish Choudhary  
MBBS, MD (Radio-Diagnosis)  
Fetal Medicine Consultant  
FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain  
MBBS, DNB, (Radio-Diagnosis)  
RMC No. 21687

Dr. Navneet Agarwal  
MD, DNB (Radio Diagnosis)  
RMC No. 33613/14911

Dr. Poorvi Malik  
MBBS, MD, DNB (Radio Diagnosis)  
RMC No. 21505



Date :- 24/02/2024 09:19:09  
**NAME :- Mr. NARENDRA SINGH**  
Sex / Age :- Male 46 Yrs 7 Mon 13 Days  
Company :- MediWheel

Patient ID :- 12235997  
Ref. By Doctor:-BOB  
Lab/Hosp :-

Final Authentication : 24/02/2024 12:47:02

BOB PACKAGE ABOVE 40MALE

**USG WHOLE ABDOMEN**

**Liver** is of normal size. **Echo-texture is bright.** No focal space occupying lesion is seen within liver parenchyma. Intra hepatic biliary channels are not dilated. Portal vein diameter is normal.

**Gall bladder** is of normal size. Wall is not thickened. No calculus or mass lesion is seen in gall bladder. Common bile duct is not dilated.

**Pancreas** is of normal size and contour. Echo-pattern is normal. No focal lesion is seen within pancreas.

**Spleen** is of normal size and shape. Echotexture is normal. No focal lesion is seen.

**Kidneys** are normally sited and are of normal size and shape. Cortico-medullary echoes are normal. No focal lesion is seen. Collecting system does not show any dilatation or calculus.

**Urinary bladder** is partially distended.

**Prostate is enlarged in size (~ 27 gms)** with normal echo-texture and outline. No enlarged nodes are visualised. No retro-peritoneal lesion is identified. No significant free fluid is seen in peritoneal cavity.

**IMPRESSION:**

- \* Grade I fatty liver.
- \* Mild prostatomegaly.

Needs clinical correlation.

\*\*\* End of Report \*\*\*



# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur  
 Tele : 0141-2293346, 4049787, 9887049787  
 Website : www.drgoyalspathlab.com E-mail : drgoyalpiyush@gmail.com



Date :- 24/02/2024 09:19:09	Patient ID :- 12235997
NAME :- Mr. NARENDRA SINGH	Ref. By Doctor:-BOB
Sex / Age :- Male 46 Yrs 7 Mon 13 Days	Lab/Hosp :-
Company :- MediWheel	

Final Authentication : 24/02/2024 13:56:15

BOB PACKAGE ABOVE 40MALE  
 2D ECHO OPTION TMT (ADULT/CHILD)

### 2D-ECHOCARDIOGRAPHY M.MODE WITH DOPPLER STUDY:

#### FAIR TRANSTHORACIC ECHOCARDIOGRAPHIC WINDOW MORPHOLOGY:

MITRAL VALVE	NORMAL	TRICUSPID VALVE	NORMAL
AORTIC VALVE	NORMAL	PULMONARY VALVE	NORMAL

#### M.MODE EXAMINATION:

AO	27	mm	LA	31	Mm	IVS-D	9	mm
IVS-S	16	mm	LVID	33	Mm	LVSD	20	mm
LVPW-D	9	mm	LVPW-S	16	Mm	RV	-	mm
RVWT		mm	EDV		MI	LVVS		ml
LVEF	70 %		RWMA			ABSENT		

#### CHAMBERS:

LA	NORMAL	RA	NORMAL
LV	NORMAL	RV	NORMAL
PERICARDIUM	NORMAL		

#### COLOUR DOPPLER:

MITRAL VALVE				
E VELOCITY	0.74	m/sec	PEAK GRADIENT	Mm/hg
A VELOCITY	0.57	m/sec	MEAN GRADIENT	Mm/hg
MVA BY PHT		Cm2	MVA BY PLANIMETRY	Cm2
MITRAL REGURGITATION		ABSENT		
AORTIC VALVE				
PEAK VELOCITY	1.0	m/sec	PEAK GRADIENT	mm/hg
AR VMAX		m/sec	MEAN GRADIENT	mm/hg
AORTIC REGURGITATION		ABSENT		
TRICUSPID VALVE				
PEAK VELOCITY	0.50	m/sec	PEAK GRADIENT	mm/hg
MEAN VELOCITY		m/sec	MEAN GRADIENT	mm/hg
VM <sub>ax</sub> VELOCITY				-
TRICUSPID REGURGITATION		ABSENT		
PULMONARY VALVE				
PEAK VELOCITY	0.89	M/sec.	PEAK GRADIENT	Mm/hg
MEAN VELOCITY			MEAN GRADIENT	Mm/hg
PULMONARY REGURGITATION		ABSENT		

Transcript by.

Dr. Piyush Goyal  
 M.B.B.S., D.M.R.D.  
 RMC Reg No. 017996

Dr. Ashish Choudhary  
 MBBS, MD (Radio Diagnosis)  
 Fetal Medicine Consultant  
 FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain  
 MBBS, DNB, (Radio-Diagnosis)  
 RMC No. 21687

Dr. Navneet Agarwal  
 MD, DNB (Radio Diagnosis)  
 RMC No. 33613/14911

Dr. Poorvi Malik  
 MBBS, MD, DNB (Radio Diagnosis)  
 RMC No. 21505

# Dr. Goyal's

## Path Lab & Imaging Centre

B-51, Ganesh Nagar, Near Metro Pillar No. 109-110, New Sanganer Road, Jaipur  
Tele : 0141-2293346, 4049787, 9887049787  
Website : www.drgoyalspathlab.com E-mail : drgoyalpiyush@gmail.com



Date :- 24/02/2024 09:19:09	Patient ID :- 12235997
<b>NAME :- Mr. NARENDRA SINGH</b>	Ref. By Doctor:-BOB
Sex / Age :- Male 46 Yrs 7 Mon 13 Days	Lab/Hosp :-
Company :- MediWheel	

Final Authentication : 24/02/2024 13:56:15

### Impression--

1. Normal LV size & contractility
2. No RWMA, LVEF 70 %.
3. Normal cardiac chamber.
4. Normal valve
5. No clot, no vegetation, no pericardial effusion.

  
(Cardiologist)

\*\*\* End of Report \*\*\*

Transcript by.

Dr. Piyush Goyal  
M.B.B.S., D.M.R.D.  
RMC Reg No. 017898

Dr. Ashish Choudhary  
MBBS, MD (Radio Diagnosis)  
Fetal Medicine Consultant  
FMF ID - 260517 | RMC No 22430

Dr. Abhishek Jain  
MBBS, DNB, (Radio-Diagnosis)  
RMC No. 21687

Dr. Navneet Agarwal  
MD, DNB (Radio Diagnosis)  
RMC No. 33813/14911

Dr. Poorvi Malik  
MBBS, MD, DNB (Radio Diagnosis)  
RMC No. 21505

