



Subject: Aadhaar parth

From: PARTH NAIR <parthnair99@gmail.com>

Date: 19/11/2022, 08:12 am
To: kannurreception.ddrc@srl.in











DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

ACCESSION NO: 4053VK001846 AGE: 29 Years SEX: Male

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units

MEDIWHEEL HEALTH CHEKUP BELOW 40(M)TMT

OPTHAL

OPTHAL COMPLETED

TREADMILL TEST

TREADMILL TEST COMPLETED

PHYSICAL EXAMINATION

PHYSICAL EXAMINATION COMPLETED



Page 1 Of 9

Scan to View Report





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

AGE: 29 Years ACCESSION NO: 4053VK001846 SEX: Male

RECEIVED: 19/11/2022 10:16 20/11/2022 19:37 DRAWN: REPORTED:

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Units

MEDIWHEEL HEALTH CHEKUP BELOW 40(M)TMT

BUN/CREAT RATIO

BUN/CREAT RATIO 15 5.00 - 15.00

CREATININE, SERUM

0.90 **CREATININE** 18 - 60 yrs : 0.9 - 1.3 mg/dL

GLUCOSE, POST-PRANDIAL, PLASMA RESULT PENDING

GLUCOSE, FASTING, PLASMA

GLUCOSE, FASTING, PLASMA 98 Diabetes Mellitus: > or = 126. mg/dL

Impaired fasting Glucose/ Prediabetes: 101 - 125. Hypoglycemia : < 55.

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD

GLYCOSYLATED HEMOGLOBIN (HBA1C) Normal : 4.0 - 5.6%.%

Non-diabetic level : < 5.7%. Diabetic : >6.5%

Glycemic control goal

More stringent goal : < 6.5 %. General goal : < 7%. Less stringent goal : < 8%.

Glycemic targets in CKD :-If eGFR > 60 : < 7%. If eGFR < 60: 7 - 8.5%.

CORONARY RISK PROFILE (LIPID PROFILE), SERUM

CHOLESTEROL Desirable: < 200 mg/dL

Borderline: 200-239 High : >or= 240

TRIGLYCERIDES Normal 118 : < 150 mg/dL

> High : 150-199

Hypertriglyceridemia: 200-499 Very High: > 499

Low General range: 40-60 HDL CHOLESTEROL 33

DIRECT LDL CHOLESTEROL 119 Optimum : < 100 mg/dL

Above Optimum: 100-139 Borderline High: 130-159 High : 160-189

Very High : >or= 190





mg/dL





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

ACCESSION NO: 4053VK001846 AGE: 29 Years SEX: Male

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	Results			Units
NON HDL CHOLESTEROL	144	High	Desirable-Less than 130 Above Desirable-130-159 Borderline High-160-189 High-190-219 Very High- >or =220	mg/dL
CHOL/HDL RATIO	5.4	High	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0 High Risk	
LDL/HDL RATIO	3.6	High	0.5-3 Desirable/Low risk 3.1-6 Borderline/Moderate risk >6.0 High Risk	
VERY LOW DENSITY LIPOPROTEIN	23.5		= 30</td <td>mg/dL</td>	mg/dL
LIVER FUNCTION TEST WITH GGT				
BILIRUBIN, TOTAL	0.80		General Range : < 1.1	mg/dL
BILIRUBIN, DIRECT	0.21		General Range : < 0.2	mg/dL
BILIRUBIN, INDIRECT	0.59		0.00 - 0.60	mg/dL
TOTAL PROTEIN	7.7		Ambulatory: 6.4 - 8.3 Recumbant: 6 - 7.8	g/dL
ALBUMIN	4.4		20-60yrs: 3.5 - 5.2	g/dL
GLOBULIN	3.3		2.0 - 4.0	g/dL
ALBUMIN/GLOBULIN RATIO	1.3		1.0 - 2.0	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	28		Adults: < 40	U/L
ALANINE AMINOTRANSFERASE (ALT/SGPT)	35		Adults: < 45	U/L
ALKALINE PHOSPHATASE	96		Adult(<60yrs): 40 - 130	U/L
GAMMA GLUTAMYL TRANSFERASE (GGT)	17		Adult(male) : < 60	U/L
TOTAL PROTEIN, SERUM				
TOTAL PROTEIN	7.7		Ambulatory: 6.4 - 8.3 Recumbant: 6 - 7.8	g/dL
URIC ACID, SERUM				
URIC ACID	6.5		Adults: 3.4-7	mg/dL
ABO GROUP & RH TYPE, EDTA WHOLE BLOOD				
ABO GROUP	TYPE O			
RH TYPE	POSITIVE			
BLOOD COUNTS				
HEMOGLOBIN	15.3		13.0 - 17.0	g/dL
RED BLOOD CELL COUNT	4.89		4.5 - 5.5	mil/μL
WHITE BLOOD CELL COUNT	9.16		4.0 - 10.0	thou/µL





Page 3 Of 9





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

ACCESSION NO: 4053VK001846 AGE: 29 Years SEX: Male

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	Results			Units
PLATELET COUNT	309		150 - 410	thou/µL
RBC AND PLATELET INDICES				
HEMATOCRIT	45.6		40 - 50	%
MEAN CORPUSCULAR VOL	93.3		83 - 101	fL
MEAN CORPUSCULAR HGB.	31.2		27.0 - 32.0	pg
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION MEAN PLATELET VOLUME	33.59.9		31.5 - 34.5 6.8 - 10.9	g/dL fL
WBC DIFFERENTIAL COUNT - NLR	9.9		0.0 - 10.9	IL.
SEGMENTED NEUTROPHILS	64		40 - 80	%
ABSOLUTE NEUTROPHIL COUNT	5.86		2.0 - 7.0	π thou/μL
LYMPHOCYTES	28		20 - 40	%
ABSOLUTE LYMPHOCYTE COUNT	2.56		1 - 3	thou/µL
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	2.4		1 3	τιιου, με
EOSINOPHILS	6		1 - 6	%
ABSOLUTE EOSINOPHIL COUNT	0.55	High	0.02 - 0.50	thou/µL
MONOCYTES	1	_	2 - 10	%
ABSOLUTE MONOCYTE COUNT	0.09		0.20 - 1.00	thou/µL
BASOPHILS	1		0 - 2	%
ERYTHRO SEDIMENTATION RATE, BLOOD				
SEDIMENTATION RATE (ESR)	10		0 - 14	mm at 1 hr
STOOL: OVA & PARASITE	RESULT PENDING			
SUGAR URINE - POST PRANDIAL	RESULT PENDING			
THYROID PANEL, SERUM				
Т3	129.30		80.00 - 200.00	ng/dL
T4	9.90		5.10 - 14.10	μg/dl
TSH 3RD GENERATION	2.250		21-50 yrs : 0.4 - 4.2	μIU/mL
URINE ANALYSIS				
COLOR	PALE YELLOW			
APPEARANCE	CLEAR			
PH	6.0		4.7 - 7.5	
SPECIFIC GRAVITY	1.020		1.003 - 1.035	
PROTEIN	DETECTED (SMALL)	NOT DETECTED	
NITRITE	NOT DETECTED		NOT DETECTED	



Page 4 Of 9





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

4053VK001846 AGE: 29 Years ACCESSION NO: SEX: Male

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	Results		Units
WBC	1-2	0-5	/HPF
RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
CHEMICAL EXAMINATION, URINE			
GLUCOSE	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BILIRUBIN	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
MICROSCOPIC EXAMINATION, URINE			
EPITHELIAL CELLS	DETECTED (OCCASIONAL)	NOT DETECTED	/HPF
SERUM BLOOD UREA NITROGEN			
BLOOD UREA NITROGEN	13	Adult(<60 yrs) : 6 to 20	mg/dL
SUGAR URINE - FASTING			
SUGAR URINE - FASTING	NOT DETECTED	NOT DETECTED	

Interpretation(s)

CREATININE, SERUM-

Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
 Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:
• Myasthenia Gravis

• Muscular dystrophy
GLUCOSE, FASTING, PLASMAADA 2012 guidelines for adults as follows:
Pre-diabetics: 100 - 125 mg/dL
Diabetic: > or = 126 mg/dL

(Ref: Tietz 4th Edition & ADA 2012 Guidelines)

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased

any condition that afters the life span of the red blood cells has the potential to after the Grib level. Samples from patients with heritory the anemia. Samples from patients with polycythemia or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of testing such as glycated serum protein (fructosamine) should be considered.



Page 5 Of 9





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

29 Years 4053VK001846 AGE: SEX: Male ACCESSION NO:

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Units

"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient

References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.
- 2. Forsham PH. Diabetes Mellitus:A rational plan for management. Postgrad Med 1982, 71,139-154.
- 3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184. CORONARY RISK PROFILE (LIPID PROFILE), SERUM-

Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don't cause any signs or symptoms, so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn't need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination provides valuable information for the assessment of coronary heart disease risk. It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely. HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease. Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL).

NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.

Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings.

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult. TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome,Protein-losing enteropathy etc. URIC ACID, SERUM-Causes of Increased levels

- Dietary
 High Protein Intake.
- Prolonged Fasting, Rapid weight loss
- Gout

Lesch nyhan syndrome. Type 2 DM.

Metabolic syndrome.

Causes of decreased levels

- Low Zinc IntakeOCP's
- Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

Drink plenty of fluids



Page 6 Of 9







DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

4053VK001846 29 Years AGE: SEX: Male ACCESSION NO:

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Units

- · Limit animal proteins
- · High Fibre foods
- Vit C Intake

• Antioxidant rich foods ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-

Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.

The test is performed by both forward as well as reverse grouping methods.

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

WBC DIFFERENTIAL COUNT - NLR-

The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504

This ratio element is a calculated parameter and out of NABL scope.

ERYTHRO SEDIMENTATION RATE, BLOOD
Erythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0 -1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as polkilocytosis, spherocytosis or sickle cells.

- 1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition
 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition" THYROID PANEL, SERUM-

Triiodothyronine T3, is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

Enculating infinite is nee and protogramy active. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Levels in TOTAL T4 TSH3G TOTAL T3 Pregnancy (µg/dL) (µIU/mL) (ng/dL) First Trimester 6.6 - 12.4 6.6 - 15.5 0.1 - 2.5 0.2 - 3.0 81 - 190 100 - 260 2nd Trimester 3rd Trimester 6.6 - 15.5 0.3 - 3.0 100 - 260
Below mentioned are the guidelines for age related reference ranges for T3 and T4. T4 T3

(μg/dL) 1-3 day: 8.2 - 19.9 (ng/dL) New Born: 75 - 260 1 Week: 6.0 - 15.9

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

Reference:

- 1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.
- Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.
 Behrman R.E. Kilegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition MICROSCOPIC EXAMINATION, URINE-

Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever



Page 7 Of 9

Scan to View Details





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

4053VK001846 AGE: 29 Years ACCESSION NO: SEX: Male

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SFLF CLIENT PATIENT ID:

Test Report Status Results Units

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders. Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine. Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and

proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

- High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal
- Renal Failure

Post Renal • Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

- Liver disease
- SIADH.

SUGAR URINE - FASTING-METHOD: DIPSTICK/BENEDICT'S TEST



Page 8 Of 9





DDRC SRL DIAGNOSTICS

KANNUR KERALA, INDIA Tel: 93334 93334

Email: customercare.ddrc@srl.in

PATIENT NAME: PARTH MADHU NAIR PATIENT ID: PARTM1911934053

ACCESSION NO: 4053VK001846 AGE: 29 Years SEX: Male

DRAWN: RECEIVED: 19/11/2022 10:16 REPORTED: 20/11/2022 19:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Units

MEDIWHEEL HEALTH CHEKUP BELOW 40(M)TMT

ECG WITH REPORT

REPORT

COMPLETED

USG ABDOMEN AND PELVIS

REPORT

COMPLETED

CHEST X-RAY WITH REPORT

REPORT

COMPLETED

End Of Report
Please visit www.srlworld.com for related Test Information for this accession

JINSHA KRISHNAN

LAB TECHNICIAN

DR.INDUSARATH S
CONSULTANT PATHOLOGIST

NIMISHA K LAB TECHNICIAN

time

SREENA A LAB TECHNICIAN





Page 9 Of 9



OPTHALMOLOGY REPORT

TO WHOM-SO-EVER IT MAY CONCERN

This is to certify that I have examined Mr. PARTH MADHU NAIR, 29 years Male on 19.11.2022 and his visual standards are as follows:

erron nerice	OD	OS
UNCORRECTED DISTANCE VISUAL ACUITY	6/24	6/24
UNCORRECTED NEAR VISUAL ACUITY	N6	N6
VISION WITH PRESENT GLASS	6/6(P)	6/6(P)
BEST CORRECTED VISUAL ACUITY	6/6(P)	6/6(P)
COLOUR VISION	NORMAL	NORMAL

NOTE: HISTORY OF SPECS SINCE 14 YEARS, LAST CHANGED 3 YEARS BACK
NO RELEVANT MEDICAL HISTORY

VIMEGA.V OPTOMETRIST DATE: 19.11.2022



INDIA'S LEADING DIAGNOSTICS NETWORK

Name	Mr. PARTH MADHU NAIR	Age/Sex	29/Male
Ref: By:	MEDIWEEL	Date	19.11.2022

ULTRASOUND SCAN OF ABDOMEN AND PELVIS

(With relevant image copies)

LIVER: Normal in size and **shows diffusely increased echotexture**. No e/o focal parenchymal lesions / IHBD. PV, HV & IVC are within normal limits.

GB: Normally distended, normal wall thickness. No e/o calculi/polyps/pericholecystic collections.

CBD: Normal

PANCREAS: Head and body visualized, and are of normal size and echotexture. No e/o focal/diffuse parenchymal lesions/ductal dilatation/calculi. Tail could not be visualized due to poor acoustic window.

SPLEEN: Normal in size and echotexture. Splenic vein shows normal diameter.

KIDNEYS: Both kidneys are normal in size and echotexture. No e/o calculi/

hydronephrosis/ focal lesions/ perinephric collections.

RIGHT KIDNEY: Measures 113 x 50 mms **LEFT KIDNEY:** Measures 106 x 52 mms

UB: Moderately distended, shows normal wall thickness. No e/o calculi/

growth/diverticulae. Both UV junctions are within normal limits.

PROSTATE: 17 cc, normal in size and echotexture.

No e/o intraperitoneal free fluid/ abdominal lymphadenopathy /mass lesion.

IMPRESSION:

- > GRADE I FATTY LIVER.
- > NO OTHER SONOLOGICALLY DETECTED ABNORMALITY.

Dr. P.NIYAZI NASIR MBBS, DMRD

(Because of technical and technological limitation complete diagnosis cannot be assured on imaging sonography. Clinical correlation, consultation if required repeat imaging required in the event of controversies. This document is not for legal purposes).

Dr. P. NIYAZI NASIR. MBBS, DMRD
REG. No. 41419
CONSULTANT RADIOLOGIST
DDRC SRL DIAGNOSTIC (P) LTD.
KANNUR





Name	Mr. PARTH MADHU NAIR	Age/Sex	29/Male
Ref: By:	MEDIWHEEL	Date	19.11.2022

Thanks for referral

CHEST X-RAY - PA VIEW

Trachea is central. Carina and principal bronchi are normal.

Cardio-thoracic ratio is within normal limits.

Both lungs show normal Broncho-vascular markings. No definite focal opacities noted.

No volume loss in either hemithorax.

No definite mediastinal widening or other abnormalities noted.

CP angles, diaphragm, bony cage and soft tissue shadows - not remarkable.

IMPRESSION:

Normal X-ray chest

DR. P. NIYAZI NASIR, MBBS, DMRD

(Because of technical and technological limitation complete diagnosis cannot be assured on imaging sonography. Clinical correlation, consultation if required repeat imaging required in the event of controversies. This document is not for legal purposes).

Dr. P. NIYAZI NASIR. MBBS, DMRD REG. No. 41419 CONSULTANT RADIOLOGIST DDRC SRL DIAGNOSTIC (P) LTD. KANNUR