





CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL

ESTATE, S.V. ROAD, GOREGAON (W) Mumbai, 400062

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: **0002VC075624** AGE: 30 Years SEX: Female

REPORTED: DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u> Results	Biological Reference Interval Units
---	-------------------------------------

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

BLOOD COUNTS,EDTA WHOLE BLOOD				
HEMOGLOBIN	12.4		12.0 - 15.0	g/dL
METHOD : PHOTOMETRIC MEASUREMENT				3/
RED BLOOD CELL COUNT	5.34	High	3.8 - 4.8	mil/µL
METHOD : COULTER PRINCIPLE				
WHITE BLOOD CELL COUNT	8.40		4.0 - 10.0	thou/µL
METHOD: COULTER PRINCIPLE				
PLATELET COUNT	324		150 - 410	thou/µL
METHOD: ELECTRONIC IMPEDENCE & MICROSCOPY				
RBC AND PLATELET INDICES				
HEMATOCRIT	39.2		36.0 - 46.0	%
METHOD: CALCULATED PARAMETER				
MEAN CORPUSCULAR VOL	73.4	Low	83.0 - 101.0	fL
METHOD : DERIVED PARAMETER FROM RBC HISTOGRAM				
MEAN CORPUSCULAR HGB.	23.2	Low	27.0 - 32.0	pg
METHOD: CALCULATED PARAMETER				
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION	31.6		31.5 - 34.5	g/dL
METHOD: CALCULATED PARAMETER				
MENTZER INDEX	13.8			
RED CELL DISTRIBUTION WIDTH	15.3	High	11.6 - 14.0	%
METHOD: DERIVED PARAMETER FROM RBC HISTOGRAM				
MEAN PLATELET VOLUME	8.8		6.8 - 10.9	fL
METHOD: DERIVED PARAMETER FROM PLATELET HISTOGRAM				
WBC DIFFERENTIAL COUNT - NLR				
SEGMENTED NEUTROPHILS	72		40 - 80	%
METHOD: VCSN TECHNOLOGY/ MICROSCOPY				
ABSOLUTE NEUTROPHIL COUNT	6.05		2.0 - 7.0	thou/µL
METHOD: CALCULATED PARAMETER				
LYMPHOCYTES	21		20 - 40	%
METHOD: VCSN TECHNOLOGY/ MICROSCOPY				
ABSOLUTE LYMPHOCYTE COUNT	1.76		1.0 - 3.0	thou/µL
METHOD: CALCULATED PARAMETER				
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	3.4			
METHOD : CALCULATED				
EOSINOPHILS	1		1.0 - 6.0	%



Page 1 Of 15 Scan to View Report







CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS**:

PRIYANKA N MESHRAM

SRL Ltd
PRIME SQUARE BUILDING,PLOT NO 1,GAIWADI INDUSTRIAL
ESTATE,S.V. ROAD,GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results		Biological Reference Inte	erval Units
METHOD: VCSN TECHNOLO	OGY/ MICROSCOPY				
ABSOLUTE EOSINOPHI	IL COUNT	0.08		0.02 - 0.50	thou/µL
METHOD : CALCULATED PAR	RAMETER				
MONOCYTES		5		2.0 - 10.0	%
METHOD: VCSN TECHNOLO	OGY/ MICROSCOPY				
ABSOLUTE MONOCYTE	COUNT	0.42		0.2 - 1.0	thou/µL
METHOD : CALCULATED PAR	RAMETER				
BASOPHILS		1		0 - 1	%
METHOD: VCSN TECHNOLO	OGY/ MICROSCOPY				
ABSOLUTE BASOPHIL	COUNT	0.08		0.02 - 0.10	thou/µL
METHOD : CALCULATED PAR	RAMETER				
MORPHOLOGY					
RBC		Mild anisopoikilo	cytosis. Mi	crocytic hypochromic with over	alocytes.
METHOD : MICROSCOPIC EX	XAMINATION				
WBC		Normal morphol	ogy		
METHOD : MICROSCOPIC EX	XAMINATION				
PLATELETS		Adequate			
METHOD : ELECTRONIC IMP	PEDENCE & MICROSCOPY				
* ERYTHRO SEDIME	NTATION RATE, BLOOD				
SEDIMENTATION RATE	E (ESR)	24	High	0 - 20	mm at 1 hr
METHOD : AUTOMATED (PH	OTOMETRICAL CAPILLARY STOPP	ED FLOW KINETIC ANALYSIS)			
GLUCOSE, FASTING,	PLASMA				
GLUCOSE, FASTING, P	LASMA	83		74 - 99	mg/dL
METHOD : SPECTROPHOTON					5, -
	IOGLOBIN, EDTA WHO	LE BLOOD			
GLYCOSYLATED HEMO		5.2		Non-diabetic: < 5.7 Pre-diabetics: 5.7 - 6.4 Diabetics: > or = 6.5 ADA Target: 7.0 Action suggested: > 8.0	%
METHOD : ION- EXCHANGE	HPLC			Action suggested. > 0.0	
MEAN PLASMA GLUCOS		102.5		< 116.0	mg/dL
METHOD : CALCULATED PAR		-			J, -
GLUCOSE, POST-PRA	NDIAL, PLASMA				
GLUCOSE, POST-PRAN	•	80		70 - 139	mg/dL
METHOD : SPECTROPHOTON	,	00		, 0 133	mg, ac

CORONARY RISK PROFILE (LIPID PROFILE), SERUM











CLIENT CODE: C000138356
CLIENT'S NAME AND ADDRESS:

PRIYANKA N MESHRAM

SRL Ltd
PRIME SQUARE BUILDING,PLOT NO 1,GAIWADI INDUSTRIAL

ESTATE, S.V. ROAD, GOREGAON (W) Mumbai, 400062

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Un
01101 F07FD 01	4.40	5
CHOLESTEROL	148	Desirable cholesterol level mg/d < 200 Borderline high cholesterol 200 - 239 High cholesterol > / = 240
METHOD : SPECTROPHOTOMETRY, ENZYMATIC COLORI	·	•
TRIGLYCERIDES	101	Normal: < 150 mg/d Borderline high: 150 - 199 High: 200 - 499 Very High: >/= 500
METHOD: SPECTROPHOTOMETRY, ENZYMATIC ENDPOI		Law I UDI I I I I I I I I I I I I I I I I I
HDL CHOLESTEROL	37	Low Low HDL cholesterol mg/d < 40 High HDL cholesterol > / = 60
METHOD: SPECTROPHOTOMETRY, HOMOGENEOUS DIF	RECT ENZYMATIC COLORIMETRIC	
DIRECT LDL CHOLESTEROL	107	Optimal : < 100 mg/d Near optimal/above optimal : 100 - 129 Borderline high : 130 - 159 High : 160 - 189 Very high : $> / = 190$
METHOD: SPECTROPHOTOMETRY, HOMOGENEOUS EN	ZYMATIC COLORIMETRIC	
NON HDL CHOLESTEROL	111	Desirable : < 130 mg/d Above Desirable : $130 - 159$ Borderline High : $160 - 189$ High : $190 - 219$ Very high : $> / = 220$
METHOD: CALCULATED PARAMETER		
CHOL/HDL RATIO	4	Low Risk : 3.3 - 4.4 Average Risk : 4.5 - 7.0 Moderate Risk : 7.1 - 11.0 High Risk : > 11.0
METHOD : CALCULATED PARAMETER		
LDL/HDL RATIO	2.9	Desirable/Low Risk: 0.5 - 3.0 Borderline/Moderate Risk: 3.1 - 6.0 High Risk: > 6.0
METHOD: CALCULATED PARAMETER		
VERY LOW DENSITY LIPOPROTEIN METHOD: CALCULATED PARAMETER	20.2	< or = 30.0 mg/d
LIVER FUNCTION PROFILE, SERUM		
BILIRUBIN, TOTAL	0.35	Upto 1.2 mg/d
METHOD: SPECTROPHOTOMETRY COLORIMETRIC -DIA		Opto 1.2 Ilig/u

METHOD: SPECTROPHOTOMETRY, COLORIMETRIC -DIAZO METHOD











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

SRL Ltd
PRIME SQUARE BUILDING,PLOT NO 1,GAIWADI INDUSTRIAL
ESTATE,S.V. ROAD,GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax:

CIN - U74899PB1995PLC045956

PATIENT ID : **PRIYF02079127**

PATIENT NAME: PRIYANKA N MESHRAM

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results	Biological Reference	Interval Units
BILIRUBIN, DIRECT		0.18	0.0 - 0.2	mg/dL
METHOD : SPECTROPHOTO				
BILIRUBIN, INDIRECT		0.17	0.1 - 1.0	mg/dL
METHOD : CALCULATED PA	RAMETER			
TOTAL PROTEIN		7.1	6.0 - 8.0	g/dL
	METRY, COLORIMETRIC -BI	URET, REAGENT BLANK, SERUM BLANK		
ALBUMIN		4.3	3.97 - 4.94	g/dL
METHOD : SPECTROPHOTO	METRY, BROMOCRESOL GRI	EEN(BCG) - DYE BINDING		
GLOBULIN		2.8	2.0 - 3.5	g/dL
METHOD : CALCULATED PA	RAMETER			
ALBUMIN/GLOBULIN F	OITA	1.5	1.0 - 2.1	RATIO
METHOD : CALCULATED PA	RAMETER			
ASPARTATE AMINOTR	ANSFERASE (AST/SO	GOT) 15	Upto 32	U/L
METHOD : SPECTROPHOTO	METRY, WITHOUT PYRIDOX	AL PHOSPHATE ACTIVATION(P5P) - IFCC		
ALANINE AMINOTRAN	SFERASE (ALT/SGPT)) 11	Upto 33	U/L
METHOD : SPECTROPHOTO	METRY, WITHOUT PYRIDOX	AL PHOSPHATE ACTIVATION(P5P) - IFCC		
ALKALINE PHOSPHATA	ASE	74	35 - 104	U/L
METHOD : SPECTROPHOTO	METRY, PNPP, AMP BUFFER	- IFCC		
GAMMA GLUTAMYL TR	` ,	9	< 40	U/L
METHOD : SPECTROPHOTO	METRY, ENZYMATIC COLOR	IMETRIC - G-GLUTAMYL-CARBOXY-NITRO	ANILIDE - IFCC	
LACTATE DEHYDROGE	NASE	150	< 223	U/L
METHOD : SPECTROPHOTO	METRY, LACTATE TO PYRUV	ATE - UV-IFCC		
SERUM BLOOD UREA	A NITROGEN			
BLOOD UREA NITROG	EN	6	6 - 20	mg/dL
METHOD : SPECTROPHOTO	METRY, UREASE -COLORIMI	ETRIC		
CREATININE, SERUI	М			
CREATININE		0.62	0.60 - 1.10	mg/dL
METHOD : SPECTROPHOTO	METRY, JAFFE'S ALKALINE F	PICRATE KINETIC - RATE BLANKED - IFCC	-IDMS STANDARIZED	
BUN/CREAT RATIO				
BUN/CREAT RATIO		9.00	8 - 15	
METHOD : CALCULATED PA	RAMETER			
URIC ACID, SERUM				
URIC ACID		4.3	2.4 - 5.7	mg/dL
METHOD : SPECTROPHOTO	METRY, ENZYMATIC COLOR			3,
TOTAL PROTEIN, SE				
TOTAL PROTEIN		7.1	6.0 - 8.0	g/dL
	METRY COLORIMETRIC -RI	URET, REAGENT BLANK, SERUM BLANK	0.0 0.0	g/uL
HEITIOD . SPECINOFIIOTO	TIETRI, COLORIFIETRIC "DI	ORET, REAGENT BEANK, SERON BEANK		











CLIENT CODE: C000138356
CLIENT'S NAME AND ADDRESS:

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING,PLOT NO 1,GAIWADI INDUSTRIAL

ESTATE, S. V. ROAD, GOREGAON (W) Mumbai, 400062 MAHARASHTRA, INDIA

Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status	<u>Final</u>	Results		Biological Reference	Interval Units
ALDUMIN CEDUM					
ALBUMIN, SERUM		4.2		2.07. 4.04	- / -
ALBUMIN	METRY PROMOCRESOL OF	4.3		3.97 - 4.94	g/dL
METHOD : SPECTROPHOTON	METRY, BROMOCRESOL GI	REEN(BCG) - DYE BINDING			
GLOBULIN					
GLOBULIN		2.8		2.0 - 3.5	g/dL
METHOD : CALCULATED PAR					
ELECTROLYTES (NA/	/K/CL), SERUM				
SODIUM		135	Low	136 - 145	mmol/L
METHOD : ISE INDIRECT					
POTASSIUM		4.20		3.5 - 5.1	mmol/L
METHOD: ISE INDIRECT					
CHLORIDE		102		98 - 106	mmol/L
METHOD: ISE INDIRECT					
URINALYSIS					
COLOR		PALE YELLOW			
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY				
APPEARANCE		CLEAR			
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY				
PH		6.0		4.7 - 7.5	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY- DO	UBLE INDICATOR METHOD			
SPECIFIC GRAVITY		<=1.005		1.003 - 1.035	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY- PKA	A CHANGE OF AN IONIC POLYELECTROLYTE			
GLUCOSE		NOT DETECTED		NOT DETECTED	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY, DO	UBLE SEQUENTIAL ENZYME REACTION-GOI	D/POD		
PROTEIN		NOT DETECTED		NOT DETECTED	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY - PR	OTEIN-ERROR-OF-INDICATOR PRINCIPLE			
KETONES		NOT DETECTED		NOT DETECTED	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY, RO	THERA'S PRINCIPLE			
BLOOD		NOT DETECTED		NOT DETECTED	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY, PER	OXIDASE LIKE ACTIVITY OF HAEMOGLOBI	N		
BILIRUBIN		NOT DETECTED		NOT DETECTED	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY, DIA	ZOTIZATION- COUPLING OF BILIRUBIN WI	TH DIAZ	OTIZED SALT	
UROBILINOGEN		NORMAL		NORMAL	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY - EH	RLICH REACTION			
NITRITE		NOT DETECTED		NOT DETECTED	
METHOD : REFLECTANCE SF	PECTROPHOTOMETRY, COI	NVERSION OF NITRATE TO NITRITE			
PUS CELL (WBC'S)		1-2		0-5	/HPF
·					











CLIENT CODE: C000138356
CLIENT'S NAME AND ADDRESS:

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING,PLOT NO 1,GAIWADI INDUSTRIAL

3rd Trimester: 0.21 - 3.15

ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference	Interval Units
EPITHELIAL CELLS	1-2	0-5	/HPF
METHOD: MICROSCOPIC EXAMINATION			
ERYTHROCYTES (RBC'S)	NOT DETECTED	NOT DETECTED	/HPF
CASTS	NOT DETECTED		
METHOD: MICROSCOPIC EXAMINATION			
CRYSTALS	NOT DETECTED		
METHOD: MICROSCOPIC EXAMINATION			
BACTERIA	NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAMINATION			

Comments

URINALYSIS: MICROSCOPIC EXAMINATION OF URINE IS CARRIED OUT ON CENTRIFUGED URINARY SEDIMENT.

NOTE: KINDLY EXERT CAUTION DURING INTERPRETATION OF FINDINGS REPORTED IN URINALYSIS WHERE IN THE SAMPLE IS MORE THAN TWO HOURS OLD.

THYROID PANEL, SERUM

Т3	108.0	Non-Pregnant Women 80.0 - 200.0 Pregnant Women 1st Trimester105.0 - 230.0 2nd Trimester129.0 - 262.0 3rd Trimester135.0 - 262.0	ng/dL
METHOD: COMPETITIVE ELECTROCHEMILUMINESCENCE I	IMMUNOASSAY		
T4	6.67	Non-Pregnant Women 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70	μg/dL
METHOD: COMPETITIVE ELECTROCHEMILUMINESCENCE I	MMUNOASSAY		
TSH 3RD GENERATION	1.880	Non Pregnant Women 0.27 - 4.20 Pregnant Women 1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10	μIU/mL

 ${\tt METHOD: SANDWICH\ ELECTROCHEMILUMINESCENCE\ IMMUNOASSAY}$

STOOL: OVA & PARASITE

COLOUR BROWN
CONSISTENCY SEMI FORMED
ODOUR FAECAL

MUCUS NOT DETECTED NOT DETECTED











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING,PLOT NO 1,GAIWADI INDUSTRIAL

ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>I</u>	<u>inal</u>	Results	Biological Reference In	terval Units
VISIBLE BLOOD		ABSENT	ABSENT	
POLYMORPHONUCLEAR L	EUKOCYTES	0 - 1	0 - 5	/HPF
METHOD: MICROSCOPIC EXAM	INATION			
RED BLOOD CELLS		NOT DETECTED	NOT DETECTED	/HPF
METHOD: MICROSCOPIC EXAM	INATION			
MACROPHAGES		NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAM				
CHARCOT-LEYDEN CRYST	_	NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAM	INATION			
TROPHOZOITES		NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAM	INATION			
CYSTS		NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAM	INATION			
OVA		NOT DETECTED		
METHOD: MICROSCOPIC EXAM	INATION			
LARVAE		NOT DETECTED	NOT DETECTED	
METHOD: MICROSCOPIC EXAM	INATION			
ADULT PARASITE		NOT DETECTED		
METHOD : MICROSCOPIC EXAM	INATION			
OCCULT BLOOD		NOT DETECTED	NOT DETECTED	
METHOD : MODIFIED GUAIAC M				
ABO GROUP & RH TYPE	E, EDTA WHOLE BLOOD			
ABO GROUP		Α		
METHOD : HAEMAGGLUTINATIO	ON (AUTOMATED)			
RH TYPE		POSITIVE		
METHOD : HAEMAGGLUTINATIO	ON (AUTOMATED)			
* XRAY-CHEST				
IMPRESSION		NO ABNORMALITY DE	TECTED	
TMT OR ECHO				
TMT OR ECHO		*ON HOLD*		
* ECG				
ECG		WITHIN NORMAL LIM	ITS	
* MEDICAL HISTORY				
RELEVANT PRESENT HIST	TORY	FULLY VACCINATED FO	OR COVID 19 ACK MEDICATION BACK	
DELEVANT BACT LITERS				

JAUNDICE 5 YRS BACK

GALL BLADDER STONE 5 YRS BACK



RELEVANT PAST HISTORY









CLIENT CODE: C000138356
CLIENT'S NAME AND ADDRESS:

PRIYANKA N MESHRAM

SRL Ltd

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL

ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: **0002VC075624** AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
RELEVANT PERSONAL HISTORY	NOT SIGNIFICANT	
MENSTRUAL HISTORY (FOR FEMALES)	REGULAR	
LMP (FOR FEMALES)	22/03/2022	
RELEVANT FAMILY HISTORY	NOT SIGNIFICANT	
HISTORY OF MEDICATIONS	NOT SIGNIFICANT	
* ANTHROPOMETRIC DATA & BMI		
HEIGHT IN METERS	1.53	mts
WEIGHT IN KGS.	56	Kgs
ВМІ	24	BMI & Weight Status as follows: kg/sqmts Below 18.5: Underweight 18.5 - 24.9: Normal 25.0 - 29.9: Overweight 30.0 and Above: Obese
* CENEDAL EVAMINATION		

* GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE	NORMAL
PHYSICAL ATTITUDE	NORMAL
GENERAL APPEARANCE / NUTRITIONAL STATUS	HEALTHY
BUILT / SKELETAL FRAMEWORK	AVERAGE
FACIAL APPEARANCE	NORMAL
SKIN	NORMAL
UPPER LIMB	NORMAL
LOWER LIMB	NORMAL
NECK	NORMAL

NECK LYMPHATICS / SALIVARY GLANDS NOT ENLARGED OR TENDER

THYROID GLAND NOT ENLARGED CAROTID PULSATION NORMAL

TEMPERATURE NORMAL

PULSE 76/MIN REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID

BRUIT

RESPIRATORY RATE NORMAL

* CARDIOVASCULAR SYSTEM

BP 100/70 MM HG mm/Hg

(SUPINE)

PERICARDIUM NORMAL APEX BEAT NORMAL

HEART SOUNDS S1, S2 HEARD NORMALLY









PRIYF02079127

CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

ACCESSION NO: 0002VC075624

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL

PATIENT ID:

ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM

AGE: 30 Years

30/03/2022 15:37 DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED:

SEX: Female

DRAWN . 29/03/2022 09:30	RECEIVED : 29/03/2022 09:40	REFORTED: 30/03/2022 13.37
REFERRING DOCTOR: SELF		CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
MURMURS	ABSENT	
* RESPIRATORY SYSTEM		
SIZE AND SHAPE OF CHEST	NORMAL	
MOVEMENTS OF CHEST	SYMMETRICAL	
BREATH SOUNDS INTENSITY	NORMAL	
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)	
ADDED SOUNDS	ABSENT	
* PER ABDOMEN		
APPEARANCE	NORMAL	
VENOUS PROMINENCE	ABSENT	
LIVER	NOT PALPABLE	
SPLEEN	NOT PALPABLE	
HERNIA	ABSENT	
* CENTRAL NERVOUS SYSTEM		
HIGHER FUNCTIONS	NORMAL	
CRANIAL NERVES	NORMAL	

CEREBELLAR FUNCTIONS **NORMAL** SENSORY SYSTEM **NORMAL** MOTOR SYSTEM **NORMAL REFLEXES** NORMAL

* MUSCULOSKELETAL SYSTEM

SPINE NORMAL JOINTS NORMAL

* BASIC EYE EXAMINATION

CONJUNCTIVA NORMAL EYELIDS NORMAL EYE MOVEMENTS **NORMAL CORNEA NORMAL**

DISTANT VISION RIGHT EYE WITH GLASSES REDUCE VISUAL ACUITY (6/9) DISTANT VISION LEFT EYE WITH GLASSES REDUCE VISUAL ACUITY (6/18) WITHIN NORMAL LIMIT (N6) NEAR VISION RIGHT EYE WITH GLASSES NEAR VISION LEFT EYE WITH GLASSES WITHIN NORMAL LIMIT (N6)

NORMAL (17/17) COLOUR VISION

* BASIC ENT EXAMINATION











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL

ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final**

EXTERNAL EAR CANAL NORMAI TYMPANIC MEMBRANE NORMAL

NOSE NO ABNORMALITY DETECTED

SINUSES NORMAL

THROAT NO ABNORMALITY DETECTED

NOT ENLARGED TONSILS

* BASIC DENTAL EXAMINATION

TEETH NORMAL GUMS HEALTHY

* SUMMARY

RELEVANT HISTORY NOT SIGNIFICANT

RELEVANT GP EXAMINATION FINDINGS REDUCE VISUAL ACUITY DISTANT VISION BOTH EYES

RELEVANT LAB INVESTIGATIONS RAISED ESR (24)

LOW HDL CHOLESTEROL (37)

LOW SODIUM (135)

RELEVANT NON PATHOLOGY DIAGNOSTICS USG-NO ABNORMALITIES DETECTED REMARKS / RECOMMENDATIONS RAISED ESR,LOW HDL CHOLESTEROL

FOLLOW UP WITH PHYSICIAN

Interpretation(s)
BLOOD COUNTS,EDTA WHOLE BLOOD-

The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-

Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

diagnosing a case of beta thalassaemia trait.
WBC DIFFERENTIAL COUNT - NLR-

The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope. ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0 -1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

Reference:

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition

2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin
3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition" GLUCOSE, FASTING, PLASMA-

ADA 2021 guidelines for adults, after 8 hrs fasting is as follows: Pre-diabetics: 100 - 125 mg/dL











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL ESTATE, S.V. ROAD, GOREGAON (W) Mumbai, 400062

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final**

Diabetic: > or = 126 mg/dL

GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood, the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks.

Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased

glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells.

Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia, increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of testing such as glycated serum protein (fructosamine) should be considered.

"Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient

References

- 1. Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006,
- 2. Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.
- 2. Forsial From Diabetes Melitus. A radioial plan for inal agent multi-rostylau med 1962, 11,139-134.

 3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184.

 GLUCOSE, POST-PRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75grams of glucose in 300 ml water, over a period of 5

CORONARY RISK PROFILE (LIPID PROFILE), SERUM-

Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don't cause any signs or so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn't need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination provides valuable information for the assessment of coronary heart disease risk. It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely.HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease. Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL). NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.

Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings.

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult. LIVER FUNCTION PROFILE, SERUM-

attaches sugar molecules to bilirubin.

LIVER FUNCTION PROFILE Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units **Final**

anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc

SERUM BLOOD UREA NITROGEN-

Causes of Increased levels

Pre renal

- High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal
 Renal Failure

• Malignancy, Nephrolithiasis, Prostatism

Causes of decreased levels

Liver disease

STADH.

CREATININE, SERUM-

Higher than normal level may be due to:

- Blockage in the urinary tract
 Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
 Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia GravisMuscular dystrophy

URIC ACID, SERUM-

Causes of Increased levels

Dietary

- High Protein Intake.
- Prolonged Fasting,Rapid weight loss.
- Gout

Lesch nyhan syndrome.

Type 2 DM. Metabolic syndrome.

Causes of decreased levels

- Low Zinc Intake
- OCP's
- Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels

- Drink plenty of fluids
- Limit animal proteinsHigh Fibre foods
- Vit C Intake
- · Antioxidant rich foods

TOTAL PROTEIN, SERUM-

Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and alobulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units Final

syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-

Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc. ELECTROLYTES (NA/K/CL), SERUM-

Sodium levels are Increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is common in vomiting, diarrhea, alcoholism, folic acid deficiency and primary aldosteronism. Hyperkalemia may be seen in end-stage renal failure, hemolysis, trauma, Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion. Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hyperfuction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonged vomiting,

URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise.

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders. Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine. Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and

proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia THYROID PANEL, SERUM-

Triiodothyronine T3, is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3

Levels in TOTAL T4 TSH3Ġ TOTAL T3 (µg/dL) (µIU/mL) (ng/dL) Pregnancy First Trimester 6.6 - 12.4 6.6 - 15.5 0.1 - 2.5 81 - 190 100 - 260 2nd Trimester 3rd Trimester 6.6 - 15.5 0.3 - 3.0 100 - 260
Below mentioned are the guidelines for age related reference ranges for T3 and T4.

T3 T4 (ng/dL) (µg/dL) New Born: 75 - 260 1-3 day: 8.2 - 19.9 1 Week: 6.0 - 15.9

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group.

Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

Reference:

- 1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.
- 2. Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.
 3. Behrman R.E. Kilegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition

STOOL: OVA & PARASITE-

Acute infective diarrhoea and gastroenteritis (diarrhoea with vomiting) are major causes of ill health and premature death in developing countries. Loss of water and electrolytes from the body can lead to severe dehydration which if untreated, can be rapidly fatal in young children, especially that are malnourished, hypoglycaemic, and

Laboratory diagnosis of parasitic infection is mainly based on microscopic examination and the gross examination of the stool specimen. Depending on the nature of the parasite, the microscopic observations include the identification of cysts, ova, trophozoites, larvae or portions of adult structure. The two classes of parasites that cause human infection are the Protozoa and Helminths. The protozoan infections include amoebiasis mainly caused by Entamoeba histolytica and giardiasis caused by Giardia











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL ESTATE, S.V. ROAD, GOREGAON (W)

Mumbai, 400062 MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Biological Reference Interval Units <u>Final</u>

lamblia. The common helminthic parasites are Trichuris trichiura, Ascaris lumbricoides, Strongyloides stercoralis, Taenia sp. etc

ABO GROUP & RH TYPE, EDTA WHOLE BLOODBlood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for

The test is performed by both forward as well as reverse grouping methods.

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.











CLIENT CODE: C000138356 **CLIENT'S NAME AND ADDRESS:**

PRIYANKA N MESHRAM

PRIME SQUARE BUILDING, PLOT NO 1, GAIWADI INDUSTRIAL

ESTATE, S.V. ROAD, GOREGAON (W) Mumbai, 400062

MAHARASHTRA, INDIA Tel: 9111591115, Fax: CIN - U74899PB1995PLC045956

PATIENT NAME: PRIYANKA N MESHRAM PATIENT ID: PRIYF02079127

ACCESSION NO: 0002VC075624 AGE: 30 Years SEX: Female

DRAWN: 29/03/2022 09:38 RECEIVED: 29/03/2022 09:40 REPORTED: 30/03/2022 15:37

REFERRING DOCTOR: SELF CLIENT PATIENT ID:

Test Report Status Results Units **Final**

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

* ULTRASOUND ABDOMEN

ULTRASOUND ABDOMEN

NO ABNORMALITIES DETECTED

End Of Report

Please visit www.srlworld.com for related Test Information for this accession TEST MARKED WITH '*' ARE OUTSIDE THE NABL ACCREDITED SCOPE OF THE LABORATORY.

Kshama P. MD (Reg No. MMC2000/02/0552) **Biochemist**

Dr. J N Shukla ,MBBS, AFIH **Consultant Physician**

Dr. Swati Karmarkar, MD, DNB, DMRD **Consultant Radiologist**

Dr. Sukanya Verma (Reg.No.MMC2012/03/0443) **Consultant Microbiologist**

CONDITIONS OF LABORATORY TESTING & REPORTING

- 1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
- 2. All Tests are performed and reported as per the turnaround time stated in the SRL Directory of services (DOS).
- 3. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
- 4. A requested test might not be performed if:
- a. Specimen received is insufficient or inappropriate specimen quality is unsatisfactory
 - b. Incorrect specimen type
- c. Request for testing is withdrawn by the ordering doctor or patient
- d. There is a discrepancy between the label on the specimen container and the name on the test requisition form

- 5. The results of a laboratory test are dependent on the quality of the sample as well as the assay technology.
- 6. Result delays could be because of uncontrolled circumstances. e.g. assay run failure.
- 7. Tests parameters marked by asterisks are excluded from the "scope" of NABL accredited tests. (If laboratory is accredited).
- 8. Laboratory results should be correlated with clinical information to determine Final diagnosis.
- 9. Test results are not valid for Medico- legal purposes. 10. In case of gueries or unexpected test results please call at SRL customer care (91115 91115). Post proper investigation repeat analysis may be carried out.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII, Mohali 160062



