




Garg Pathology

Certified by :
National Accreditation Board For Testing & Calibration Laboratories
ISO 9001:2008
Garden House Colony, Near Nai Sarak, Garh Road, Meerut
Ph.: 0121-2600454, 8979608687, 9837772828

DR. MONIKA GARG
M.D. (Path) Gold Medalist
Former Pathologist :
St. Stephan's Hospital, Delhi

| | | |
|--|-------------------|---|
| PUID : 221008/601 | C. NO: 601 | Collection Time : 08-Oct-2022 8:39AM |
| Patient Name : Mrs. NEHA AGARWAL 30Y / Female | | Receiving Time : 08-Oct-2022 8:48AM |
| Referred By : Dr. BANK OF BARODA | | Reporting Time : 08-Oct-2022 11:55AM |
| Sample By : | | Centre Name : Garg Pathology Lab - TPA |
| Organization : | |  |

| Investigation | Results | Units | Biological Ref-Interval |
|---------------|---------|-------|-------------------------|
|---------------|---------|-------|-------------------------|

BIOCHEMISTRY (SERUM)

THYRIOD PROFILE*

| | | | |
|---|-------|-------|-----------|
| Triiodothyronine (T3) * (ECLIA) | 1.471 | ng/dl | 0.79-1.58 |
|---|-------|-------|-----------|

| | | | |
|------------------------------------|-------|-------|----------|
| Thyroxine (T4) * (ECLIA) | 9.965 | ug/dl | 4.9-11.0 |
|------------------------------------|-------|-------|----------|

| | | | |
|---|-------|--------|-----------|
| THYROID STIMULATING HORMONE (T) (ECLIA) | 3.062 | uIU/ml | 0.38-5.30 |
|---|-------|--------|-----------|

Normal Range:-

| | |
|--------------|----------|
| 1 TO 4 DAYS | 2.7-26.5 |
| 4 TO 30 DAYS | 1.2-13.1 |

Hyperthyroid patient have suppressed TSH values, with the exception of those few individuals whos have hyperthyroidism caused by TSH producing pituitary tumor or other rare disordes such as pituitary resistance to thyroid hormones. Subclinical hyperthyroidism is defined as low TSH with levels of T4 and T3 within the reference interval. In most patients with hypothyroidism,serum TSH results are markedly elevated, but results are low in individuals with hypothyroidism caused by pituitary or hypothalamic disorders. An important cause of both increased and decreased TSH results is NTI. Patients with NTI tend to have low TSH results during their acute illness ,then TSH rises to within or above the reference range with resolution of the underlying illness,and finally returns to within the reference range. The situation is complicated because drugs,including glucagon and dopamine,suppress TSH . Sensitive TSH assays are helpful in evaluation of treatment with thyroid hormone both for replacement therapy and suppressive doses for malignant thyroid disease.

-----{END OF REPORT }-----



*THIS TEST IS NOT UNDER NABL SCOPE

Checked By Technician:

Page 1 of 1

Dr. Monika Garg
MBBS, MD(Path)
(Consultant Pathologist)

२१ सँडे सुविधा उपलब्ध है।

