Name:Mrs. SINDHU SRIVASTAVAPID No.:MED111622596SID No.:223006996Age / Sex:41 Year(s) / FemaleType:OPRef. Dr:MediWheel	Register On         : 29/0           Collection On         : 29/0           Report On         : 29/0	04/2023 8:28 AM 04/2023 9:31 AM /05/2023 7:45 PM 05/2023 12:13 PM	DIAGNOSTICS
Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BLOOD GROUPING AND Rh TYPING (EDTA Blood/Agglutination) INTERPRETATION: Reconfirm the Blood g Complete Blood Count With - ESR	'B' 'Positive' group and Typing before b	lood transfusion	
Haemoglobin (EDTA Blood'Spectrophotometry)	11.6	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocri (EDTA Blood/Derived from Impedance)	t <b>36.8</b>	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.06	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	90.6	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	) 28.6	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	31.5	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	14.0	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	44.39	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	7640	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	66.2	%	40 - 75
Lymphocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	25.1	%	20 - 45







APPROVED BY

The results pertain to sample tested.

Page 1 of 8

Name	:	Mrs. SINDHU SRIVASTAVA				
PID No.	:	MED111622596	Register On	:	29/04/2023 8:28 AM	
SID No.	:	223006996	<b>Collection On</b>	:	29/04/2023 9:31 AM	
Age / Sex	:	41 Year(s) / Female	Report On	:	02/05/2023 7:45 PM	mec
Туре	:	OP	Printed On	:	04/05/2023 12:13 PM	DIAGNO
Ref. Dr	:	MediWheel				

Investigation	<u>Observed</u> <u>Value</u>	Unit	Biological Reference Interval
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	4.8	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	3.8	%	01 - 10
Basophils (EDTA Blood/Impedance Variation & Flow Cytometry)	0.1	%	00 - 02
INTERPRETATION: Tests done on Automated F	ive Part cell counte	r. All abnormal results are r	eviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	5.06	10^3 / µl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	1.92	10^3 / µl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.37	10^3 / µl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.29	10^3 / µl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.01	10^3 / µl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	168	10^3 / µl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	13.7	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.23	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Blood/Automated - Westergren method)	34	mm/hr	< 20







APPROVED BY

The results pertain to sample tested.

Page 2 of 8

Name	: Mrs. SINDHU SRIVAST	AVA	
PID No.	: MED111622596	Register On : 29/04/2023 8:28 AM	
SID No.	: 223006996	Collection On : 29/04/2023 9:31 AM	$\mathbf{O}$
Age / Sex	: 41 Year(s) / Female	Report On : 02/05/2023 7:45 PM	medall
Туре	: OP	Printed On : 04/05/2023 12:13 PM	DIAGNOSTICS

#### Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
BUN / Creatinine Ratio	13.4		6.0 - 22.0
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	102.7	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

**INTERPRETATION:** Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose, Fasting (Urine) (Urine - F/GOD - POD)	Negative		Negative
Glucose Postprandial (PPBS) (Plasma - PP/GOD-PAP)	117.2	mg/dL	70 - 140

#### **INTERPRETATION:**

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV/derived)	8.8	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.67	mg/dL	0.6 - 1.1

**INTERPRETATION:** Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin, cefazolin, ACE inhibitors, angiotensin II receptor antagonists, N-acetylcysteine, chemotherapeutic agent such as flucytosine etc.

Uric Acid (Serum/ <i>Enzymatic</i> )	3.5	mg/dL	2.6 - 6.0
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	0.60	mg/dL	0.1 - 1.2
Dr ANITHA Consultant Pathologist MCI -112788 VERIFIED BY			DR.SUNDAR ELAYAPERUMAL MD, CIC CONSULTANT MICROBIOLOGIST REG ND, 41854 APPROVED BY

The results pertain to sample tested.

Page 3 of 8

Name	: Mrs. SINDHU SRIVASTAVA		
PID No.	: MED111622596	Register On	: 29/04/2023 8:28 AM
SID No.	: 223006996	<b>Collection On</b>	: 29/04/2023 9:31 AM
Age / Sex	: 41 Year(s) / Female	Report On	: 02/05/2023 7:45 PM
Туре	: OP	Printed On	: 04/05/2023 12:13 PM
Ref. Dr	: MediWheel		



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.14	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/ <i>Derived</i> )	0.46	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/ <i>Modified IFCC</i> )	19.9	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/ <i>Modified IFCC</i> )	17.2	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	18.4	U/L	< 38
Alkaline Phosphatase (SAP) (Serum/ <i>Modified IFCC)</i>	84.3	U/L	42 - 98
Total Protein (Serum/ <i>Biuret</i> )	7.11	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	3.98	gm/dL	3.5 - 5.2
Globulin (Serum/Derived)	3.13	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.27		1.1 - 2.2
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	173.6	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/ <i>GPO-PAP with ATCS)</i>	54.4	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500
Edmittin	∎sa2	K.	Ehr

Consultant Pathologist MCI -112788 VERIFIED BY





APPROVED BY

The results pertain to sample tested.

Dr ANITHA

Page 4 of 8

Name	: Mrs. SINDHU SRIVASTAVA			
PID No.	: MED111622596	Register On : 2	29/04/2023 8:28 AM	
SID No.	: 223006996	Collection On :	29/04/2023 9:31 AM	
Age / Sex	: 41 Year(s) / Female	Report On :	02/05/2023 7:45 PM	medall
Туре	: OP	Printed On :	04/05/2023 12:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			
<u>Investiga</u>	ition	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
increasing variation to	as much as 5 to 10 times the fasting bo. There is evidence recommending for metabolic syndrome, as non-fasti	levels, just a few hour triglycerides estimation	s after eating. Fasting tright on in non-fasting condition	change drastically in response to food, yceride levels show considerable diurnal for evaluating the risk of heart disease and irculating level of triglycerides during most
HDL Cho (Serum/Imr	plesterol nunoinhibition)	51.2	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cho (Serum/Ca.		111.5	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL C (Serum/Ca	holesterol	10.9	mg/dL	< 30
Non HDI (Serum/Ca.	L Cholesterol lculated)	122.4	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >=220

**INTERPRETATION:** 1.Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

3.4

Total Cholesterol/HDL Cholesterol
Ratio
(Serum/Calculated)





Optimal: < 3.3 Low Risk: 3.4 - 4.4 Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0



APPROVED BY

The results pertain to sample tested.

Page 5 of 8

Name	: Mrs. SINDHU SRIVASTAVA			
PID No.	: MED111622596	Register On	: 29/04/2023 8:28 AM	
SID No.	: 223006996	<b>Collection On</b>	: 29/04/2023 9:31 AM	$\mathbf{O}$
Age / Sex	: 41 Year(s) / Female	Report On	: 02/05/2023 7:45 PM	medall
Туре	: OP	Printed On	: 04/05/2023 12:13 PM	DIAGNOSTICS

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Triglyceride/HDL Cholesterol Ratio (TG/HDL) (Serum/ <i>Calculated</i> )	1.1		Optimal: < 2.5 Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0
LDL/HDL Cholesterol Ratio (Serum/Calculated)	2.2		Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 High Risk: > 6.0
<u>Glycosylated Haemoglobin (HbA1c)</u>			
HbA1C (Whole Blood/HPLC)	5.7	%	Normal: 4.5 - 5.6 Prediabetes: 5 7 - 6 4

(Whole Blood/HPLC)Prediabetes: 5.7 - 6.4Diabetic: >= 6.5

Estimated Average Glucose	116.89	mg/dL
(Whole Blood)		

## **INTERPRETATION:** Comments

: MediWheel

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbA1C values. Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

# THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total	1.02	ng/ml	0.7 - 2.04
(Serum/Chemiluminescent Immunometric Assay			
(CLIA))			

## **INTERPRETATION:**

### **Comment :**

Ref. Dr

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.





	E	ha		2	
DR.SUN	1000		MICRO	BIOLO	), CIC

APPROVED BY

The results pertain to sample tested.

Page 6 of 8

Name	: Mrs. SINDHU SRIVASTAVA			
PID No.	: MED111622596	Register On : 29	/04/2023 8:28 AM	<b>~</b>
SID No.	: 223006996	Collection On : 29	9/04/2023 9:31 AM	$\mathbf{O}$
Age / Sex	: 41 Year(s) / Female	Report On : 02	2/05/2023 7:45 PM	medall
Туре	: OP	Printed On : 04	4/05/2023 12:13 PM	DIAGNOSTICS
Ref. Dr	: MediWheel			
Investiga	ation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
	xine) - Total emiluminescent Immunometric Assay	6.79	µg/dl	4.2 - 12.0
<b>Comment</b> Total T4 va	ariation can be seen in other conditio	n like pregnancy, drugs	nephrosis etc. In such case	es, Free T4 is recommended as it is
(Serum/Cha	ally active. yroid Stimulating Hormone) emiluminescent Immunometric Assay	1.50	µIU/mL	0.35 - 5.50
<ul> <li>(CLIA))</li> <li>INTERPRETATION:</li> <li>Reference range for cord blood - upto 20</li> <li>1 st trimester: 0.1-2.5</li> <li>2 nd trimester 0.2-3.0</li> <li>3 rd trimester : 0.3-3.0</li> <li>(Indian Thyroid Society Guidelines)</li> <li>Comment:</li> <li>1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.</li> <li>2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.</li> <li>3.Values&amp;amplt0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.</li> </ul>				
<u>Urine An</u>	alysis - Routine			
COLOUI (Urine)	R	Pale yellow		Yellow to Amber
APPEAR (Urine)	AANCE	Turbid		Clear
Protein (Urine/Prot	tein error of indicator)	Negative		Negative
Glucose (Urine/GO)	D - POD)	Negative		Negative
Pus Cells (Urine/Auto	s omated ó"Flow cytometry )	1 - 2	/hpf	NIL
	Dr ANITHA sultant Pathologist MCI -112788 ERIFIED BY			DR.SUNDAR ELAYAPERUMAL MD, CIC CONSULTANT MICROBIOLOGIST REG NO. 41854

The results pertain to sample tested.

Page 7 of 8

Name	: Mrs. SINDHU SRIVASTAVA			
PID No.	: MED111622596	Register On	: 29/04/2023 8:28 AM	
SID No.	: 223006996	<b>Collection On</b>	: 29/04/2023 9:31 AM	$\mathbf{O}$
Age / Sex	: 41 Year(s) / Female	Report On	: 02/05/2023 7:45 PM	medall
Туре	: OP	Printed On	: 04/05/2023 12:13 PM	DIAGNOSTICS

# Ref. Dr : MediWheel

Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	Biological Reference Interval
Epithelial Cells (Urine/Automated 6"Flow cytometry)	2 - 3	/hpf	NIL
RBCs (Urine/Automated 6"Flow cytometry )	NIL	/hpf	NIL
Casts (Urine/Automated 6"Flow cytometry )	NIL	/hpf	NIL
Crystals (Urine/Automated ó"Flow cytometry)	NIL	/hpf	NIL

OthersAmmonium urate(Urine)crystal present .

**INTERPRETATION:** Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.





-- End of Report --



APPROVED BY

The results pertain to sample tested.

Page 8 of 8

Name	: Mrs. SINDHU SRIVASTAVA	Register On	: 29/04/2023 8:28 AM
PID No.	: MED111622596	Collection On	: 29/04/2023 9:31 AM
SID No.	: 223006996	Report On	: 02/05/2023 7:45 PM
Age / Sex	: 41 Year(s) / Female	Printed On	<sup>:</sup> 04/05/2023 12:13 PM
Ref. Dr	: MediWheel	OP / IP	: OP



PAP Smear by LBC( Liquid based Cytology )

SPECIMEN NO: Cy 1076/2023

# MICROSCOPIC FINDINGS:

ADEQUACY: Satisfactory.

PREDOMINANT CELLS: Superficial and intermediate cells.

BACKGROUND: Neutrophils.

**ORGANISMS:** No specific organisms.

IMPRESSION:

Negative for intraepithelial lesion/ malignancy.





APPROVED BY

-- End of Report --

The results pertain to sample tested.

Page 1 of 1