Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	31240600976
Patient Episode	: H03000063919	Collection Date :	22 Jun 2024 10:00
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 11:15	Reporting Date :	22 Jun 2024 12:32

Department of Transfusion Medicine (Blood Bank)

BLOOD GROUPING, RH TYPING & ANTIBODY SCREEN (TYPE & SCREEN) Specimen-Blood

Blood Group & Rh Typing (Agglutinaton by gel/tube technique)

Blood Group & Rh typing O Rh(D) Positive

Antibody Screening (Microtyping in gel cards using reagent red cells)

Final Antibody Screen Result Negative

Technical Note: ABO grouping and Rh typing is done by cell and serum grouping by microplate / gel technique. Antibody screening is done using a 3 cell panel of reagent red cells coated with Rh, Kell,Duffy,Kidd, Lewis, P,MNS,Lutheran and Xg antigens using gel technique.

Page1 of 4

-----END OF REPORT-----

Dr Himanshu Lamba

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	32240611606
Patient Episode	: H03000063919	Collection Date :	22 Jun 2024 10:00
Referred By Receiving Date	: HEALTH CHECK MHD: 22 Jun 2024 10:40	Reporting Date :	22 Jun 2024 14:15

BIOCHEMISTRY

		Specimen: EDTA Whole blood
HbA1c (Glycosylated Hemoglobin)	7.8 # %	
		HbAlc in % Non diabetic adults : < 5.7 % Prediabetes (At Risk) : 5.7 % - 6.4 % Diabetic Range : > 6.5 %
Estimated Average Glucose (eAG)	177	mg/dl

Use :

1.Monitoring compliance and long-term blood glucose level control in patients with diabetes. 2.Index of diabetic control (direct relationship between poor control and development of complications).

3. Predicting development and progression of diabetic microvascular complications.

Limitations :

A1C values may be falsely elevated or decreased in those with chronic kidney disease.
False elevations may be due in part to analytical interference from carbamylated hemoglobin formed in the presence of elevated concentrations of urea, with some assays.
False decreases in measured A1C may occur with hemodialysis and altered red cell turnover, especially in the setting of erythropoietin treatment

References : Rao.L.V., Michael snyder.L. (2021). Wallach's Interpretation of Diagnostic Tests. 11th Edition. Wolterkluwer. NaderRifai, Andrea Rita Horvath, Carl T.wittwer. (2018) Teitz Text book

of Clinical Chemistry and Molecular Diagnostics.First edition,Elsevier,South Asia.

Page2 of 4

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	32240611606
Patient Episode	: H03000063919	Collection Date :	22 Jun 2024 10:00
Referred By Receiving Date	HEALTH CHECK MHD22 Jun 2024 10:29	Reporting Date :	22 Jun 2024 12:22

BIOCHEMISTRY

mg/dl

[<200]

153

Lipid Profile (Serum)

TOTAL CHOLESTEROL (CHOD/POD)

			Moderate risk:200-239 High risk:>240
TRIGLYCERIDES (GPO/POD)	174	# mg/dl	[<150]
			Borderline high:151-199
			High: 200 - 499
			Very high:>500
HDL - CHOLESTEROL (Direct)	51	mg/dl	[30-60]
Methodology: Homogenous Enzymatic			
VLDL - Cholesterol (Calculated)	35	mg/dl	[10-40]
(CALCULATED) LDL-	CHOLESTEROL	67 mg/dl	[<100]
			Near/Above optimal-100-129
			Borderline High:130-159
			High Risk:160-189
T.Chol/HDL.Chol ratio	3.0		<4.0 Optimal
			4.0-5.0 Borderline
			>6 High Risk
LDL.CHOL/HDL.CHOL Ratio	1.3		<3 Optimal
			3-4 Borderline
			>6 High Risk

Note: Reference ranges based on ATP III Classifications. Recommended to do fasting Lipid Profile after a minimum of 8 hours of overnight fasting.

Technical Notes: Lipid profile is a panel of blood tests that serves as initial broad medical screening tool for abnormalities in lipids, the results of these tests can identify certain genetic

Page3 of 4

P 011 4967 4967 E info@manipalhospitals.com Emergency 011 4040 7070 www.hcmct.in www.manipalhospitals.com/delhi/ Managed by Manipal Hospitals (Dwarka) Private Limited

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	32240611606
Patient Episode	: H03000063919	Collection Date :	22 Jun 2024 10:00
Referred By Receiving Date	HEALTH CHECK MHD22 Jun 2024 10:29	Reporting Date :	22 Jun 2024 12:22

BIOCHEMISTRY

diseases and determine approximate risks for cardiovascular disease, certain forms of pancreatitis and other diseases.

Test Name	Result	Unit	Biological Ref. Interval
TOTAL PSA, Serum (ECLIA)	1.480	ng/mL	[<2.500]

Note : PSA is a glycoprotein that is produced by the prostate gland. Normally, very little PSA is secreted in the blood. Increases in glandular size and tissue damage caused by BPH, prostatitis, or prostate cancer may increase circulating PSA levels.

Caution : Serum markers are not specific for malignancy, and values may vary by method.

Immediate PSA testing following digital rectal examination, ejaculation, prostate massage urethral instrumentation, prostate biopsy may increase PSA levels.

Some patients who have been exposed to animal antigens, may have circulating anti-animal antibodies present. These antibodies may interfere with the assay reagents to produce unreliable results.

-----END OF REPORT-----

Neefan Sugal

Page 4 of 4

Dr. Neelam Singal CONSULTANT BIOCHEMISTRY

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	:	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	:	32240611606
Patient Episode	: H03000063919	Collection Date	:	22 Jun 2024 10:00
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 10:29	Reporting Date	:	22 Jun 2024 12:27

BIOCHEMISTRY

THYROID PROFILE, Serum			ecimen Type : Serum
T3 – Triiodothyronine (ECLIA) T4 – Thyroxine (ECLIA)	0.977 7.040	ng/ml µg/dl	[0.800-2.040] [4.600-10.500]
Thyroid Stimulating Hormone (ECLIA)	1.360	µIU/mL	[0.340-4.250]

Note : TSH levels are subject to circadian variation, reaching peak levels between 2-4.a.m.and at a minimum between 6-10 pm.Factors such as change of seasons hormonal fluctuations, Ca or Fe supplements, high fibre diet, stress and illness affect TSH results.

* References ranges recommended by the American Thyroid Association

1) Thyroid. 2011 Oct;21(10):1081-125.PMID .21787128

2) http://www.thyroid-info.com/articles/tsh-fluctuating.html

Test Name	Result	Unit	Biological Ref. Interval
LIVER FUNCTION TEST (Serum)			
BILIRUBIN-TOTAL (Diazonium Ion)	1.26 #	mg/dl	[0.10-1.20]
BILIRUBIN - DIRECT (Diazotization)	0.65 #	mg/dl	[0.00-0.30]
BILIRUBIN - INDIRECT (Calculated)	0.61	mg/dl	[0.20-1.00]
SGOT/ AST (UV without P5P)	80 #	U/L	[10-50]
SGPT/ ALT (UV without P5P)	117 #	U/L	[0-41]
ALP (p-NPP,kinetic)*	93	U/L	[45-135]
TOTAL PROTEIN (Biuret)	7.6	g/dl	[7.0-9.0]
SERUM ALBUMIN (BCG-dye)	4.8	g/dl	[3.5-5.2]
SERUM GLOBULIN (Calculated)	2.8	g/dl	[1.8-3.4]
ALB/GLOB (A/G) Ratio(Calculated)	1.71		[1.10-1.80]



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	:	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	:	32240611606
Patient Episode	: H03000063919	Collection Date	:	22 Jun 2024 10:00
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 10:29	Reporting Date :	:	22 Jun 2024 12:23

BIOCHEMISTRY

Technical Notes:

Liver function test aids in diagnosis of various pre hepatic, hepatic and post hepatic causes of dysfunction like hemolytic anemia's, viral and alcoholic hepatitis and cholestasis of obstructive causes.

Test Name	Result	Unit B:	iological Ref. Interval
KIDNEY PROFILE (Serum)			
BUN (Urease/GLDH)	6.00	mg/dl	[6.00-20.00]
SERUM CREATININE (Jaffe's method)	0.60 #	mg/dl	[0.80-1.60]
SERUM URIC ACID (Uricase)	5.2	mg/dl	[3.5-7.2]
SERUM CALCIUM (NM-BAPTA)	9.44	mg/dl	[8.00-10.50]
SERUM PHOSPHORUS (Molybdate, UV)	2.0 #	mg/dl	[2.5-4.5]
SERUM SODIUM (ISE)	133.0 #	mmol/l	[134.0-145.0]
SERUM POTASSIUM (ISE)	4.76	mmol/l	[3.50-5.20]
SERUM CHLORIDE (ISE Indirect)	95.7	mmol/L	[95.0-105.0]
eGFR	120.6	ml/min/1.73sq	.m [>60.0]

Technical Note

eGFR which is primarily based on Serum Creatinine is a derivation of CKD-EPI 2009 equation normalized to1.73 sq.m BSA and is not applicable to individuals below 18 years. eGFR tends to be less accurate when Serum Creatinine estimation is indeterminate e.g. patients at extremes of muscle mass, on unusual diets etc. and samples with severe Hemolysis / Icterus / Lipemia.

Page2 of 7

-----END OF REPORT-----

Neefan Suga

Dr. Neelam Singal CONSULTANT BIOCHEMISTRY

P 011 4967 4967 E info@manipalhospitals.com Emergency 011 4040 7070 www.hcmct.in www.manipalhospitals.com/delhi/ Managed by Manipal Hospitals (Dwarka) Private Limited



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age	:	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No	:	32240611607
Patient Episode	: H03000063919	Collection Date	:	22 Jun 2024 10:00
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 10:29	Reporting Date	:	22 Jun 2024 21:59

BIOCHEMISTRY

Specimen Type : Plasma

GLUCOSE-Fasting	(Hexokinase)	201 #	mg/dl	[74-106]	
				Page 3 o:	f 7

-----END OF REPORT-----

Neefam Suga

Dr. Neelam Singal CONSULTANT BIOCHEMISTRY



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age	:	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No	:	33240606808
Patient Episode	: H03000063919	Collection Date	:	22 Jun 2024 10:01
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 10:40	Reporting Date	:	22 Jun 2024 13:26

HAEMATOLOGY

ERYTHROCYTE SEDIMENTATION RATE (Automated) Specimen-Whole Blood

ESR	2.0	mm/1sthour	[0.0-10.0]

Interpretation :

Erythrocyte sedimentation rate (ESR) is a non-specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants (e.g. pyogenic infections, inflammation and malignancies). The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week postpartum.

ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives).

It is especially low (0 -1mm) in polycythemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

Test Name	Result	Unit Bio	ological Ref. Interval
COMPLETE BLOOD COUNT (EDTA Blood)			
WBC Count (Flow cytometry)	5710	/cu.mm	[4000-10000]
RBC Count (Impedence)	5.89 #	million/cu.mm	[4.50-5.50]
Haemoglobin (SLS Method)	13.5	g/dL	[13.0-17.0]
Haematocrit (PCV)	42.4	00	[40.0-50.0]
(RBC Pulse Height Detector Method)			
MCV (Calculated)	72.0 #	fL	[83.0-101.0]
MCH (Calculated)	22.9 #	pg	[25.0-32.0]
MCHC (Calculated)	31.8	g/dL	[31.5-34.5]
Platelet Count (Impedence)	106000 #	/cu.mm	[150000-410000]
RDW-CV (Calculated)	17.3 #	8	[11.6-14.0]
DIFFERENTIAL COUNT			
Neutrophils (Flowcytometry)	58.3	90	[40.0-80.0]
Lymphocytes (Flowcytometry)	32.6	00	[20.0-40.0]

P 011 4967 4967 E info@manipalhospitals.com Emergency 011 4040 7070 www.hcmct.in www.manipalhospitals.com/delhi/ Managed by Manipal Hospitals (Dwarka) Private Limited



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age	:	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No	:	33240606808
Patient Episode	: H03000063919	Collection Date	e:	22 Jun 2024 10:01
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 10:40	Reporting Date	e:	22 Jun 2024 11:48

HAEMATOLOGY

Monocytes (Flowcytometry)	7.2	:	00	[2.0-10.0]
Eosinophils (Flowcytometry)	1.2	1	00	[1.0-6.0]
Basophils (Flowcytometry)	0.7 #	:	90 0	[1.0-2.0]
IG	1.60	:	00	
Neutrophil Absolute(Flouroscence f	low cytometry)	3.3	/cu mm	[2.0-7.0]x10 ³
Lymphocyte Absolute(Flouroscence f	low cytometry)	1.9	/cu mm	[1.0-3.0]x10 ³
Monocyte Absolute(Flouroscence flo	w cytometry)	0.4	/cu mm	[0.2-1.2]x10 ³
Eosinophil Absolute(Flouroscence f	low cytometry)	0.1	/cu mm	[0.0-0.5]x10 ³
Basophil Absolute(Flouroscence flo	w cytometry)	0.0	/cu mm	[0.0-0.1]x10 ³

Complete Blood Count is used to evaluate wide range of health disorders, including anemia, infection, and leukemia. Abnormal increase or decrease in cell counts as revealed may indicate that an underlying medical condition that calls for further evaluation.

-----END OF REPORT------

Dr. Priyanka Bhatia CONSULTANT PATHOLOGY

Page 5 of 7

Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age	:	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No	:	38240602111
Patient Episode	: H03000063919	Collection Date	:	22 Jun 2024 10:01
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 11:01	Reporting Date	:	22 Jun 2024 12:32

CLINICAL PATHOLOGY

Test Name	Result	Biological Ref. Interval
ROUTINE URINE ANALYSIS		
MACROSCOPIC DESCRIPTION		
Colour (Visual)	PALE YELLOW	(Pale Yellow - Yellow)
Appearance (Visual)	CLEAR	
CHEMICAL EXAMINATION		
Reaction[pH]	6.0	(5.0-9.0)
(Reflectancephotometry(Indicator Method	od))	
Specific Gravity	1.005	(1.003-1.035)
(Reflectancephotometry(Indicator Method	od))	
Bilirubin	Negative	NEGATIVE
Protein/Albumin	Negative	(NEGATIVE-TRACE)
(Reflectance photometry(Indicator Met)	nod)/Manual SSA)	
Glucose	DETECTED TRACE	(NEGATIVE)
(Reflectance photometry (GOD-POD/Bene	dict Method))	
Ketone Bodies	NOT DETECTED	(NEGATIVE)
(Reflectance photometry(Legal's Test),	/Manual Rotheras)	
Urobilinogen	NORMAL	(NORMAL)
Reflactance photometry/Diazonium salt	reaction	
Nitrite	NEGATIVE	NEGATIVE
Reflactance photometry/Griess test		
Leukocytes	NIL	NEGATIVE
Reflactance photometry/Action of Ester	rase	
BLOOD	NIL	NEGATIVE
(Reflectance photometry(peroxidase))		
MICROSCOPIC EXAMINATION (Manual) Mo	ethod: Light microscopy on	centrifuged urine
WBC/Pus Cells	1-2 /hpf	(4-6)
Red Blood Cells	NIL	(1-2)
Epithelial Cells	1-2 /hpf	(2-4)
Casts	NIL	(NIL)
Crystals	NIL	(NIL)
Bacteria	NIL	
Yeast cells	NIL	
Interpretation:		
=		

Page6 of 7



Registered Office: Sector-6, Dwarka, New Delhi 110 075

Department Of Laboratory Medicine

Name	: MR NISHU NARAYAN PRASAD	Age :	46 Yr(s) Sex :Male
Registration No	: MH013572384	Lab No :	38240602111
Patient Episode	: H03000063919	Collection Date :	22 Jun 2024 10:01
Referred By Receiving Date	: HEALTH CHECK MHD : 22 Jun 2024 11:01	Reporting Date :	22 Jun 2024 12:32

CLINICAL PATHOLOGY

URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic , urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urina tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine.

Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine.

Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise. Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys Most Common cause is bacterial urinary tract infection.

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration duri infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased Specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decrease Specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis,

bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in case of hemolytic anemia.

Page7 of 7

-----END OF REPORT-----

Dr. Priyanka Bhatia CONSULTANT PATHOLOGY



