

CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015


ABHA NO :

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED :11/02/2024 09:36:59

REPORTED :11/02/2024 15:53:12

Test Report Status <u>Final</u> Results Biological Reference Interval Units

HAEMATOLOGY - CBC				
MEDI WHEEL FULL BODY HEALTH CHECK UP BE	LOW 40 MALE			
BLOOD COUNTS,EDTA WHOLE BLOOD				
HEMOGLOBIN (HB) METHOD: CVANIDE FREE DETERMINATION	14.2	13.0 - 17.0	g/dL	
RED BLOOD CELL (RBC) COUNT METHOD: ELECTRICAL IMPEDANCE	4.80	4.5 - 5.5	mil/µL	
WHITE BLOOD CELL (WBC) COUNT METHOD: ELECTRICAL IMPEDANCE	5.30	4.0 - 10.0	thou/µL	
PLATELET COUNT METHOD: ELECTRONIC IMPEDANCE	194	150 - 410	thou/µL	
RBC AND PLATELET INDICES				
HEMATOCRIT (PCV) METHOD: CALCULATED PARAMETER	44.7	40 - 50	%	
MEAN CORPUSCULAR VOLUME (MCV) METHOD : CALCULATED PARAMETER	93.0	83 - 101	rL.	
MEAN CORPUSCULAR HEMOGLOBIN (MCH) METHOD: CALCULATED PARAMETER	29.6	27.0 - 32.0	Pg	
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC) NETHOD: CALQUATED PARAMETER	31.7	31.5 - 34.5	g/dL	
RED CELL DISTRIBUTION WIDTH (RDW) METHOD: CALCULATED PARAMETER	15.2 High	11.6 - 14.0	96	
MENTZER INDEX	19.4		_	
MEAN PLATELET VOLUME (MPV) METHOD: CALCULATED PARAMETER	12.4 High	6.8 - 10.9	rL.	
WBC DIFFERENTIAL COUNT				
NEUTROPHILS NETHOD: IMPEDIANCE WITH HYDRO FOCUS AND MICROSCOPY	58	40 - 80	96	
LYMPHOCYTES METHOD: IMPEDIANCE WITH HYDRO FOCUS AND MICROSCOPY	34	20 - 40	%	
MONOCYTES	03	2 - 10	%	







Page 1 Of 17

View Details

View Report







CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO

AGE/SEX : 37 Years Male DRAWN ;11/02/2024 09:30:00 RECEIVED: 11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

		i	
Test Report Status <u>Final</u>	Results	Biological Reference Interv	al Units
METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY			D.
EOSINOPHILS	05	1 - 6	96
METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY			
BASOPHILS	00	0 - 2	96
METHOD: IMPEDANCE WITH HYDRO FOCUS AND MICROSCOPY			
ABSOLUTE NEUTROPHIL COUNT	3.07	2.0 - 7.0	thou/µL
METHOD: CALCULATED PARAMETER			
ABSOLUTE LYMPHOCYTE COUNT	1.80	1.0 - 3.0	thou/µL
	1.00	1.0 - 3.0	tilou) pt
METHOD : CALCULATED PARAMETER		03.40	Min man Cord
ABSOLUTE MONOCYTE COUNT	0.16 Low	0.2 - 1.0	thou/µL
METHOD: CALCULATED PARAMETER:			
ABSOLUTE EOSINOPHIL COUNT	0.26	0.02 - 0.50	thou/µL
METHOD: CALCULATED PARAMETER			
ABSOLUTE BASOPHIL COUNT	0 Low	0.02 - 0.10	thou/µL
NEUTROPHIL LAMPHOCATE BATTO (NUB)	1.7		
NEUTROPHIL LYMPHOCYTE RATIO (NLR)	1.7		

Interpretation(s)
BLOOD COUNTS, BDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. Howeverafter 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool **O differentiats cases of iron deficiency anaemia(> 13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for

diagnosing a case of beta thalasseemia trait.

WBC confirmential. COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms technique from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR = 3.0.0 yea

3.3, COVID-19 patients tend to show mild disease.
(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients ; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504.
This ratio element is a calculated parameter and out of NABL scope.

Dr. Akansha Jain **Consultant Pathologist** Page 2 Of 17











CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912

PATIENT ID : ANSHM110287251 CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00

RECEIVED :11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

HAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE

BLOOD

HBA1C 5.7

Non-diabetic: < 5.7 %
Pre-diabetics: 5.7 - 6.4
Diabetics: > or = 6.5
Therapeutic goals: < 7.0
Action suggested : > 8.0
(ADA Guideline 2021)

METHOD: HIGH PERFORMANCE LIQUID CHROMATOGRAPHY (HPLC)

ESTIMATED AVERAGE GLUCOSE(EAG)

METHOD: CALCULATED PARAMETER

116.9 High

< 116.0

mg/dL

Dr. Akansha Jain Consultant Pathologist

Page 3 Of 17





View Details

View Report





CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

1AIPUR 302017 9314660100

ACCESSION NO: 0251XB000912

PATIENT ID: : ANSHM110287251

CLIENT PATIENT ID: 012402110015 ABHA NO

AGE/SEX : 37 Years Male ;11/02/2024 09:30:00 DRAWN

RECEIVED: 11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ERYTHROCYTE SEDIMENTATION RATE (ESR), EDTA BLOOD

mm at 1 hr E.S.R. 0 - 14

METHOD: AUTOMATED (PHOTOMETRICAL CAPILLARY STOPPED FLOW KINETIC ANALYSIS)*

Interpretation(s) GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:

- Evaluating the long-term control of blood glucose concentrations in diabetic patients.
- Diagnosing diabetes

Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic petients, and 2 times per year for well-controlled *VpPa 2 diabetic patients) to determine whether a patients metabolic control has remained continuously within the target range.

1. ANG (Estimated average glucose) converts percentage HAA1; to mg/dl, to compare blood glucose levels.

2. ANG glyesan evaluation of blood glucose levels for the last couple of months.

3. ANG is calculated as ANG (mg/dl) = 28.7 ** HoA1c - 46.7

HbA1c Estimation can get affected due to:

Shortened Brythrocyte survival: Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acuta blood loss, hemolytic

- 1. Snottened a Profescyte survival Any condition tract snortens environce survival or access mean environce age (e.g. recovery from access glood loss, nemo yet anemia) will falsely lower that is the track from the recommended in these patterns which indicates diabetes control over 15 days.

 2. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin.

 3. Iron deficiency anemia is reported to interfere with some assay methods, falsely increasing results.

 4. Interference of hemoglobinopathies in HbA1c estimation is seen in
- a) Homozygous hemoglobinopathy, Pructosamine is recommended for testing of HbA1c.
 b) Heterozygous state detected (D10 is corrected for Hb5 & HbC trait.)

c) HbF > 25% on alternate patform (Boronate affinity chromatography) is recommended for testing of HbA1c.Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy ERYTHROCYTE SEDIMENTATION RATE (ESR),EDTA BLOOD-TEST DESCRIPTION:

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of crythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic; it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition.CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculities, Inflammatory arthritis, Renal disease, Anemia, Halignancies and plasma cell dyscrasias, Acute allergy Tissue Injury, Pregnancy, Estrogen medication, Aging,

Finding a very accelerated ESR(>100 mm/hour) in petients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissuedisease, severe infections such as bacterial endocarditis).

In pregnancy BPE in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm /hr(95 if anemic), ESR returns to normal 4th week post partum.

Decreased in: Polycythermia vera, Sickle cell anemia

False elevated ESR: Increased fibrinogen, Drugs(Vitamin A, Dextran etc.), Hypercholesterolemia
False Decreased: Polikilocytosis, (SickleCells, spherocytes), Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine, salicylates)

1. Nathan and Oski's Hoematology of Infancy and Childhood, 5th edition; 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin; 3. The reference for the adult reference range is "Practical Haematology by Dade and Lewis, 10th edition."

Dr. Akansha Jain Consultant Pathologist





Page 4 Of 17

PERFORMED AT :



PATIENT NAME: ANSHU BOBS8959

CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

1AIPUR 302017 9314660100

ACCESSION NO: 0251XB000912

REF. DOCTOR: SELF

PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00

RECEIVED: 11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

IMMUNOHAEMATOLOGY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP TYPE B

METHOD: TUBE AGGLUTINATION

RH TYPE POSITIVE

METHOD: TUBE AGGLUTINATION

Interpretation(s)
ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified byantigers and antibodies present in the blood. Antigers are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or Ab.

Disclaimer: "Please note, as the results of previous ABO and Rhigroup (Blood Group) for pregnant women are not available, please check with the patient records for

The bask is performed by both forward as well as reverse grouping methods.

Dr. Akansha Jain **Consultant Pathologist** Page 5 Of 17











CODE/NAME & ADDRESS : C000049066 AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

1AIPUR 302017 9314660100

ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO

AGE/SEX : 37 Years Male DRAWN ;11/02/2024 09:30:00 RECEIVED: 11/02/2024 09:36:59

REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

BIOCHEMISTRY

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

GLUCOSE FASTING, FLUORIDE PLASMA

FBS (FASTING BLOOD SUGAR)

METHOD: GLUCOSE OXIDASE

METHOD: GLUCOSE OXIDASE

148 High

74 - 99

mg/dL

GLUCOSE, POST-PRANDIAL, PLASMA

PPBS(POST PRANDIAL BLOOD SUGAR)

112

70 - 140

mg/dL

LIPID PROFILE WITH CALCULATED LDL

237 High CHOLESTEROL, TOTAL

< 200 Desirable

mg/dL

200 - 239 Borderline High >/= 240 High

METHOD: CHOLESTEROL OXIDASE

228 High TRIGLYCERIDES

< 150 Normal

mg/dL

150 - 199 Borderline High 200 - 499 High

>/=500 Very High

METHOD: LIPASE/GPO-PAP NO CORRECTION

HDL CHOLESTEROL 40 < 40 Low >/=60 High mg/dL

METHOD: DIRECT CLEARANCE METHOD

CHOLESTEROL LDL

151 High

< 100 Optimal

mg/dL

mg/dL

100 - 129

Near optimal/ above optimal

130 - 159

Borderline High 160 - 189 High

>/= 190 Very High

197 High Desirable: Less than 130

> Above Desirable: 130 - 159 Borderline High: 160 - 189

High: 190 - 219

Very high: > or = 220

METHOD: CALCULATED PARAMETER

NON HDL CHOLESTEROL

Dr. Akansha Jain **Consultant Pathologist** Page 6 Of 17











CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-INAAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX : 37 Years Male
DRAWN :11/02/2024 09:30:00
RECEIVED :11/02/2024 09:36:59
REPORTED :11/02/2024 15:53:12

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
VERY LOW DENSITY LIPOPROTEIN	45.6 High	= 30.0 mg/dL</td
CHOL/HDL RATIO	5.9 High	3.3 - 4.4 Low Risk 4.5 - 7.0 Average Risk 7.1 - 11.0 Moderate Risk > 11.0
LDL/HDL RATIO	3.8 High	High Risk 0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk

Interpretation(s)

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

THE PARTY OF THE P	ALSO TIS (ALBERTOSCICIONE CHI GIOTASCOMI GI	ensely by empre a construction of amount
Risk Category		
Extreme risk group	A.CAD with > 1 feature of high risk group	
	B. CAD with > 1 feature of Very high risk g	group or recurrent ACS (within 1 year) despite LDL-C < or =
	50 mg/dl or polyvascular disease	
Very High Risk	 Established ASCVD 2. Diabetes with 2 r 	najor risk factors or evidence of end organ damage 3.
	Familial Homozygous Hypercholesterolemi	1
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ	
	damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary	
	Artery Calcium - CAC >300 AU. 7. Lipoprotein a >/= 50mg/dl 8. Non stenotic carotid plaque	
Moderate Risk	2 major ASCVD risk factors	
Low Risk	0-1 major ASCVD risk factors	
Major ASCVD (Ath	erosclerotic cardiovascular disease) Risk Fa	ctors
1. Age > or = 45 years in males and > or = 55 years in females 3. Current Cigarette smoking or tobacco use		
2. Family history of premature ASCVD 4. High blood pressure		
5. Low HDL		

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

٠	vewer treatment goals and status instantion thresholds based on the risk tategories proposed by 12-11 in 2020.				
	Risk Group	Treatment Goals		Consider Drug T	herapy
		LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
	Extreme Risk Group Category A	<50 (Optional goal	< 80 (Optional goal	>OR = 50	>OR = 80
		< OR = 30)	<or 60)<="" =="" td=""><td></td><td></td></or>		

Dr. Akansha Jain Consultant Pathologist





Page 7 Of 17

View Details

View Report







CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED :11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

Extreme Risk Group Category B	<or 30<="" =="" th=""><th><or 60<="" =="" th=""><th>> 30</th><th>>60</th></or></th></or>	<or 60<="" =="" th=""><th>> 30</th><th>>60</th></or>	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

^{*}After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.65	0 - 1	mg/dL
METHOD: DIAZO WITH SULPHANILIC ACID			
BILIRUBIN, DIRECT	0.19	0.00 - 0.25	mg/dL
METHOD: DIAZO WITH SULPHANILIC ACID BILIRUBIN, INDIRECT	0.46	0.1 - 1.0	mg/dL
METHOD : CALCULATED PARAMETER	0.40	0.1 - 1.0	mg/ac
TOTAL PROTEIN	7.4	6.4 - 8.2	g/dL
METHOD: BIURET REACTION, END POINT			-
ALBUMIN	4.5 High	3.8 - 4.4	g/dL
METHOD: BROMOCRESOL GREEN			
GLOBULIN	2.9	2.0 - 4.1	g/dL
METHOD: CALCULATED PARAMETER			RATIO
ALBUMIN/GLOBULIN RATIO METHOD: CALCULATED PARAMETER	1.6	1.0 - 2.1	PORTIO
ASPARTATE AMINOTRANSFERASE(AST/SGOT)	64 High	0 - 37	U/L
METHOD: TRIS BUFFER NO PSP IFCC / SFBC 37° C			-,-
ALANINE AMINOTRANSFERASE (ALT/SGPT)	116 High	0 - 40	U/L
METHOD: TRIS BUFFER NO PSP IFCC / SFBC 37* C			
ALKALINE PHOSPHATASE	112	39 - 117	U/L
METHOD: AMP OPTIMISED TO IFCC 37° C	467 Wieb	44 50	11.0
GAMMA GLUTAMYL TRANSFERASE (GGT) METHOD: GAMMA GLUTAMYL-3 CARBOXY-4 NITROANILIDE (IFCC) 3	167 High	11 - 50	U/L
LACTATE DEHYDROGENASE	413	230 - 460	U/L
E TOTALE DELITOROGENHOL	120	E00 100	

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN 10 5.0 - 18.0 mg/dL

METHOD: UREASE KINETIC

Dr. Akansha Jain Consultant Pathologist



Page 8 Of 17

View Details

View Report







CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-

AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID: ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED :11/02/2024 09:36:59

REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

CREATININE, SERUM

CREATININE 1.27 0.8 - 1.3 mg/dL

METHOD: ALKALINE PICRATE NO DEPROTEINIZATION

BUN/CREAT RATIO

BUN/CREAT RATIO 7.87

METHOD: CALCULATED PARAMETER

URIC ACID, SERUM

URIC ACID 7.5 High 3.4 - 7.0 mg/dL

METHOD: URICASE PEROXIDASE WITH ASCORBATE OXIDASE

TOTAL PROTEIN, SERUM

TOTAL PROTEIN 7.4 6.4 - 8.3 g/dL

METHOD: BIURET REACTION, END POINT

ALBUMIN, SERUM

ALBUMIN 4.5 High 3.8 - 4.4 g/dL

METHOD: BROMOCRESOL GREEN

GLOBULIN

GLOBULIN 2.9 2.0 - 4.1 g/dL

ELECTROLYTES (NA/K/CL), SERUM

Dr. Akansha Jain Consultant Pathologist Page 9 Of 17





View Details

View Benort







REF. DOCTOR: SELF PATIENT NAME: ANSHU BOBS8959

CODE/NAME & ADDRESS : C000049066 AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED: 11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

	i	i i	
Test Report Status <u>Final</u>	Results	Biological Reference	e Interval Units
SODIUM, SERUM	141.0	137 - 145	mmol/L
METHOD: ION-SELECTIVE ELECTRODE POTASSIUM, SERUM	4.20	3.6 - 5.0	mmol/L
METHOD: JON-SELECTIVE ELECTRODE CHLORIDE, SERUM METHOD: JON-SELECTIVE ELECTRODE	101.3	98 - 107	mmol/L

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in:CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, anti depressants (SSRI), antipsychotics.	Decreased in: Low potassium intake, prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome, osmotic diuresis (e.g., hyperglycemia), alkalosis, familial periodic paralysis, trauma (transient). Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenalinsufficiency, hyperaldosis. Drugs: chronic loxative, corticular oils, diuretics.
Increased in: Dehydration (excessives weating, severe vomiting or diarrhea), diabetes mellitus, diabetes insipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice, oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhahdemyolysis, acidosis, dehydration, renal failure, Addison's disease, BTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium-sparing diuretics, NSAIDs, beta-blockers, ACE inhibitors, highdose trimethoprim-sulfamethoxazole.	Intreased In: Runal failure, nephrotic syndrome, RTA dehydration, overtreatment with sallee, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO3-), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemi, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy [Normal serum chloride]

Interpretation(s)
GLUCOSE FASTING, FLUCRIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and sothat no glucose is excreted in the

unitive.

Increased in: Diabetes melitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs:corticosteroids,phenytoin, estrogen, thiazides.

Decreased in: Pencreatic islet cell disease with increased insulin,insulinoma,adrenocortical insufficiency,hypopituitarism,diffuse liver disease, malignancy(adrenocortical,stomach,fibrosarcoma),infant of g diabetic mother,enzyme deficiency diseases(e.g.galactosemia),Drugs-insulin,ethanol,propranolol;sulfonylureas,folbutamide,and other oral hypoglycemic agents.

NOTIE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values),there is wide fluctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

Dr. Akansha Jain **Consultant Pathologist** Page 10 Of 17











CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

1AIPUR 302017 9314660100

ACCESSION NO: 0251XB000912

: ANSHM110287251 CLIENT PATIENT ID: 012402110015

ABHA NO

PATIENT ID:

AGE/SEX : 37 Years Male ;11/02/2024 09:30:00 DRAWN

RECEIVED: 11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycsuria, Glycaemic index & reaponse to food consumed, Alimentary Hypoglycemia, Increased insulin reaponae & sensitivity etc.
GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycsuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER PUNCTION PROFILE, SERUM-

Still which is 8 yellowish pigment found in bile and is 8 breakdown product of normal heme catabolism. Bilirubin is excreted in bile and unine, and elevated levels may give yellow discoloration in joundice. Elevated levels results from increased bilirubin production (eg., hemolysis and ineffective erythropolesis), decreased bilirubin excretion (eg., obstruction and hepatitis), and abnormal bilirubin metabolism (e.g., hereditary and reconstal joundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in five and abnormal bilirubin metabolism (e.g., hereditary and reconstal joundice). Conjugated (direct) bilirubin in five and properties and indirect) bilirubin in journal hepatitis, (purg reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anomia, Transfusion reaction is a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that ches sugar molecules to bilirubin.

AST is an enzyme found in various Parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic angmia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocal ular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, is chemia to the liver, chronic

hepetitis, obstruction of bile ducts, cirrhosis.

ALP in a protein found in almost all body tessues. Tessues with higher amounts of ALP include the liver, bile ducts and bone. Blevated ALP levels are seen in Billiery obstruction. Osteoblastic bone furnisms, osteomalacia, hepatitis, Hyperparathyroldism, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoldosis etc. Lower-than-normal ALP levels seen in Hyperparathyroldisms, Leukemia, Lymphoma, Pagets disease, Rickets, Sarcoldosis etc. Lower-than-normal ALP levels seen in Hyperparathyroldisms, Mainutrition, Protein deficiency, Wilsons disease,

GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain

and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated gerum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs set.

Total Protein also known as total protein, is a biochemical feast for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiplie myeloma, Waldenstroms

disease Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Halabsorption, Hainutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

Albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hyposlipuminemia) can be caused by: Liver disease like circhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic dearance, mainutrition and wasting es-

BLOCG UREA NITROGEN (BUN), SERUM-Causes of Increased levels include Prenenal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Fallure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)
Causes of decreased level include Liverdisease, SIADH.

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM-Higher than normal level may be due to:

Blackage in the urinary triact, kidney problems, such as kidney demage or failure, infection, or reduced blood flow, Loss of body fluid (dehydration), Muscle problems, such as becaused on muscle fibers, Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preciampsia)

Lower than normal level may be due to: Myasthenia Gravis, Muscuophy

URIC ACID, SERUM-Causes of Increased levels-Chetary(High Problem Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch myhan syndrome, Type 2 DM, Metabolic syndrome Causes of decreased levels-Low Zinc intake, OCP, Multiple Sciences

TOTAL PROPEIN, SERUM-is a biochemical bast for measuring the total amount of protein in serum. Problem in the plasma is made up of albumin and globulin.

Higher them, server all levels may be due to: 100 contributes or infection, including MM and benefits the contribute or infection, including MM and benefits the residence. Mildigent these diseases

Higher-than-normal levels may be due 10: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstroms disease

Lower-than-normal levels may be due to: Agammagisbulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malebsorption, Mainutrition, Nephrotic syndrome, Protein-losing enteropathy etc.

ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver, Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, mainutrition and wasting etc.

Dr. Akansha Jain Consultant Pathologist



Page 11 Of 17







CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID: ANSHM11028725

PATIENT ID : ANSHM110287251 CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX :37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED :11/02/2024 09:36:59

REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - URINALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW

METHOD: GROSS EXAMINATION

APPEARANCE CLEAR

METHOD: GROSS EXAMINATION

CHEMICAL EXAMINATION, URINE

PH 5.5 4.7 - 7.5

METHOD: DOUBLE INDICATOR PRINCIPLE
SPECIFIC GRAVITY 1.020 1.003 - 1.035

METHOD: JONIC CONCENTRATION METHOD

PROTEIN NOT DETECTED NEGATIVE

METHOD: PROTEIN ERROR OF INDICATORS WITH REFLECTANCE

GLUCOSE NOT DETECTED NEGATIVE

METHOD: GLUCOSE OXIDASE PEROXIDASE / BENEDICTS

KETONES NOT DETECTED NOT DETECTED

METHOD: SODIUM NITROPRUSSIDE REACTION

BLOOD NOT DETECTED NEGATIVE METHOD: PEROCIDASE ANTI PEROXIDASE

BILIRUBIN NOT DETECTED NOT DETECTED

METHOD : DIPSTICK

UROBILINOGEN NORMAL NORMAL
METHOD: EHRLICH REACTION REFLECTANCE

NITRITE NOT DETECTED NOT DETECTED

METHOD: NITRATE TO NITRITE CONVERSION METHOD

LEUKOCYTE ESTERASE NOT DETECTED NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS NOT DETECTED NOT DETECTED /HPF

METHOD: MICROSCOPIC EXAMINATION

PUS CELL (WBC'S) 1-2 0-5 /HPF

METHOD: DIPSTICK, MICROSCOPY

Dr. Akansha Jain Consultant Pathologist



Page 12 Of 17

View Details

View Benort





CODE/NAME & ADDRESS : C000049066 AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100

ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015 ABHA NO

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED: 11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status <u>Final</u>	Results	Biological Reference	e Interval Units
EPITHELIAL CELLS METHOD: MICROSCOPIC EXAMINATION	1-2	0-5	/HPF
CASTS	NOT DETECTED		
METHOD: MICROSCOPIC EXAMINATION CRYSTALS METHOD: MICROSCOPIC EXAMINATION	NOT DETECTED		
BACTERIA METHOD: MICROSCOPIC EXAMINATION	NOT DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases

Dr. Akansha Jain **Consultant Pathologist**





Page 13 Of 17



C/O Aakriti Labs Pvt Ltd, 3. Mahatma Gandhi Narg,Gandhi Nagar Mod, Tonk Road Jaipur, 302015 Rajasthan, India





CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID: ANSHM110287251

CLIENT PATIENT ID: 012402110015


ABHA NO :

AGE/SEX :37 Years Male DRAWN :11/02/2024 09:30:00

RECEIVED :11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status <u>Final</u> Results Biological Reference Interval Units

Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infectionwhen present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis







Page 14 Of 17

View Details

View Report







REF. DOCTOR: SELF

PATIENT NAME: ANSHU BOBS8959

CODE/NAME & ADDRESS : C000049066

AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912

PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00

RECEIVED :11/02/2024 09:36:59 REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

CLINICAL PATH - STOOL ANALYSIS

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

PHYSICAL EXAMINATION, STOOL

COLOUR SA

METHOD: GROSS EXAMINATION

SAMPLE NOT RECEIVED

Dr. Abhishek Sharma Consultant Microbiologist



Page 15 Of 17

View Details

View Report







CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED :11/02/2024 09:36:59

REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

SPECIALISED CHEMISTRY - HORMONE

MEDI WHEEL FULL BODY HEALTH CHECK UP BELOW 40 MALE

THYROID PANEL, SERUM

ТЗ	124.87	60.0 - 181.0	ng/dL
METHOD: CHEMILUMINESCENCE T4	7.90	4.5 - 10.9	μg/dL
METHOD : CHEMILUMINESCENCE TSH (ULTRASENSITIVE)	2.910	0.550 - 4.780	μIU/mL
METHOD: CHEMILLIMINESCENCE			

Interpretation(s)

Triiodothyronine T3, Thyroxine T4, and Thyroid Stimulating Hormone TSH are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low. Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hypothyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3)
					Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1)Subclinical Hypothyroidism (2) Patient with insufficient thyroid
					hormone replacement therapy (3) In cases of Autoimmune/Hashimoto
					thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical
					inflammation, drugs like amphetamines, Iodine containing drug and
					dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre
					(3)Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid
					hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4
					replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism

Dr. Akansha Jain Consultant Pathologist



Page 16 Of 17

View Details

View Benort





CODE/NAME & ADDRESS : C000049066
AGILUS DIAGNOSTICS LIMITED-WEL WALK-IN-AAKRITI LABS PVT LTD, A-430, AGRASEN MARG

JAIPUR 302017 9314660100 ACCESSION NO: 0251XB000912 PATIENT ID : ANSHM110287251

CLIENT PATIENT ID: 012402110015

ABHA NO :

AGE/SEX : 37 Years Male DRAWN :11/02/2024 09:30:00 RECEIVED :11/02/2024 09:36:59

REPORTED :11/02/2024 15:53:12

Test Report Status Final Results Biological Reference Interval Units

6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent
					treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2.Guidlines of the American Thyroid association during pregnancy and Postpartum, 2011.

NOTE: It is advisable to detect Free T3,FreeT4 along with TSH, instead of testing for albumin bound Total T3, Total T4.TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

End Of Report
Please visit www.agilusdiagnostics.com for related Test Information for this accession

Dr. Akansha Jain Consultant Pathologist



Page 17 Of 17

View Details

View Report







3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661 www.aakritilabs.com

CIN NO.: U85195RJ2004PTC019563

PATIENT NAME: MR ANSHU BOBS AGE & SEX: 37 V/M REF. by: MEDIWHEEL DATE: 11.02.2024

USG: WHOLE ABDOMEN (Male)

LIVER

: Is enlarged in size with bright in echogenecity.

The IHBR and hepatic radicals are not dilated. No evidence of focal echopoor/echorich lesion seen.

Portal vein diameter and common bile duct appear normal.

GALL

: Is contracted (Post Prandial)

BLADDER

PANCREAS: Is normal in size, shape and echotexture. Pancreatic duct is not dilated. : Is normal in size, shape and echogenecity Spleenic hilum is not dilated.

KIDNEYS: Right Kidney:-Size: 105 x 45 mm. Left Kidney:-Size: 102 x 42 mm.

Bilateral Kidneys are normal in size, shape and echotexture, corticomedullary differentiation is fair and ratio appears normal.

Pelvi calyceal system is normal. No evidence of hydronephrosis/ nephrolithiasis.

URINARY: Bladder walls are smooth regular and normal thickness.

BLADDER: No evidence of mass or stone in bladder lumen.

PROSTATE: Is normal in size, shape and echotexture,

measures: 31 x 27 x 25 mm, wt: 11 gms.

Its capsule is intact and no evidence of focal lesion.

SPECIFIC: No evidence of retroperitoneal mass or free fluid seen in peritoneal cavity.

: NO evidence of lymphadenopathy or mass lesion in retroperitoneum.

: Visualized bowel loop appear normal Great vessels appear normal.

IMPRESSION: - Hepatomegaly with fatty changes (Grade - III)

DR NEERA MEHTA MBBS, DMRD RMCNO.005807/14853



Aakriti Labs

3 Mahatma Gandhi Marg, Gandhi Nagar Mod Tonk Road, Jaipur (Raj.) Ph.: 0141-2710661

www.aakritilabs.com CIN NO.: U85195RJ2004PTC019563



: Mr. ANSHU BOBS8959

Age/Gender: 37 Y/Male

Patient ID : 012402110015

BarcodeNo :10114575

Referred By : Self

Registration No: 75768

Registered

: 11/Feb/2024 09:30AM

Analysed Reported

: 11/Feb/2024 02:35PM

Panel

: 11/Feb/2024 02:35PM : MEDI WHEEL (ARCOFEMI

HEALTHCARE LTD)

DIGITAL X-RAY CHEST PA VIEW

Soft tissue shadow and bony cages are normal.

Trachea is central.

Bilateral lung field and both CP angle are clear.

Domes of diaphragm are normally placed.

Transverse diameter of heart appears with normal limits.

IMPRESSION:- NO OBVIOUS ABNORMALITY DETECTED.

*** End Of Report ***

Page 1 of 1




Dr. Neera Mehta M.B.B.S., D.M.R.D. RMCNO.005807/14853

ALPL policy mandates the film records to be maintained for a period of 3 months only. Kindly collect the films before this serior



MR. ANSHU /37 KIS/M/O CITS/O KG/HR 83

11-02-2024 BP: 123/72 mmHg BLC On North On HF 0:05 Hz LF 29 Hz



idhi Marg, Gandhi Nagar Mod, Tonk Road, Jaipur 302015 EMail:

III / 37 Yrs / M / 0 Cms / 0 Kg II - 02 - 2024 Refd By : BOB Examined By: No Angina / Hypercholestromia/Diabetic/Positive Estrogen/Non-Athlete



I	Elevation	METS	Rata	% THR	80	RPP	PVC
_	00.0	01.0	086	46 %	123/72	104	00
	00.0	01.0	097	53 %	123/72	119	8
	00.0	01.0	097	53%	123/72	119	8
	00.0	010	084	46 %	123/72	103	8
	00.0	01.0	084	2	123/72	103	3 :
				40.76		900	
	10,0	04.7	140	77%	123/72	172	8 8
	10.0	04.7	140	27 8 8 8	123/72	172	8 8 8
	10,0 12,0	04.7 07.1	140 169	2 2 7 8	123/72 123/72 123/72	172	8888
	10.0 12.0 14.0	04.7 07.1 07.3	140	2 2 2 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3 3	123/72 123/72 123/72 123/72	172 207 207	88888
	10.0 12.0 14.0 00.0	04.7 07.1 07.3 01.2	140 169 142	3 2 2 2 3 3	123/72 123/72 123/72 123/72 123/72	172 207 207 174	888888
	5peed(mph) 00.0 00.0 00.0	Speed(mph) Elevation 00.0 00.0 00.0 00.0 00.0 00.0 00.0	17	00.0 01.0 00.0 00.0 01.0 01.0 01.0	00.0 01.0 097 00.0 01.0 097 00.0 01.0 097 00.0 01.0 097	Elevation METs Rate % THR 00.0 .01.0 085 46 % 00.0 01.0 097 53 % 00.0 01.0 097 53 % 00.0 01.0 084 46 % 00.0 01.0 084 46 %	Elevation METs Rate % THR BP 00.0 .01.0 085 46 % 123/72 00.0 01.0 097 53 % 123/72 00.0 01.0 097 53 % 123/72 00.0 01.0 084 46 % 123/72 00.0 01.0 084 46 % 123/72

Test End Reasons Max WorkLoad Attained Initial BP (ExStrt) Initial HR (ExStrt) Exercise Time : 123/72 (mm/Hg) 7.3 Fair response to induced stress : 84 bpm 46% of Target 183 Test Complete, Heart Rate Achieved

> Max BP Attained 154/73 (mm/Hg) Max HR Attained 169 bpm 92% of Target 183

REPORT:

FINAL IMPRESSION :- TEST IS NEGATIVE FOR INDUCIBLE ISCHAEMIA

Doctor: DR

