



Visit ID : MBAR39296	Registration : 11/Mar/2023 01:28PM
UHID/MR No : ABAR.0000039284	Collected : 11/Mar/2023 01:33PM
Patient Name : Mr.SHOBHIT	Received : 11/Mar/2023 01:37PM
Age/Gender : 43 Y 0 M 0 D /M	Reported : 11/Mar/2023 02:44PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240, Sanjay Nagar Bareilly (UP)	Barcode No : A3619651

DEPARTMENT OF HORMONE ASSAYS				
Test Name	Result	Unit	Bio. Ref. Range	Method

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)				
Sample Type : SERUM				
T3	1.56	ng/ml	0.61-1.81	CLIA
T4	10.5	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	0.772	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

1. Serum T3, T4 and TSH are the measurements from three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
2. Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
3. Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
4. Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propranolol and propylthiouracil.
5. Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
6. Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
7. Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
8. TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

9. REFERENCE RANGE:

PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 - 2.500
2nd Trimester	0.200 - 3.000
3rd Trimester	0.300 - 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

1. During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
2. TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***

Miti

Dr. Miti Gupta
DNB ; MD [Pathology]



Visit ID : MBAR39297	Registration : 11/Mar/2023 01:28PM
UHID/MR No : ABAR.0000039285	Collected : 11/Mar/2023 01:33PM
Patient Name : Mrs.BHAWNA	Received : 11/Mar/2023 01:37PM
Age/Gender : 30 Y 0 M 0 D /F	Reported : 11/Mar/2023 02:44PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240, Sanjay Nagar Bareilly (UP)	Barcode No : A3619650

DEPARTMENT OF HORMONE ASSAYS				
Test Name	Result	Unit	Bio. Ref. Range	Method

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)				
Sample Type : SERUM				
T3	1.03	ng/ml	0.61-1.81	CLIA
T4	7.5	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	2.273	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

1. Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
2. Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
3. Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
4. Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propranolol and propylthiouracil.
5. Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
6. Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
7. Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
8. TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

9. REFERENCE RANGE:

PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 – 2.500
2nd Trimester	0.200 – 3.000
3rd Trimester	0.300 – 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

1. During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
2. TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***

Dr. Miti Gupta
DNB ; MD [Pathology]

Visit ID : MBAR39298	Registration : 11/Mar/2023 01:29PM
UHID/MR No : ABAR.0000039286	Collected : 11/Mar/2023 01:34PM
Patient Name : Mr.VIKRAM	Received : 11/Mar/2023 01:37PM
Age/Gender : 31 Y O M O D /M	Reported : 11/Mar/2023 02:44PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240,Sanjay Nagar Bareilly (UP)	Barcode No : A3619649

DEPARTMENT OF HORMONE ASSAYS

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)

Sample Type : SERUM

T3	1.31	ng/ml	0.61-1.81	CLIA
T4	9.2	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	2.426	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

- Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
- Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
- Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
- Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propranolol and propylthiouracil.
- Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
- Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
- Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
- TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

9. REFERENCE RANGE:


PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 - 2.500
2nd Trimester	0.200 - 3.000
3rd Trimester	0.300 - 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

- During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
- TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***



 Dr. Miti Gupta
 DNB ; MD [Pathology]


Visit ID : MBAR39299	Registration : 11/Mar/2023 01:30PM
UHID/MR No : ABAR.0000039287	Collected : 11/Mar/2023 01:34PM
Patient Name : Mrs.POOJA	Received : 11/Mar/2023 01:37PM
Age/Gender : 40 Y O M O D /F	Reported : 11/Mar/2023 02:44PM
Ref Doctor : Dr.NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240, Sanjay Nagar Bareilly (UP)	Barcode No : A3619648

DEPARTMENT OF HORMONE ASSAYS

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

THYROID PROFILE (T3,T4,ULTRASENSITIVE TSH)

Sample Type : SERUM

Test Name	Result	Unit	Bio. Ref. Range	Method
T3	0.80	ng/ml	0.61-1.81	CLIA
T4	9.4	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	7.192	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

1. Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
2. Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
3. Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
4. Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propranolol and propylthiouracil.
5. Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
6. Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
7. Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
8. TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

REFERENCE RANGE:


PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 - 2.500
2nd Trimester	0.200 - 3.000
3rd Trimester	0.300 - 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

1. During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
2. TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***



 Dr. Miti Gupta
 DNB : MD [Pathology]



Visit ID : MBAR39300	Registration : 11/Mar/2023 01:31PM
UHID/MR No : ABAR.0000039288	Collected : 11/Mar/2023 01:34PM
Patient Name : Mr. DEEPAK	Received : 11/Mar/2023 01:37PM
Age/Gender : 46 Y 0 M 0 D /M	Reported : 11/Mar/2023 02:44PM
Ref Doctor : Dr. NITIN AGARWAL	Status : Final Report
Client Name : MODERN PATH SERVICES, BAREILLY	Client Code : 2423
Client Add : 240, Sanjay Nagar Bareilly (UP)	Barcode No : A3619647

DEPARTMENT OF HORMONE ASSAYS

Test Name	Result	Unit	Bio. Ref. Range	Method
-----------	--------	------	-----------------	--------

THYROID PROFILE (T3, T4, ULTRASENSITIVE TSH)

Sample Type : SERUM				
T3	1.14	ng/ml	0.61-1.81	CLIA
T4	8.3	ug/dl	5.01-12.45	CLIA
Ultrasensitive TSH	2.275	uIU/mL	0.55-4.78	CLIA

INTERPRETATION:

1. Serum T3, T4 and TSH are the measurements form three components of thyroid screening panel and are useful in diagnosing various disorders of thyroid gland function.
2. Primary hyperthyroidism is accompanied by elevated serum T3 and T4 values along with depressed TSH levels.
3. Primary hypothyroidism is accompanied by depressed serum T3 and T4 values and elevated serum TSH levels.
4. Normal T4 levels accompanied by high T3 levels are seen in patients with T3 thyrotoxicosis. Slightly elevated T3 levels may be found in pregnancy and in estrogen therapy while depressed levels may be encountered in severe illness, malnutrition, renal failure and during therapy with drugs like propranolol and propylthiouracil.
5. Although elevated TSH levels are nearly always indicative of primary hypothyroidism, rarely they can result from TSH secreting pituitary tumors (secondary hyperthyroidism).
6. Low levels of Thyroid hormones (T3, T4 & FT3, FT4) are seen in cases of primary, secondary and tertiary hypothyroidism and sometimes in non-thyroidal illness also.
7. Increased levels are found in Grave's disease, hyperthyroidism and thyroid hormone resistance.
8. TSH levels are raised in primary hypothyroidism and are low in hyperthyroidism and secondary hypothyroidism.

9. REFERENCE RANGE:

PREGNANCY	Ultrasensitive TSH in uIU/mL
1st Trimester	0.100 - 2.500
2nd Trimester	0.200 - 3.000
3rd Trimester	0.300 - 3.000

(Reference range recommended by the American Thyroid Association)

Comments :

1. During pregnancy, Free thyroid profile (FT3, FT4 & Ultra-TSH) is recommended.
2. TSH levels are subject to circadian variation, reaches peak levels between 2-4 AM and at a minimum between 6-10 PM. The variation of the day has influence on the measured serum TSH concentrations.

*** End Of Report ***

Miti


Dr. Miti Gupta
DNB ; MD [Pathology]

10mm/mV 25mm/sec 25Hz

BPL CARDIART 610BT

10mm/mV 25mm/sec 25Hz

BPL CARDIART



Normal

डॉ. विविन अग्रवाल
11/3/23
बसु गंगु विभागा

Pat. ID. Pooja Agr

11/3/23

Pat. ID.



CARDIART

CARDIART

BPL CARDIART 6108T

10mm/mV 25mm/sec 25Hz

aVL




Pat. ID.

BPL CARDIART 6108T


BPL

aVF



10mm/mV 25mm/sec 25Hz

V1



Pat. ID.

V2



BPL

BPL 6108T

10mm/mV 25mm/sec 25Hz


V3



Pat. ID.....


BPL BPL CARDIART 6108T

V4



10mm/mV 25mm/sec 25Hz


V5



Pat. ID.....

BPL BPL CARDIART 6108T

V6



CARDIART



PARAS MRI & ULTRASOUND CENTRE

MOST ADVANCED 32 CHANNEL 3T 3D WHOLE BODY MRI

261, ASHAPURAM, OPP. DR. BASU EYE HOSPITAL, STADIUM ROAD, BAREILLY

• Helpline : 7300761761 • E-mail : parasmribly@gmail.com

REPORT

4D / 5D ULTRASOUND

COLOR DOPPLER

TVS/ TRUS

MUSCULOSKELETAL USG

Date : 11.3.2023
Name : SHOBIT AGARWAL 42Y/M
Ref.By : APPLE CARDIAC CARE

ULTRASOUND WHOLE ABDOMEN

LIVER - Liver is normal in size and outline. It shows increased echogenicity. No obvious focal pathology is seen. The intra hepatic biliary radicals are not dilated. PV -normal.

GALL BLADDER -Gall Bladder is normal in size, has normal wall thickness with no evidence of calculi. Fat planes between GB and liver are well maintained. The CBD appears normal.

PANCREAS - Pancreas is normal in size and echogenicity. Its outlines are distinct. No obvious focal lesion, calcification or ductile dilatation is seen.

SPLEEN - Spleen is normal in size and echogenicity. There is no evidence of collaterals

KIDNEYS - Both kidneys are normal in position, outline and echogenicity. No evidence of calculi is seen. CMD is maintained. No evidence of hydronephrosis is seen on either side.

URINARY BLADDER -Urinary Bladder is normal in size and outline. There is no evidence of any obvious intraluminal or paramedical pathology. **Wall is not thickened. Both VUJ clear.**

PROSTATE- Normal in size and echotexture.

No evidence of ascites/pleural effusion/ adenopathy is seen. Bowel loops are not dilated. Bilateral iliac fossa appears normal

IMPRESSION:



❖ Grade I fatty liver.

Adv- clinical correlation.

Dr. Puja Tripathi

M.B.B.S., M.D.

MBBS, MD (Radiodiagnosis, SGPGI)




NAME	Mr. SHOBHIT AGARWAL	AGE/SEX	42 Y/M
Reff. By	Dr. NITIN AGARWAL (DM)	DATE	11/03/2023

ECHOCARDIOGRAPHY AND COLOUR DOPPLER STUDY

<u>MEASUREMENTS</u>	<u>VALUE</u>	<u>NORMAL DIMENSIONS</u>
LVID (d)	4.5	cm (3.7 –5.6 cm)
LVID (s)	2.5	cm (2.2 –3.9 cm)
RVID (d)	2.4	cm (0.7 –2.5 cm)
IVS (ed)	1.0	cm (0.6 –1.1 cm)
LVPW (ed)	1.0	cm (0.6 –1.1 cm)
AO	2.0	cm (2.2 –3.7 cm)
LA	3.2	cm (1.9 –4.0 cm)
<u>LV FUNCTION</u>		
EF	60	% (54 –76 %)
FS	30	% (25 –44 %)

- LEFT VENTRICLE : No regional wall motion abnormality
 No concentric left Ventricle Hypertrophy
- MITRAL VALVE : Thin, PML moves posteriorly during Diastole
 No SAM, No Subvalvular pathology seen.
 No mitral valve prolapse calcification .
- TRICUSPID VALVE : Thin, opening wells. No calcification, No doming .
 No Prolapse.
 Tricuspid inflow velocity= 0.7 m/sec
- AORTIC VALVE : Thin, tricuspid, opening well, central closer,
 no flutter.
 No calcification
 Aortic velocity = 1.3 m/sec
- PULMONARY VALVE : Thin, opening well, Pulmonary artery is normal
 EF slope is normal.
 Pulmonary Velocity = 0.9 m/sec



ON DOPPLER INTERROGATION THERE WAS :

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

MITRAL FLOW

E= 0.8 m/sec

A= 0.6 m/sec

ON COLOUR FLOW:

- No mitral regurgitation
- No tricuspid regurgitation
- No aortic regurgitation
- No pulmonary regurgitation

COMMENTS:

- No LA /LV clot
- No pericardial effusion
- No intracardiac mass
- IAS/IVS Intact
- Inferior vena cava – normal in size with normal respiratory variation

FINAL IMPRESSION


- NO REGIONAL WALL MOTION ABNORMALITY
- NORMAL LV DIASTOLIC FUNCTION
- NORMAL LV SYSTOLIC FUNCTION (LVEF~60%)
- NORMAL CARDIAC CHAMBER DIMENSIONS
- NORMAL VALVULAR COLOUR FLOW PATTERN

**DR.NITIN AGARWAL
DM (Cardiology)
Consultant Cardiologist**

This opinion is to be correlated with the clinically findings and if required, please re-evaluate / reconfirm with further investigation.

11 MAR 2023 03:05pm


B F P G 32K CFM F 2.5 MHz G 40K
TEI D 17 CM XY C PRF 4.2KHZ
PRC 6-S-H PRS 2 PRC 2-L-H PRS 3
PST 1 MF H



11 MAR 2023 03:05pm

B F P G 40K
TEI D 17 CM XY C
PRC 6-S-L PRS A
PST 1 SV 4-112mm


PW F 2.5 MHz G 64K
PRF 4.8KHZ
PRC 6-1
PST 2 MF 300 Hz



11 MAR 2023 03:05pm


B F P G 40K
TEI D 17 CM XY C
PRC 6-S-H PRS A
PST 1

M G 40K
PRC 7-3
PST 2



11 MAR 2023 03:05pm


B F P G 40K
TEI D 17 CM XY C
PRC 6-S-L PRS A
PST 1



11 MAR 2023 03:05pm

B F P G 50K
TEI D 17 CM XY C
PRC 6-S-L PRS A
PST 1 SV 4-81mm

PW F 2.5 MHz G 64K
PRF 6.7KHZ
PRC 6-1
PST 2 MF 300 Hz





॥ ॐ गणेशाय नमः ॥

GANESH DIAGNOSTIC

DR. LOKESH GOYAL

MBBS (KGMC), MD (RADIOLOGY)

CONSULTANT INTERVENTIONAL RADIOLOGIST
FORMER SR. REGISTRAR - APOLLO HOSPITAL, NEW DELHI
LIFE MEMBER OF IRIA

Timings : 9:00 am to 9:00 pm, Sunday 9.00 am to 3.00 pm ☎ 8392957683, 6395228718

MR. SHOBHIT AGARWAL
DR. NITIN AGARWAL, DM

11-03-2023

REPORT

EXAMINATION PERFORMED: X-RAY CHEST

B/L lung fields are clear

Both of the CP angles are clear.

Both hila show a normal pattern .

Cardiac and mediastinal borders appear normal.

Visualized bony thorax and soft tissue of the chest wall appear normal.

IMPRESSION ---NO SIGNIFICANT ABNORMALITY IS SEEN


Not for medico-legal purpose

DR LOKESH GOYAL
MD
RADIOLOGIST

डिजिटल एक्स-रे, मल्टी स्लाइस
सी. टी. स्कैन सुविधा उपलब्ध है।



NOT VALID FOR
MEDICO LEGAL PURPOSE





Reg.NO. : 113
 NAME : **Mr. SHOBHIT AGARWAL**
 REFERRED BY : Dr.Nitin Agarwai (D M)
 SAMPLE : BLOOD

DATE : **11/03/2023**
 AGE : 43 Yrs.
 SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
BLOOD UREA	27	mg/dL.	10-40
<p>* Low serum urea is usually associated with status of overhydration severe hepatic failure. * A urea level of 10-45 mg/dl indicates normal glomerular function and a level of 100-250 mg/dl indicates a serious impairment of renal function. In chronic renal failure , urea correlates better with the symptoms of uremia than does serum creatinine. * Urine/Serum urea is more than 9 in prerenal and less than 3 in renal uremia.</p>			
SERUM CREATININE	0.9	mg/dL.	0.5-1.4
URIC ACID	7.0	mg/dl	3.5-8.0

CLINICAL SIGNIFICANCE:

Analysis of synovial fluid plays a major role in the diagnosis of joint disease.

SERUM SODIUM (Na)	141	m Eq/litre.	135 - 155
SERUM POTASSIUM (K)	4.0	m Eq/litre.	3.5 - 5.5
SERUM CALCIUM	9.4	mg/dl	8.5 - 10.5

Report is not valid for medicolegal purpose



Reg.NO. : 113
NAME : **Mr. SHOBHIT AGARWAL**
REFERRED BY : Dr.Nitin Agarwal (D M)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 43 Yrs.
SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
LIPID PROFILE			
SERUM CHOLESTEROL	197	mg/dL.	130 200
SERUM TRIGLYCERIDE	168	mg/dl.	30 - 160
HDL CHOLESTEROL	47	mg/dL.	30-70
VLDL CHOLESTEROL	33.6	mg/dL.	15 - 40
LDL CHOLESTEROL	106.40	mg/dL.	00-130
CHOL/HDL CHOLESTEROL RATIO	3.98	mg/dl	
LDL/HDL CHOLESTEROL RATIO	2.26	mg/dl	

INTERPRETATION

TRIGLYCERIDE level > 250mg/dL is associated with an approximately 2-fold greater risk of coronary vascular disease. Elevation of triglycerides can be seen with obesity, medication, fast less than 12 hrs., alcohol intake, diabetes melitus, and pancreatitis. CHOLESTEROL, its fractions and triglycerides are the important plasma lipids in defining cardiovascular risk factors and in the management of cardiovascular disease. Highest acceptable and optimum values of cholesterol vary with age. Values above 220 mgm/dl are associated with increased risk of CHD regardless of HDL & LDL values. HDL-CHOLESTEROL level <35 mg/dL is associated with an increased risk of coronary vascular disease even in the face of desirable levels of cholesterol and LDL - cholesterol. LDL - CHOLESTEROL & TOTAL CHOLESTEROL levels can be strikingly altered by thyroid, renal and liver disease as well as hereditary factors. Based on total cholesterol, LDL- cholesterol, and total cholesterol/HDL - cholesterol ratio, patients may be divided into the three risk categories.

Report is not valid for medicolegal purpose

Reg.NO. : 113
 NAME : **Mr. SHOBHIT AGARWAL**
 REFERRED BY : Dr. Nitin Agarwal (D.M.)
 SAMPLE : BLOOD

DATE : **11/03/2023**
 AGE : 43 Yrs.
 SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
LIVER PROFILE			
SERUM BILIRUBIN			
TOTAL	0.8	mg/dL	0.3-1.2
DIRECT	0.5	mg/dL	0.2-0.6
INDIRECT	0.3	mg/dL	0.1-0.4
SERUM PROTEINS			
Total Proteins	7.0	Gm/dL	6.4 - 8.3
Albumin	4.2	Gm/dL	3.5 - 5.5
Globulin	2.8	Gm/dL	2.3 - 3.5
A : G Ratio	1.5		0.0-2.0
SGOT	38	IU/L	0-40
SGPT	33	IU/L	0-40
SERUM ALK.PHOSPHATASE	67	IU/L	00-115

NORMAL RANGE : BILIRUBIN TOTAL

Premature infants. 0 to 1 day: <8 mg/dL Premature infants. 1 to 2 days: <12 mg/dL Adults: 0.3-1 mg/dL.
 Premature infants. 3 to 5 days: <16 mg/dL Neonates, 0 to 1 day: 1.4-8.7 mg/dL
 Neonates, 1 to 2 days: 3.4-11.5 mg/dL Neonates, 3 to 5 days: 1.5-12 mg/dL Children 6 days to 18 years: 0.3-1.2 mg/dL

COMMENTS-

Total and direct bilirubin determination in serum is used for the diagnosis, differentiation and follow-up of jaundice. Elevation of SGPT is found in liver and kidney diseases such as infectious or toxic hepatitis, LM and cirrhosis. Organs rich in SGOT are heart, liver and skeletal muscles. When any of these organs are damaged, the serum SGOT level rises in proportion to the severity of damage. Elevation of Alkaline Phosphatase in serum or plasma is found in hepatitis, biliary obstructions, hyperparathyroidism, steatorrhea and bone diseases.

URINE EXAMINATION

Report is not valid for medicolegal purpose

Reg.NO. : 113
 NAME : **Mr. SHOBHIT AGARWAL**
 REFERRED BY : Dr. Nitin Agarwal (D.M.)
 SAMPLE : BLOOD

DATE : **11/03/2023**
 AGE : 43 Yrs.
 SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
URINE EXAMINATION REPORT			
PHYSICAL EXAMINATION			
pH	6.0		
TRANSPARENCY			
Volume	25	ml	
Colour	Light Yellow		
Appearance	Clear		Nil
Sediments	Nil		
Specific Gravity	1.020		1.015-1.025
Reaction	Acidic		
BIOCHEMICAL EXAMINATION			
UROBILINOGEN	Nil		NIL
BILIRUBIN	Nil		NEGATIVE
URINE KETONE	Nil		NEGATIVE
Sugar	Nil		Nil
Albumin	Nil		Nil
Phosphates	Absent		Nil
MICROSCOPIC EXAMINATION			
Red Blood Cells	Nil	/H.P.F.	
Pus Cells	1-2	/H.P.F.	
Epithelial Cells	1-2	/H.P.F.	
Crystals	NIL		NIL
Casts	NIL	/H.P.F.	
Bacteria	NIL		
Other	NIL		

BIOCHEMICAL

Report is not valid for medicolegal purpose

Reg.NO. : 113
NAME : **Mr. SHOBHIT AGARWAL**
REFERRED BY : Dr.Nitin Agarwal (D.M)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 43 Yrs.
SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
Prostatic Specific Antigen	2.7	ng/ml	0-4

Prostatic Specific Antigen (P.S.A)


Comment : The fact of PSA is unique to prostate tissue makes it a suitable marker for monitoring men with cancer of the prostate. PSA is also useful for determining possible recurrence after therapy. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy.

* Quality controlled report with external quality assurance

BIOCHEMISTRY

Gamma Glutamyl Transferase (GGT)	20	U/L	7-32
----------------------------------	----	-----	------

--{End of Report}--



Dr. Shweta Agarwal
MD(Pathology), Apple Pathology
Bareilly (UP)

Report is not valid for medicolegal purpose

Department of Apple Cardiac Care
Ekta Nagar, Stadium Road,
Apple Care Hospital,
Bareilly - 243 122 (U.P.) India
Tel. : 07599031977, 09458888448



Reg.NO. : 113
NAME : **Mr. SHOBHIT AGARWAL**
REFERRED BY : Dr.Nitin Agarwal (D M)
SAMPLE : BLOOD

DATE : **11/03/2023**
AGE : 43 Yrs.
SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
------------------	----------------	--------------	------------------------------

HAEMATOLOGY

GLYCOSYLATED HAEMOGLOBIN

5.7

EXPECTED RESULTS :

Non diabetic patients : 4.0% to 6.0%
Good Control : 6.0% to 7.0%
Fair Control : 7.0% to -8%
Poor Control : Above 8%

***ADA: American Diabetes Association**

The glycosylated hemoglobin assay has been validated as a reliable indicator of mean blood glucose levels for a period of 8-12 week period prior to HBA1C determination. ADA recommends the testing twice a year in patients with stable blood glucose, and quarterly, if treatment changes, or if blood glucose levels are unstable.

METHOD : ADVANCED IMMUNO ASSAY.

--{End of Report}--

Dr. Shweta Agarwal
MD(Pathology), Apple Pathology
Bareilly (UP)

Report is not valid for medicolegal purpose


Lab. Timings : 9.00 a.m. to 8.00 p.m. Sunday : 10.00 a.m. to 2.00 p.m.
Home Sample Collection Facility Available

Reg. NO. : 113
NAME : Mr. SHOBHIT AGARWAL
REFERRED BY : Dr. Nitin Agarwal (D.M.)
SAMPLE : BLOOD

DATE : 11/03/2023
AGE : 43 Yrs.
SEX : MALE

<u>TEST NAME</u>	<u>RESULTS</u>	<u>UNITS</u>	<u>BIOLOGICAL REF. RANGE</u>
BLOOD SUGAR P.P.	BIOCHEMISTRY 131	mg/dl	80-140

--(End of Report)--


Dr. Shweta Agarwal
MD(Pathology), Apple Pathology
Bareilly (UP)

Report is not valid for medicolegal purpose

Dr. Nitin Agarwal

MD, DM (Cardiology)
Consultant Interventional Cardiologist
Cell : +91-94578 33777

Formerly at :
Escorts Heart Institute & Research Centre, Delhi
Dr. Ram Manohar Lohia Hospital, Delhi



APPLE
CARDIAC CARE
DR. NITIN AGARWAL'S HEART CLINIC

Shobhit
Agarwal
42/117

1113/23

120/30

461

22

Aspirin

Q

Exerc

Life style

Aspirin

Q

42 - 106

A-3, EKTA NAGAR, (OPP. CARE HOSPITAL) STADIUM ROAD, NEAR DELAPEER CHAURAHA, BAREILLY - 243 122 (U.P.)

OPD Timings : 12.00 Noon to 04.00 pm, Sunday : 12.00 Noon to 3.00 pm

नम्बर लगाने के लिए फोन करें : 09458888448, 07599031977

VALID FOR 5 DAYS.


पर्चा पाँच दिन के लिये मान्य

10mm/mV 25mm/sec 25Hz

BPL CARDIART 610BT

10mm/mV 25mm/sec 25Hz

BPL CARDIART



Normal

डॉ. विजय कुमार
 25/3/23
 बसु गंगु विद्यालय

Pat. ID. Pooja Aoo

11/3/23

Pat. ID.



CARDIART

CARDIART

BPL CARDIART 6108T

10mm/mV 25mm/sec 25Hz

aVL





Pat. ID.

BPL CARDIART 6108T

BPL


aVF



Pat. ID.

10mm/mV 25mm/sec 25Hz

V1



V2




BPL

BPL 6108T

10mm/mV 25mm/sec 25Hz


V3



Pat. ID.....


BPL BPL CARDIART 6108T

V4



10mm/mV 25mm/sec 25Hz


V5



Pat. ID.....

BPL BPL CARDIART 6108T

V6



CARDIART

10mm/mV 25mm/sec 25Hz

BPL CARDIART 6108T

10mm/mV 25mm/sec 25Hz

BPL CARDIART 6108T

10mm/mV 25mm/sec 25Hz




I

II

III

aVR

aVL



Pat. ID. Shobhit


11/3/23

Normal
Dr. Nitin Sharma
11/3/23

Pat. ID.

Pat. ID.

CARDIART



61087

10mm/mV 25mm/sec \approx 25Hz

BPL CARDIART 6108T

10mm/mV 25mm/sec \approx 25Hz


BPL CARDIART 6108T

V3

V4

V5

V6



Pat. ID.....

Pat. ID.....

REPORT