PID No. : MED111034817

: 422024610

: MediWheel

Age / Sex : 30 Year(s) / Female

: OP

SID No.

Type

Ref. Dr

Register On : 26/03/2022 10:41 AM

Collection On : 26/03/2022 12:01 PM

Report On : 27/03/2022 5:20 PM

Printed On : 30/03/2022 2:33 PM



Investigation HAEMATOLOGY	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood'Spectrophotometry)	11.6	g/dL	12.5 - 16.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood/Derived from Impedance)	37.3	%	37 - 47
RBC Count (EDTA Blood/Impedance Variation)	4.46	mill/cu.mm	4.2 - 5.4
Mean Corpuscular Volume(MCV) (EDTA Blood/Derived from Impedance)	84.0	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood/Derived from Impedance)	26.0	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood/Derived from Impedance)	31.0	g/dL	32 - 36
RDW-CV (EDTA Blood/Derived from Impedance)	15.2	%	11.5 - 16.0
RDW-SD (EDTA Blood/Derived from Impedance)	44.69	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood/Impedance Variation)	6200	cells/cu.mm	4000 - 11000
Neutrophils (EDTA Blood/Impedance Variation & Flow Cytometry)	50.0	%	40 - 75
Lymphocytes	39.1	%	20 - 45



Cytometry)

(EDTA Blood/Impedance Variation & Flow

DR SHAMIM JAVED MD PATHOLOGY KMC-88902

PID No. : MED111034817

: 422024610

Age / Sex : 30 Year(s) / Female

Type : OP

SID No.

Ref. Dr : MediWheel

Register On : 26/03/2022 10:41 AM

Collection On : 26/03/2022 12:01 PM

Report On : 27/03/2022 5:20 PM

Printed On : 30/03/2022 2:33 PM



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Eosinophils (EDTA Blood/Impedance Variation & Flow Cytometry)	4.5	%	01 - 06
Monocytes (EDTA Blood/Impedance Variation & Flow Cytometry)	5.5	%	01 - 10
Basophils (Blood/Impedance Variation & Flow Cytometry)	0.9	%	00 - 02
Absolute Neutrophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	3.10	10^3 / μl	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	2.42	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood/Impedance Variation & Flow Cytometry)	0.28	10^3 / μ1	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.34	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood/Impedance Variation & Flow Cytometry)	0.06	10^3 / μl	< 0.2
Platelet Count (EDTA Blood/Impedance Variation)	326	10^3 / μl	150 - 450
MPV (EDTA Blood/Derived from Impedance)	8.9	fL	8.0 - 13.3
PCT (EDTA Blood/Automated Blood cell Counter)	0.29	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood/Modified Westergren)	7	mm/hr	< 20





PID No. : MED111034817 **Register On** : 26/03/2022 10:41 AM : 422024610 SID No. Collection On : 26/03/2022 12:01 PM Age / Sex : 30 Year(s) / Female

Report On

Printed On

Type : OP

Ref. Dr : MediWheel

Investigation	<u>Observed</u> Value	<u>Unit</u>	<u>Biological</u> Reference Interval
BIOCHEMISTRY	<u> </u>		<u></u>

: 27/03/2022 5:20 PM

: 30/03/2022 2:33 PM

Liver Function Test

0.3	mg/dL	0.1 - 1.2
0.1	mg/dL	0.0 - 0.3
0.2	mg/dL	0.1 - 1.0
7.1	gm/dL	6.0 - 8.0
4.3	gm/dL	3.5 - 5.2
2.8	gm/dL	2.3 - 3.6
1.5		1.1 - 2.2
17	U/L	5 - 40
8	U/L	5 - 41
55	U/L	42 - 98
13	U/L	< 38
	0.1 0.2 7.1 4.3 2.8 1.5 17 8	0.1 mg/dL 0.2 mg/dL 7.1 gm/dL 4.3 gm/dL 2.8 gm/dL 1.5 17 U/L 8 U/L 55 U/L



VERIFIED BY



Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/Cholesterol oxidase/Peroxidase)	167	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/Glycerol phosphate oxidase / peroxidase)	71	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

: 30/03/2022 2:33 PM

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the "usual" circulating level of triglycerides during most part of the day.

HDL Cholesterol (Serum/Immunoinhibition)	54	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 50 - 59 High Risk: < 50
LDL Cholesterol (Serum/Calculated)	98.8	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	14.2	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	113.0	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



VERIFIED BY



PID No. : MED111034817 Register On : 26/03/2022 10:41 AM : 422024610 SID No. Collection On : 26/03/2022 12:01 PM

Age / Sex : 30 Year(s) / Female Report On 27/03/2022 5:20 PM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 30/03/2022 2:33 PM

Total Cholesterol/HDL Cholesterol Ratio 3.1 Optimal: < 3.3Low Risk: 3.4 - 4.4 (Serum/Calculated)

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 1.3 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0 High Risk: > 5.0

LDL/HDL Cholesterol Ratio 1.8 Optimal: 0.5 - 3.0 Borderline: 3.1 - 6.0 (Serum/Calculated)

High Risk: > 6.0



(Serum/Calculated)

VERIFIED BY



PID No. : MED111034817 Register On : 26/03/2022 10:41 AM

Printed On

Type : OP

SID No.

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.8	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

: 30/03/2022 2:33 PM

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 119.76 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbAlc.



VERIFIED BY



Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Ur</u>	<u>Biological</u>
	Value	Reference Interval

: 30/03/2022 2:33 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.27 ng/mL 0.7 - 2.04

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is Metabolically active.

T4 (Thyroxine) - Total 8.64 μg/dL 4.2 - 12.0

(Serum/CMIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

TSH (Thyroid Stimulating Hormone) 2.75 µIU/mL 0.35 - 5.50

(Serum/Chemiluminescent Microparticle

Immuno as say (CMIA))

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester : 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values&lt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



PID No. : MED111034817 Register On : 26/03/2022 10:41 AM : 422024610 Collection On : 26/03/2022 12:01 PM SID No. Age / Sex : 30 Year(s) / Female

Report On

Printed On



Type : OP

Ref. Dr : MediWheel

<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

: 27/03/2022 5:20 PM

: 30/03/2022 2:33 PM

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION

Colour (Urine)	Pale Yellow		
Appearance (Urine)	Clear		Clear
Volume (Urine)	20	mL	
<u>CHEMICAL EXAMINATION(Automated-Urineanalyser)</u>			
pH (Urine/AUTOMATED URINANALYSER)	5.0		4.5 - 8.0
Specific Gravity (Urine)	1.025		1.002 - 1.035
Ketones (Urine)	Negative		Negative
Urobilinogen (Urine/AUTOMATED URINANALYSER)	0.2		0.2 - 1.0
Blood (Urine/AUTOMATED URINANALYSER)	Negative		Negative

Negative Nitrite $(Urine/A\,UTOMATED\,\,URINANALYSER)$ Negative Bilirubin (Urine/AUTOMATED URINANALYSER)

Negative

Negative

Negative Protein Negative (Urine)

Consultant Microbiologist

VERIFIED BY



 PID No.
 : MED111034817
 Register On
 : 26/03/2022 10:41 AM

 SID No.
 : 422024610
 Collection On
 : 26/03/2022 12:01 PM

 Age / Sex
 : 30 Year(s) / Female
 Report On
 : 27/03/2022 5:20 PM

Printed On



Type : OP

Ref. Dr : MediWheel

: 30/03/2022 2:33 PM

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glucose (Urine)	Negative		Negative
Leukocytes (Urine)	Negative	leuco/uL	Negative
MICROSCOPY(URINE DEPOSITS)			
Pus Cells (Urine/Flow cytometry)	3-5	/hpf	3-5
Epithelial Cells (Urine)	1-2	/hpf	1-2
RBCs (Urine/Flow cytometry)	Nil	/hpf	NIL
Others (Urine)	Nil		Nil
Casts (Urine/Flow cytometry)	Nil	/hpf	0 - 1
Crystals (Urine)	Nil		NIL



VERIFIED BY



PID No. Register On : MED111034817 : 26/03/2022 10:41 AM

: 422024610 Collection On : 26/03/2022 12:01 PM

Printed On

Age / Sex : 30 Year(s) / Female Report On : 27/03/2022 5:20 PM

: OP

SID No.

Type

Ref. Dr : MediWheel



Investigation <u>Observed</u> <u>Unit</u> **Biological** <u>Value</u> Reference Interval

: 30/03/2022 2:33 PM

IMMUNOHAEMATOLOGY

'A' 'Negative' BLOOD GROUPING AND Rh TYPING

(EDTA Blood/Agglutination)



VERIFIED BY



 PID No.
 : MED111034817
 Register On
 : 26/03/2022 10:41 AM

 SID No.
 : 422024610
 Collection On
 : 26/03/2022 12:01 PM

Printed On

Type : OP

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
BIOCHEMISTRY			
BUN / Creatinine Ratio	15.7		6 - 22
Glucose Fasting (FBS) (Plasma - F/GOD - POD)	79	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126

30/03/2022 2:33 PM

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

Glucose Fasting - Urine	Negative		Negative
(Urine - F)			
Glucose Postprandial (PPBS)	96	mg/dL	70 - 140

(Plasma - PP/GOD - POD)

INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Glucose Postprandial - Urine (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease-GLDH)	11	mg/dL	7.0 - 21
Creatinine (Serum/Jaffe Kinetic)	0.7	mg/dL	0.6 - 1.1

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists,N-acetylcyteine , chemotherapeutic agent such as flucytosine etc.

Uric Acid 4.2 mg/dL 2.6 - 6.0

(Serum/Uricase/Peroxidase)



VERIFIED BY



APPROVED BY

-- End of Report --

Name	MEENAKSHI S	ID	MED111034817
Age & Gender	30/FeMale		26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2 D ECHOCARDIOGRAPHIC STUDY

M mode measurement:

AORTA : 2.0cms

LEFT ATRIUM : 2.9cms

AVS :----

LEFT VENTRICLE (DIASTOLE) : 3.3cms

(SYSTOLE) : 2.2cms

VENTRICULAR SEPTUM (DIASTOLE) : 1.0cms

(SYSTOLE) : 1.3cms

POSTERIOR WALL (DIASTOLE) : 1.1cms

(SYSTOLE) : 1.3cms

EDV: 45ml

ESV : 15ml

FRACTIONAL SHORTENING : 35%

EJECTION FRACTION : 65%

EPSS :---

RVID : 1.7cms

DOPPLER MEASUREMENTS:

MITRAL VALVE : E' 0.60 m/s A' 0.49 m/s NO MR

AORTIC VALVE : 1.09 m/s NO AR

TRICUSPID VALVE : E' 1.77 m/s A' - m/s NO TR

PULMONARY VALVE : 0.65 m/s NO PR

Name	MEENAKSHI S	ID	MED111034817
Age & Gender	30/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



2D ECHOCARDIOGRAPHY FINDINGS:

Left ventricle : Normal size, Normal systolic function.

No regional wall motion abnormalities.

Left Atrium : Normal.

Right Ventricle : Normal.

Right Atrium : Normal.

Mitral valve : Normal, No mitral valve prolapsed.

Aortic valve : Normal, Trileaflet.

Tricuspid valve : Normal.

Pulmonary valve : Normal.

IAS : Intact.

IVS : Intact.

Pericardium : No pericardial effusion.

IMPRESSION:

- > NORMAL SIZED CARDIAC CHAMBERS.
- > NORMAL LV SYSTOLIC FUNCTION. EF: 65%.
- > NO REGIONAL WALL MOTION ABNORMALITIES.
- > NORMAL VALVES.
- > NO CLOTS / PERICARDIAL EFFUSION / VEGETATION.

DR. K.S. SUBRAMANI. MBBS, MD, DM (CARDIOLOGY) FESC SENIOR CONSULTANT INTERVENTIONAL CARDIOLOGIST

Name	MEENAKSHI S	ID	MED111034817
Age & Gender	30/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



Kss/an

- Note:

 * Report to be interpreted by qualified medical professional.

 * To be correlated with other clinical findings.
- * Parameters may be subjected to inter and intra observer variations.

Name	MEENAKSHI S	ID	MED111034817
Age & Gender	30/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER shows normal in shape, size and has uniform echopattern. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER shows normal shape and has clear contents. Gall bladder wall is of normal thickness. CBD is of normal calibre.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

No demonstrable Para -aortic lymphadenopathy.

KIDNEYS move well with respiration and have normal shape, size and echopattern. Cortico- medullary differentiations are well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney	9.4	1.1
Left Kidney	10.4	0.7

URINARY BLADDER shows normal shape and wall thickness. It has clear contents. No evidence of diverticula.

UTERUS is anteverted and normal in size. It has uniform myometrial echopattern.

Endometrial thickness measures 6mm

Uterus measures as follows: LS: 8.7cms AP: 3.4cms TS: 5.0cms.

OVARIES are normal in size, shape and echotexture. No focal lesion seen.

Ovaries measure as follows: **Right ovary**: 2.8 x 2.0cms **Left ovary**: 3.0 x 1.6cms

POD & adnexae are free.

No evidence of ascites/pleural effusion.

IMPRESSION:

> NO SIGNIFICANT ABNORMALITY DETECTED.

DR. H.K. ANAND

DR. APARNA

Name	MEENAKSHI S	ID	MED111034817
Age & Gender	30/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel	·	,



CONSULTANT RADIOLOGISTS:

A/da

Name	MEENAKSHI S	ID	MED111034817
Age & Gender	30/FeMale	Visit Date	26-03-2022 00:00:00
Ref Doctor Name	MediWheel		



X - RAY CHEST PA VIEW

Bilateral lung fields appear normal.

Cardiac size is within normal limits.

Bilateral hilar regions appear normal.

Bilateral domes of diaphragm and costophrenic angles are normal.

Visualised bones and soft tissues appear normal.

Impression: Essentially normal study.