



DIAGNOSTIC REPORT

CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

PATIENT NAME : RIKKEN ETTE

ACCESSION NO : 0321VC002554	AGE : 33 Years SEX : Female	
DRAWN :	RECEIVED : 26-03-2022 11:46	REPORTED : 01-04-2022 13:44
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :

 Test Report Status
 Final
 Results
 Biological Reference Interval
 Units


MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

BLOOD COUNTS, EDTA WHOLE BLOOD					
HEMOGLOBIN	10.7	Low	12.0 - 15.0	g/dL	
RED BLOOD CELL COUNT	4.37		3.8 - 4.8	mil/µL	
WHITE BLOOD CELL COUNT	9.59		4.0 - 10.0	thou/µL	
PLATELET COUNT	414	High	150 - 410	thou/µL	
RBC AND PLATELET INDICES					
HEMATOCRIT	32.7	Low	36.0 - 46.0	%	
MEAN CORPUSCULAR VOL	74.8	Low	83.0 - 101.0	fL	
MEAN CORPUSCULAR HGB.	24.6	Low	27.0 - 32.0	pg	
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION	32.8		31.5 - 34.5	g/dL	
MENTZER INDEX	17.1				
RED CELL DISTRIBUTION WIDTH	16.6	High	11.6 - 14.0	%	
MEAN PLATELET VOLUME	8.2		6.8 - 10.9	fL	
WBC DIFFERENTIAL COUNT - NLR					
SEGMENTED NEUTROPHILS	60		40 - 80	%	
LYMPHOCYTES	32		20 - 40	%	
EOSINOPHILS	01		1.0 - 6.0	%	
MONOCYTES	07		2.0 - 10.0	%	
BASOPHILS	00		0 - 1	%	
DIFFERENTIAL COUNT PERFORMED ON:	EDTA SMEAR				
MORPHOLOGY					
RBC	HYPOCHROMIC MIC	ROCYTI	C RBCS SEEN.		
WBC	NORMAL MORPHOLOGY				
PLATELETS	ADEQUATE				
REMARKS	NO PREMATURE CE DETECTED.	lls af	RE SEEN. MALARIAL PARASITE N	ОТ	
ERYTHRO SEDIMENTATION RATE, BLOOD					

SEDIMENTATION RATE (ESR)	30	High 0 - 20	mm at 1 hr
GLUCOSE, FASTING, PLASMA			
GLUCOSE, FASTING, PLASMA	99	74 - 99	mg/dL
GLYCOSYLATED HEMOGLOBIN, EDTA W	HOLE BLOOD		







Patient Ref. No. 777000001890805



DIAGNOSTIC REPORT









RIKKF070788321

CLIENT CODE : C000138364

CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
GRAND MALL, OPPOSITE SBI ZONAL OFFICE, SM ROAD, AMBAWADI,
AHMEDABAD, 380015
GUJRAT, INDIA
Tel: 079-48912999,079-48913999,079-48914999
Email : customercare.ahmedabad@srl.in

PATIENT ID :

REPORTED : 01-04-2022 13:44

CLIENT PATIENT ID :

PATIENT NAME : RIKKEN ETTE


ACCESSION NO :	0321VC002554	AGE :	33 Years	SEX : Female
DRAWN :		RECE	IVED : 26-03	-2022 11:46

REFERRING DOCTOR : SELF

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units	
ALANINE AMINOTRANSFERASE (ALT/SGPT)	17	0 - 33 U/L	
ALKALINE PHOSPHATASE	86	35 - 104 U/L	
GAMMA GLUTAMYL TRANSFERASE (GGT)	25	5 - 36 U/L	
LACTATE DEHYDROGENASE	136	135 - 214 U/L	
SERUM BLOOD UREA NITROGEN	150	155 211 0/2	
BLOOD UREA NITROGEN	10	6 - 20 mg/dL	
CREATININE, SERUM	10		
CREATININE	0.78	0.60 - 1.10 mg/dL	
BUN/CREAT RATIO	0.70		
BUN/CREAT RATIO	12.82	5.0 - 15.0	
URIC ACID, SERUM	12.02	5.0 15.0	
URIC ACID	5.5	2.4 - 5.7 mg/dL	
ELECTROLYTES (NA/K/CL), SERUM	5.5	2. 4 5.7 mg/dL	
SODIUM	139.6	136- 145 mmol/L	
POTASSIUM	4.18	3.50- 5.10 mmol/L	
CHLORIDE	104.5	98 - 107 mmol/L	
URINALYSIS	104.5	96 - 107 minor/L	
COLOR	Yellow		
APPEARANCE	Clear		
PH	6.0	4.7 - 7.5	
SPECIFIC GRAVITY	1.025	1.003 - 1.035	
GLUCOSE	NOT DETECTED	NOT DETECTED	
PROTEIN	NOT DETECTED	NOT DETECTED	
KETONES	NOT DETECTED	NOT DETECTED	
BLOOD	NOT DETECTED	NOT DETECTED	
BILIRUBIN	NOT DETECTED	NOT DETECTED	
UROBILINOGEN	NORMAL	NORMAL	
	NOT DETECTED	NOT DETECTED	
PUS CELL (WBC'S)	1-2	0-5 /HPF	
EPITHELIAL CELLS	3-5	0-5 /HPF	
ERYTHROCYTES (RBC'S)	NOT DETECTED	NOT DETECTED /HPF	
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED	NOT DETECTED	
BACTERIA	NOT DETECTED	NOT DETECTED	













Patient Ref. No. 777000001890805



RIKKF070788321

CLIENT CODE : C000138364

CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA DELHI INDIA 8800465156

SRL LTD
GRAND MALL, OPPOSITE SBI ZONAL OFFICE, SM ROAD, AMBAWADI,
AHMEDABAD, 380015
GUJRAT, INDIA
Tel : 079-48912999,079-48913999,079-48914999
Email : customercare.ahmedabad@srl.in

PATIENT ID :

01-04-2022 13:44

PATIENT NAME : RIKKEN ETTE

ACCESSION NO :	0321VC002554	AGE :	33 Years	SEX : Female		
DRAWN :		RECE	IVED : 26-03	3-2022 11:46	REPORTED :	01-04-2022
REFERRING DOCT	OR: SELF				CLIEN	T PATIENT ID :

Test Report Status <u>Final</u>	Results	Biological Reference Interval	Units
GENERAL APPEARANCE / NUTRITIONAL STATUS	OBESE		
BUILT / SKELETAL FRAMEWORK	AVERAGE		
FACIAL APPEARANCE	NORMAL		
SKIN	NORMAL		
UPPER LIMB	NORMAL		
LOWER LIMB	NORMAL		
NECK	NORMAL		
NECK LYMPHATICS / SALIVARY GLANDS	NOT ENLARGED OR TENDE	R	
THYROID GLAND	NOT ENLARGED		
TEMPERATURE	NORMAL		
PULSE	78/MIN		
RESPIRATORY RATE	NORMAL		
CARDIOVASCULAR SYSTEM			
ВР	122/76 MM HG (SITTING)		mm/Hg
PERICARDIUM	NORMAL		
APEX BEAT	NORMAL		
HEART SOUNDS	S1, S2 HEARD NORMALLY		
MURMURS	ABSENT		
RESPIRATORY SYSTEM			
SIZE AND SHAPE OF CHEST	NORMAL		
MOVEMENTS OF CHEST	SYMMETRICAL		
BREATH SOUNDS INTENSITY	NORMAL		
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)		
ADDED SOUNDS	ABSENT		
PER ABDOMEN			
APPEARANCE	NORMAL		
LIVER	NOT PALPABLE		
SPLEEN	NOT PALPABLE		
CENTRAL NERVOUS SYSTEM			
HIGHER FUNCTIONS	NORMAL		
CRANIAL NERVES	NORMAL		
CEREBELLAR FUNCTIONS	NORMAL		
SENSORY SYSTEM	NORMAL		











CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 9800465156 8800465156

SRL LTD
GRAND MALL, OPPOSITE SBI ZONAL OFFICE, SM ROAD, AMBAWADI,
AHMEDABAD, 380015
GUJRAT, INDIA
Tel: 079-48912999,079-48913999,079-48914999
Email : customercare.ahmedabad@srl.in

PATIENT ID : RIKKF070788321

PATIENT NAME : RIKKEN ETTE

ACCESSION NO :	0321VC002554	AGE: 33 Ye	ears SEX : F	emale		
DRAWN :		RECEIVED :	26-03-2022 1	1:46	REPORTED :	01-04-2022 13:44
REFERRING DOCT	OR: SELF				CLIENT	PATIENT ID :

REFERRING DOCTOR : SELF

Fest Report Status <u>Final</u>	Results	Biological Reference Interval Units
10TOR SYSTEM	NORMAL	
REFLEXES	NORMAL	
MUSCULOSKELETAL SYSTEM		
SPINE	NORMAL	
OINTS	NORMAL	
BASIC EYE EXAMINATION		
DISTANT VISION RIGHT EYE WITH GLASSES	WITH GLASSES NORM	1AL
DISTANT VISION LEFT EYE WITH GLASSES	WITH GLASSES NORM	1AL
NEAR VISION RIGHT EYE WITHOUT GLASSES	WITHIN NORMAL LIMI	IT
NEAR VISION LEFT EYE WITHOUT GLASSES	WITHIN NORMAL LIMI	IT
COLOUR VISION	NORMAL	
SUMMARY		
RELEVANT HISTORY	NOT SIGNIFICANT	
RELEVANT GP EXAMINATION FINDINGS	NOT SIGNIFICANT	
RELEVANT LAB INVESTIGATIONS	HEMOGLOBIN:- LOW,	MCV:- LOW, MCH:- LOW
	CBC:- PLATELET COUN	NT - HIGH
	ESR:- HIGH	
	HDL:- LOW	
	PPBS:- HIGH	
RELEVANT NON PATHOLOGY DIAGNOSTICS	HBA1C:- HIGH, MEAN NO ABNORMALITIES D	PLASMA GLUCOSE:- HIGH DETECTED
REMARKS / RECOMMENDATIONS	1) HEMOGLOBIN:- LO	W, MCV:- LOW, MCH:- LOW
	ADV:- TAKE MORE DIE	ETARY IRON
	2) ESR:- HIGH, CBC:-	PLATELET COUNT - HIGH
	ADV:- PHYSICIAN OPI	NION
	3) PPBS:- HIGH, HBA1	1C:- HIGH, MEAN PLASMA GLUCOSE:- HIGH
		E OF SWEET, SUGAR, STARCH IN DIET, REGULAF REPEAT FBS, PPBS AND HBA1C AND NION









CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
GRAND MALL, OPPOSITE SBI ZONAL OFFICE, SM ROAD, AMBAWADI,
AHMEDABAD, 380015
GUJRAT, INDIA
Tel : 079-48912999,079-48913999,079-48914999
Email : customercare.ahmedabad@srl.in

Test Report Status Final	Results	Biological Reference Interval Units
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
DRAWN :	RECEIVED : 26-03-2022 11:46	REPORTED : 01-04-2022 13:44
ACCESSION NO : 0321VC002554	AGE : 33 Years SEX : Female	
PATIENT NAME : RIKKEN ETTE		PATIENT ID : RIKKF070788321

Comments

OUR PANEL DOCTORS FOR NON-PATHOLOGY TESTS:-

GENERAL PHYSICIAN:- DR. PRIYANK KAPADIYA (M.B.B.S DNB MEDICINE)

RADIOLOGIST:- DR. KALPANA MODI (M.D.RADIOLOGY) // DR. SAHIL N SHAH (M.D.RADIOLOGY)

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology. RBC AND PLATELET INDICES-

Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait (<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT - NLR-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients ; A.-P. Yang, et al.; International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope. ERYTHRO SEDIMENTATION RATE, BLOOD-

Erythrocyte sedimentation rate (ESR) is a non - specific phenomena and is clinically useful in the diagnosis and monitoring of disorders associated with an increased production of acute phase reactants. The ESR is increased in pregnancy from about the 3rd month and returns to normal by the 4th week post partum. ESR is influenced by age, sex, menstrual cycle and drugs (eg. corticosteroids, contraceptives). It is especially low (0 -1mm) in polycythaemia, hypofibrinogenemia or congestive cardiac failure and when there are abnormalities of the red cells such as poikilocytosis, spherocytosis or sickle cells.

Reference :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACC Press, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th Edition"

GLUCOSE, FASTING, PLASMA-

ADA 2021 guidelines for adults, after 8 hrs fasting is as follows: Pre-diabetics: 100 - 125 mg/dL

Diabetic: > or = 126 mg/dL GLYCOSYLATED HEMOGLOBIN, EDTA WHOLE BLOOD-

Glycosylated hemoglobin (GHb) has been firmly established as an index of long-term blood glucose concentrations and as a measure of the risk for the development of complications in patients with diabetes mellitus. Formation of GHb is essentially irreversible, and the concentration in the blood depends on both the life span of the red blood cell (average 120 days) and the blood glucose concentration. Because the rate of formation of GHb is directly proportional to the concentration of glucose in the blood,

the GHb concentration represents the integrated values for glucose over the preceding 6-8 weeks. Any condition that alters the life span of the red blood cells has the potential to alter the GHb level. Samples from patients with hemolytic anemias will exhibit decreased glycated hemoglobin values due to the shortened life span of the red cells. This effect will depend upon the severity of the anemia. Samples from patients with polycythemia or post-splenectomy may exhibit increased glycated hemoglobin values due to a somewhat longer life span of the red cells. Glycosylated hemoglobins results from patients with HbSS, HbCC, and HbSC and HbD must be interpreted with caution, given the pathological processes, including anemia,

increased red cell turnover, transfusion requirements, that adversely impact HbA1c as a marker of long-term glycemic control. In these conditions, alternative forms of testing such as glycated serum protein (fructosamine) should be considered. "Targets should be individualized; More or less stringent glycemic goals may be appropriate for individual patients. Goals should be individualized based on duration of

diabetes, age/life expectancy, comorbid conditions, known CVD or advanced microvascular complications, hypoglycemia unawareness, and individual patient considerations.

References

Tietz Textbook of Clinical Chemistry and Molecular Diagnostics, edited by Carl A Burtis, Edward R.Ashwood, David E Bruns, 4th Edition, Elsevier publication, 2006, 879-884.

2. Forsham PH. Diabetes Mellitus: A rational plan for management. Postgrad Med 1982, 71,139-154.

3. Mayer TK, Freedman ZR: Protein glycosylation in Diabetes Mellitus: A review of laboratory measurements and their clinical utility. Clin Chim Acta 1983, 127, 147-184. GLUCOSE, POST-PRANDIAL, PLASMA-ADA Guidelines for 2hr post prandial glucose levels is only after ingestion of 75grams of glucose in 300 ml water, over a period of 5 minutes

CORONARY RISK PROFILE (LIPID PROFILE), SERUM .-





Scan to View Report





CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHEEL) F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156

SRL LTD
GRAND MALL, OPPOSITE SBI ZONAL OFFICE, SM ROAD, AMBAWADI,
AHMEDABAD, 380015
GUJRAT, INDIA
Tel: 079-48912999,079-48913999,079-48914999
Email : customercare.ahmedabad@srl.in

Test Report Status <u>Final</u>	Results	Biological Reference Interval Units
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
DRAWN :	RECEIVED : 26-03-2022 11:46	REPORTED : 01-04-2022 13:44
ACCESSION NO : 0321VC002554	AGE : 33 Years SEX : Female	
PATIENT NAME : RIKKEN ETTE		PATIENT ID : RIKKF070788321

Serum cholesterol is a blood test that can provide valuable information for the risk of coronary artery disease This test can help determine your risk of the build up of plaques in your arteries that can lead to narrowed or blocked arteries throughout your body (atherosclerosis). High cholesterol levels usually don't cause any signs or symptoms, so a cholesterol test is an important tool. High cholesterol levels often are a significant risk factor for heart disease and important for diagnosis of hyperlipoproteinemia, atherosclerosis, hepatic and thyroid diseases.

Serum Triglyceride are a type of fat in the blood. When you eat, your body converts any calories it doesn't need into triglycerides, which are stored in fat cells. High triglyceride levels are associated with several factors, including being overweight, eating too many sweets or drinking too much alcohol, smoking, being sedentary, or having diabetes with elevated blood sugar levels. Analysis has proven useful in the diagnosis and treatment of patients with diabetes mellitus, nephrosis, liver obstruction, other diseases involving lipid metabolism, and various endocrine disorders. In conjunction with high density lipoprotein and total serum cholesterol, a triglyceride determination provides valuable information for the assessment of coronary heart disease risk. It is done in fasting state.

High-density lipoprotein (HDL) cholesterol. This is sometimes called the ""good"" cholesterol because it helps carry away LDL cholesterol, thus keeping arteries open and blood flowing more freely.HDL cholesterol is inversely related to the risk for cardiovascular disease. It increases following regular exercise, moderate alcohol consumption and with oral estrogen therapy. Decreased levels are associated with obesity, stress, cigarette smoking and diabetes mellitus.

SERUM LDL The small dense LDL test can be used to determine cardiovascular risk in individuals with metabolic syndrome or established/progressing coronary artery disease, individuals with triglyceride levels between 70 and 140 mg/dL, as well as individuals with a diet high in trans-fat or carbohydrates. Elevated sdLDL levels are associated with metabolic syndrome and an 'atherogenic lipoprotein profile', and are a strong, independent predictor of cardiovascular disease. Elevated levels of LDL arise from multiple sources. A major factor is sedentary lifestyle with a diet high in saturated fat. Insulin-resistance and pre-diabetes have also been implicated, as has genetic predisposition. Measurement of sdLDL allows the clinician to get a more comprehensive picture of lipid risk factors and tailor treatment accordingly. Reducing LDL levels will reduce the risk of CVD and MI.

Non HDL Cholesterol - Adult treatment panel ATP III suggested the addition of Non-HDL Cholesterol as an indicator of all atherogenic lipoproteins (mainly LDL and VLDL). NICE guidelines recommend Non-HDL Cholesterol measurement before initiating lipid lowering therapy. It has also been shown to be a better marker of risk in both primary and secondary prevention studies.

Recommendations:

Results of Lipids should always be interpreted in conjunction with the patient's medical history, clinical presentation and other findings.

NON FASTING LIPID PROFILE includes Total Cholesterol, HDL Cholesterol and calculated non-HDL Cholesterol. It does not include triglycerides and may be best used in patients for whom fasting is difficult. LIVER FUNCTION PROFILE, SERUM-

LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels results from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, (indirect) bilirubin in Viral hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors &Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver,liver cancer,kidney failure,hemolytic anemia,pancreatitis,hemochromatosis. AST levels may also increase after a heart attack or strenuous activity.ALT test measures the amount of this enzyme in the blood.ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic

hepatitis, obstruction of bile ducts, cirrhosis. ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease,Rickets,Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia,Malnutrition,Protein deficiency,Wilson's disease.GGT is an enzyme found in cell membranes of many tissues mainly in the liver,kidney and pancreas.It is also found in other tissues including intestine,spleen,heart, brain and seminal vesicles.The highest concentration is in the kidney,but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also and pancreas.Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc.Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum.Protein in the plasma is made up of albumin and globulin.Higher-than-normal levels may be due to:Chronic inflammation or infection, including HIV and hepatitis B or C,Multiple myeloma,Waldenstrom's disease.Lower-than-normal levels may be due to:Chronic inflammation or infection, including HIV and hepatitis B or C,Multiple myeloma,Waldenstrom's disease.Lower-than-normal levels may be due to: Agammaglobulinemia,Bleeding (hemorrhage),Burns,Glomerulonephritis,Liver disease, Malabsorption,Malnutrition,Nephrotic syndrome,Protein-losing enteropathy etc.Human serum albumin is the most abundant protein in human blood plasma.It is produced in the liver.Albumin constitutes about half of the blood serum protein.Low blood albumin levels (hypoalbuminemia) can be caused by:Liver disease like cirrhosis of the liver, nephrotic syndrome,protein-losing enteropathy,Burns,hemodilution,increased vascular permeability or decreased lymphatic clearance,malnutrition and wasting etc SERUM BLOOD UREA NITROGEN-Councer disease like cirrhosis of the Councer disease and the disease disease.

Causes of Increased levels

Pre renal

High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal
 Renal Failure

Post Renal

• Malignancy, Nephrolithiasis, Prostatism



Scan to View Report

Page 8 Of 11

DIAGNOSTIC REPORT			≧SRL
P	atient Ref. No. 77700000189080	5	
CLIENT CODE: C000138364			Diagnostics
CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHE F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156	EL)	SRL LTD GRAND MALL, OPPOSITE SBI ZONAL AHMEDABAD, 380015 GUJRAT, INDIA Tel : 079-48912999,079-48913999,0 Email : customercare.ahmedabad@sr	79-48914999
PATIENT NAME : RIKKEN ETTE		PATIENT ID	RIKKF070788321
ACCESSION NO : 0321VC002554	AGE: 33 Years SEX: Fen	nale	
DRAWN :	RECEIVED : 26-03-2022 11:4	46 REPORTED : 01-04-	-2022 13:44
REFERRING DOCTOR : SELF		CLIENT PATIEN	T ID :
Test Report Status <u>Final</u>	Results	Biological Referen	ce Interval Units
Causes of decreased levels			

 Liver disease SIADH.

ſ ſ 8 I

1

CREATININE, SERUM-

Higher than normal level may be due to:

Blockage in the urinary tract

Kidney problems, such as kidney damage or failure, infection, or reduced blood flow

Loss of body fluid (dehydration)

Muscle problems, such as breakdown of muscle fibers

• Problems during pregnancy, such as seizures (eclampsia)), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

• Myasthenia Gravis • Muscular dystrophy URIC ACID, SERUM-Causes of Increased levels Dietary High Protein Intake. Prolonged Fasting, Rapid weight loss. Gout Lesch nyhan syndrome. Type 2 DM. Metabolic syndrome

Causes of decreased levels

Low Zinc Intake

- OCP's
- Multiple Sclerosis

Nutritional tips to manage increased Uric acid levels • Drink plenty of fluids • Limit animal proteins

· High Fibre foods

• Vit C Intake

Antioxidant rich foods
 ELECTROLYTES (NA/K/CL), SERUM-

Sodium levels are Increased in dehydration, cushing's syndrome, aldosteronism & decreased in Addison's disease, hypopituitarism, liver disease. Hypokalemia (low K) is common in vomiting, diarrhea, alcoholism, folic acid deficiency and primary aldosteronism. Hyperkalemia may be seen in end-stage renal failure, hemolysis, trauma, Addison's disease, metabolic acidosis, acute starvation, dehydration, and with rapid K infusion.Chloride is increased in dehydration, renal tubular acidosis (hyperchloremia metabolic acidosis), acute renal failure, metabolic acidosis associated with prolonged diarrhea and loss of sodium bicarbonate, diabetes insipidus, adrenocortical hyperfuction, salicylate intoxication and with excessive infusion of isotonic saline or extremely high dietary intake of salt. Chloride is decreased in overhydration, chronic respiratory acidosis, salt-losing nephritis, metabolic alkalosis, congestive heart failure, Addisonian crisis, certain types of metabolic acidosis, persistent gastric secretion and prolonged vomiting,

URINALYSIS-Routine urine analysis assists in screening and diagnosis of various metabolic, urological, kidney and liver disorders

Protein: Elevated proteins can be an early sign of kidney disease. Urinary protein excretion can also be temporarily elevated by strenuous exercise, orthostatic proteinuria, dehydration, urinary tract infections and acute illness with fever

Glucose: Uncontrolled diabetes mellitus can lead to presence of glucose in urine. Other causes include pregnancy, hormonal disturbances, liver disease and certain medications.

Ketones: Uncontrolled diabetes mellitus can lead to presence of ketones in urine. Ketones can also be seen in starvation, frequent vomiting, pregnancy and strenuous exercise

Blood: Occult blood can occur in urine as intact erythrocytes or haemoglobin, which can occur in various urological, nephrological and bleeding disorders.

Leukocytes: An increase in leukocytes is an indication of inflammation in urinary tract or kidneys. Most common cause is bacterial urinary tract infection.

Nitrite: Many bacteria give positive results when their number is high. Nitrite concentration during infection increases with length of time the urine specimen is retained in bladder prior to collection.

pH: The kidneys play an important role in maintaining acid base balance of the body. Conditions of the body producing acidosis/ alkalosis or ingestion of certain type of food can affect the pH of urine.

Specific gravity: Specific gravity gives an indication of how concentrated the urine is. Increased specific gravity is seen in conditions like dehydration, glycosuria and proteinuria while decreased specific gravity is seen in excessive fluid intake, renal failure and diabetes insipidus. Bilirubin: In certain liver diseases such as biliary obstruction or hepatitis, bilirubin gets excreted in urine.

Urobilinogen: Positive results are seen in liver diseases like hepatitis and cirrhosis and in cases of hemolytic anemia THYROID PANEL, SERUM-

Triiodothyronine T3, is a thyroid hormone. It affects almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate. Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Thyroxine T4, Thyroxine's principal function is to stimulate the metabolism of all cells and tissues in the body. Excessive secretion of thyroxine in the body is





Test Report Status Final	Results	Biological Reference Interval Units
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
DRAWN :	RECEIVED : 26-03-2022 11:46	REPORTED : 01-04-2022 13:44
PATIENT NAME : RIKKEN ETTE ACCESSION NO : 0321VC002554	AGE : 33 Years SEX : Femal	PATIENT ID : RIKKF070788321 e
CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWHE F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156	G Al G Te	RL LTD RAND MALL, OPPOSITE SBI ZONAL OFFICE,SM ROAD, AMBAWADI, HMEDABAD, 380015 UJRAT, INDIA el : 079-48912999,079-48913999,079-48914999 mail : customercare.ahmedabad@srl.in
	iii <u>2.5 - Frank B. 5</u> - Frank B. 5 - Frank B. 111 atient Ref. No. 777000001890805	Diagnostics
	CSKL	

hyperthyroidism, and deficient secretion is called hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hypothyroidism, TSH levels are low.

TSH & Total T3

Below mentioned are	the guidelines for Pre	egnancy related re	eference ranges for Tota	I T4, ⁻
Levels in	TOTAL T4	TSH3G	TOTAL T3	
Pregnancy	(µg/dL)	(µIU/mL)	(ng/dL)	
First Trimester	6.6 - 12.4	0.1 - 2.5	81 - 190	
2nd Trimester	6.6 - 15.5	0.2 - 3.0	100 - 260	
3rd Trimester	6.6 - 15.5	0.3 - 3.0	100 - 260	
Below mentioned are	the guidelines for ag	e related reference	e ranges for T3 and T4.	
Т3	T	4		
(ng/dL)	(µg/0	dL)		
New Born: 75 - 260	1-3 day: 8	.2 - 19.9		
	1 Week: 6.0	- 15.9		

Final

NOTE: TSH concentrations in apparently normal euthyroid subjects are known to be highly skewed, with a strong tailed distribution towards higher TSH values. This is well documented in the pediatric population including the infant age group. Kindly note: Method specific reference ranges are appearing on the report under biological reference range.

Reference:

1. Burtis C.A., Ashwood E. R. Bruns D.E. Teitz textbook of Clinical Chemistry and Molecular Diagnostics, 4th Edition.

Gowenlock A.H. Varley's Practical Clinical Biochemistry, 6th Edition.
 Behrman R.E. Kilegman R.M., Jenson H. B. Nelson Text Book of Pediatrics, 17th Edition

DIAGNOSTIC REPORT

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A,B,O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same.'

The test is performed by both forward as well as reverse grouping methods. MEDICAL

THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOLABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.





DIAGNOSTIC REPORT		SRL
	Patient Ref. No. 777000001890805	
CLIENT CODE: C000138364		Diagnostics
CLIENT'S NAME AND ADDRESS : ACROFEMI HEALTHCARE LTD (MEDIWH F-703, LADO SARAI, MEHRAULI SOUTH WEST DELHI NEW DELHI 110030 DELHI INDIA 8800465156	IEEL)	SRL LTD GRAND MALL, OPPOSITE SBI ZONAL OFFICE,SM ROAD, AMBAWADI, AHMEDABAD, 380015 GUJRAT, INDIA Tel : 079-48912999,079-48913999,079-48914999 Email : customercare.ahmedabad@srl.in
PATIENT NAME : RIKKEN ETTE		PATIENT ID : RIKKF070788321
ACCESSION NO : 0321VC002554	AGE: 33 Years SEX: Fem	ale
DRAWN :	RECEIVED : 26-03-2022 11:4	6 REPORTED : 01-04-2022 13:44
REFERRING DOCTOR : SELF		CLIENT PATIENT ID :
Test Report Status <u>Final</u>	Results	Biological Reference Interval Units

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE

ULTRASOUND ABDOMEN

ULTRASOUND ABDOMEN

NO ABNORMALITIES DETECTED IN PRESENT EXAMINATION

End Of Report Please visit www.srlworld.com for related Test Information for this accession



Dr.Priyank Kapadia Physician



Dr Kalpana Modi Radiologist

Dr.Sahil .N.Shah **Consultant Radiologist**

Grient

Dr.Swati M.Shah **Consultant Pathologist**

CONDITIONS OF LABORATORY TESTING & REPORTING			
 It is presumed that the test sample belongs to the patient named or identified in the test requisition form. All Tests are performed and reported as per the turnaround time stated in the SRL Directory of services (DOS). SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity. A requested test might not be performed if: a. Specimen received is insufficient or inappropriate specimen quality is unsatisfactory b. Incorrect specimen type c. Request for testing is withdrawn by the ordering doctor or patient d. There is a discrepancy between the label on the specimen container and the name on the test requisition form 	 The results of a laboratory test are dependent on the quality of the sample as well as the assay technology. Result delays could be because of uncontrolled circumstances. e.g. assay run failure. Tests parameters marked by asterisks are excluded from the "scope" of NABL accredited tests. (If laboratory is accredited). Laboratory results should be correlated with clinical information to determine Final diagnosis. Test results are not valid for Medico- legal purposes. In case of queries or unexpected test results please call at SRL customer care (Toll free: 1800-222-000). Post proper investigation repeat analysis may be carried out. 		
	SRL Limited		
	Fortis Hospital, Sector 62, Phase VIII,		

Mohali 160062





Scan to View Report