



Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:28
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 2024-09-22 09:52:17
UHID/MR NO	: ALDP.0000149782	Received	: 2024-09-22 09:52:17
Visit ID	: ALDP0228152425	Reported	: 23/Sep/2024 12:10:52
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CARDIOLOGY-ECG MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

ECG / EKG

Sinus, Regular	
74	/mt
74	/mt
Normal	
Normal	
Normal Normal Normal	
Normal	
Normal	
Normal	
	74 74 Normal Normal Normal Normal Normal Normal

9. 1 – v <u>FINAL IMPRESSION</u>

ECG Within Normal Limits: Sinus Rhythm, Sinus Arrhythmia Seen. Please correlate clinically.











Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:27
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 09:45:56
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 11:15:29
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 13:40:26
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing) , Blood				
Blood Group	0			erythrocyte Magnetized Technology / Tube Agglutina
Rh (Anti-D)	POSITIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY / TUBE AGGLUTINA
Complete Blood Count (CBC), Whole Blood				
Haemoglobin	13.60	g/dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRIC METHOD (CYANIDE-FREE REAGENT)
TLC (WBC)	4,600.00	/Cu mm	4000-10000	IMPEDANCE METHOD
DLC				
Polymorphs (Neutrophils)	65.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	29.00	%	20-40	FLOW CYTOMETRY
Monocytes	4.00	%	2-10	FLOW CYTOMETRY
Eosinophils	2.00	%	1-6	FLOW CYTOMETRY
Basophils ESR	0.00	%	< 1-2	FLOW CYTOMETRY
Observed	6.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5	



80-91 Yr 15.8







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:27
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 09:45:56
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 11:15:29
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 13:40:26
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF HAEMATOLOGY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95	
			if anaemic)	
Corrected	-	Mm for 1st hr.	< 20	
PCV (HCT)	42.00	%	40-54	
Platelet count				
Platelet Count	1.85	LACS/cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.10	fL	9-17	ELECTRONIC IMPEDANCE
P-LCR (Platelet Large Cell Ratio)	-	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.25	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	13.60	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBC Count				
RBC Count	4.51	Mill./cu mm	3.7-5.0	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	94.60	fl	80-100	CALCULATED PARAMETER
MCH	30.30	pg	27-32	CALCULATED PARAMETER
MCHC	32.00	%	30-38	CALCULATED PARAMETER
RDW-CV	13.50	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	47.70	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	2,990.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	92.00	/cu mm	40-440	

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:28
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 09:45:56
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 11:15:29
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 13:50:24
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Uni	t Bio. Ref. Interv	al Method
GLUCOSE FASTING, Plasma Glucose Fasting	82.30	mg/dl	< 100 Normal 100-125 Pre-diabetes ≥ 126 Diabetes	GOD POD

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Glucose PP	107.90	mg/dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

GLYCOSYLATED HAEMOGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	4.60	% NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	27.10	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	86	mg/dl	

Interpretation:

<u>NOTE</u>:-

• eAG is directly related to A1c.



View Reports on

Chandan 24x7 App





T



CHANDAN DIAGNOSTIC CENTRE

Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:28
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 09:45:56
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 11:15:29
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 13:50:24
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	Degree of Glucose Control Unit
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. **Some danger of hypoglycemic reaction in Type 1 diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

<u>Clinical Implications:</u>

*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

*With optimal control, the HbA 1c moves toward normal levels.

*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated *Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

*Pregnancy d. chronic renal failure. Interfering Factors:

*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood Urea Nitrogen)	9.10	mg/dL	7.0-23.0	CALCULATED
Sample:Serum				



Page 5 of 12

View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-970	073	Registered	d On	: 22/Sep/2024 09	
Age/Gender	: 29 Y 11 M 0 D /F		Collected		: 22/Sep/2024 09	
UHID/MR NO Visit ID	: ALDP.0000149782 : ALDP0228152425		Received Reported		: 22/Sep/2024 11 : 22/Sep/2024 13	
	: Dr. MEDIWHEEL-ARCO	ΕΕΜΙ ΗΕΔΙ ΤΗ				:50:24
Ref Doctor	CARE LTD -		Status		: Final Report	
		DEPARTMEN				
	MEDIWHE	EEL BANK OF B				
Test Name		Result	ι	Jnit	Bio. Ref. Interval	Method
Interpretation:	UN levels oon he seen in th	o following.				
Note: Elevated Bu	UN levels can be seen in th	ie tonowing:				
High-protein diet, D	Pehydration, Aging, Certain m	edications, Burns	, Gastrointesti	mal (GI) b	leeding.	
LOW BUN levels c	an be seen in the following	; :				
Low-protein diet, ov	verhydration, Liver disease.					
reatinine		0.71	mg/dl	0.5-1.2	.0 M	ODIFIED JAFFES
ample:Serum		0.71	mg/dl	0.5-1.2	0 M	ODIFIED JAFFES
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine c	single creatinine value must b gher creatinine concentration. concentration. Serum creatini ildly and may result in anoma	e interpreted in lig The trend of serur ne concentrations lous values if seru	ht of the patien n creatinine co may increase m samples ha	nts muscle oncentratio when an A ve heterop	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo	a greater muscle important than is taken. The assay
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi lipemic.	ther creatinine concentration.	e interpreted in lig The trend of serur ne concentrations	ht of the patien n creatinine co may increase	nts muscle oncentration when an A	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine c could be affected mi lipemic. Interpretation: Note:-	ther creatinine concentration.	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57	ht of the patien n creatinine co may increase m samples ha	nts muscle oncentratio when an A ve heterop	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine c could be affected mi lipemic. Interpretation: Note:- Elevated uric acid	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 3.57	ht of the patien n creatinine co may increase m samples ha mg/dl	nts muscle oncentratic when an A ve heterop 2.5-6.0	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine c could be affected mi lipemic. Interpretation: Note:- Elevated uric acid	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma l levels can be seen in the f rotein diet, alcohol), Chronic	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 3.57	ht of the patien n creatinine co may increase m samples ha mg/dl	nts muscle oncentratic when an A ve heterop 2.5-6.0	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo	a greater muscle important than is taken. The assay olyzed, icteric or
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pu- FT (WITH GAMM	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma l levels can be seen in the f rotein diet, alcohol), Chronic	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 3.57	ht of the patien n creatinine co may increase m samples ha mg/dl	nts muscle oncentratic when an A ve heterop 2.5-6.0	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo) UF	a greater muscle important than is taken. The assay olyzed, icteric or
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pu- FT (WITH GAMM SGOT / Aspartate A	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma I levels can be seen in the f rotein diet, alcohol), Chronic A GT) , <i>Serum</i>	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 Ollowing: kidney disease, H	ht of the patien n creatinine co may increase m samples har mg/dl	nts muscle oncentratio when an A ve heterop 2.5-6.0 Obesity.	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo) UF	a greater muscle important than is taken. The assay olyzed, icteric or RICASE
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pu- FT (WITH GAMIM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT)	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma I levels can be seen in the f rotein diet, alcohol), Chronic A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 following: kidney disease, H 18.00 10.20 13.30	ht of the patie n creatinine co may increase m samples ha mg/dl fypertension, C U/L U/L IU/L	nts muscle oncentratio when an A ve heterop 2.5-6.0 Obesity. < 35 < 40 11-50	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hema) UF	a greater muscle important than is taken. The assay olyzed, icteric or RICASE
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pu- FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma I levels can be seen in the f rotein diet, alcohol), Chronic A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 following: kidney disease, H 18.00 10.20 13.30 7.27	ht of the patien n creatinine co may increase m samples har mg/dl Typertension, C U/L U/L U/L IU/L gm/dl	nts muscle oncentratio when an A ve heterop 2.5-6.0 Obesity. Obesity. < 35 < 40 11-50 6.2-8.0	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo) UF 	a greater muscle important than is taken. The assay olyzed, icteric or RICASE
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pu- FT (WITH GAIMIM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein Albumin	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma I levels can be seen in the f rotein diet, alcohol), Chronic A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 following: kidney disease, H 18.00 10.20 13.30 7.27 4.32	ht of the patie n creatinine co may increase m samples har mg/dl fypertension, C U/L U/L U/L IU/L gm/dl gm/dl	nts muscle oncentratio when an <i>A</i> ve heterop 2.5-6.0 Obesity. Cobesity. < 35 < 40 11-50 6.2-8.0 3.4-5.4	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo) UF) UF F(F(OF B B.	a greater muscle important than is taken. The assay olyzed, icteric or RICASE
ample:Serum Interpretation: The significance of s mass will have a hig absolute creatinine of could be affected mi- lipemic. Iric Acid ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-pu- FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	ther creatinine concentration. concentration. Serum creatini ildly and may result in anoma I levels can be seen in the f rotein diet, alcohol), Chronic A GT) , <i>Serum</i> Aminotransferase (AST)	e interpreted in lig The trend of serur ne concentrations lous values if seru 3.57 following: kidney disease, H 18.00 10.20 13.30 7.27	ht of the patien n creatinine co may increase m samples har mg/dl fypertension, C U/L U/L IU/L gm/dl	nts muscle oncentratio when an A ve heterop 2.5-6.0 Obesity. Obesity. < 35 < 40 11-50 6.2-8.0	e mass. A patient with ons over time is more ACE inhibitor (ACE) hilic antibodies, hemo) UF) UF 	a greater muscle important than is taken. The assay olyzed, icteric or RICASE









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:28
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 09:45:56
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 11:15:29
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 13:50:24
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Ur	nit Bio. Ref. Inte	rval Method
Alkaline Phosphatase (Total)	61.00	U/L	42.0-165.0	PNP/AMP KINETIC
Bilirubin (Total)	0.54	mg/dl	0.3-1.2	JENDRASSIK & GROF
Bilirubin (Direct)	0.15	mg/dl	< 0.30	JENDRASSIK & GROF
Bilirubin (Indirect)	0.39	mg/dl	< 0.8	JENDRASSIK & GROF
LIPID PROFILE (MINI), Serum				
Cholesterol (Total)	176.00	mg/dl	<200 Desirable 200-239 Borderline H > 240 High	CHOD-PAP igh
HDL Cholesterol (Good Cholesterol)	59.20	mg/dl	30-70	DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	96	mg/dl	< 100 Optimal 100-129 Nr. Optimal/Above Optir 130-159 Borderline H 160-189 High > 190 Very High	
VLDL	20.94	mg/dl	10-33	CALCULATED
Triglycerides	104.70	mg/dl	< 150 Normal 150-199 Borderline H 200-499 High >500 Very High	GPO-PAP igh

AS

Dr.Akanksha Singh (MD Pathology)











Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:27
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 14:01:28
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 14:12:40
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 16:39:44
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE, Urine				
Color	LIGHT YELLOW			
Specific Gravity	1.005			
Reaction PH	Acidic (6.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	< 10 Absent	DIPSTICK
			10-40 (+)	
			40-200 (++) 200-500 (+++)	
			> 500 (+++)	
Sugar	ABSENT	gms%	< 0.5 (+)	DIPSTICK
	7.002.111	9111070	0.5-1.0 (++)	
			1-2 (+++)	
			>2 (++++)	
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial cells	1-3/h.p.f			MICROSCOPIC
	·			EXAMINATION
Pus cells	1-3/h.p.f			
RBCs	ABSENT			MICROSCOPIC
				EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC
				EXAMINATION
Others	ABSENT			

Urine Microscopy is done on centrifuged urine sediment.









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:27
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 14:01:28
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 14:12:40
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 16:39:44
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

	MEDIWHEEL BAINS OF BARODA FEMALE ADOVE 40 TRS					
Test Name	Result	Unit	Bio. Ref. Interval	Method		
SUGAR, FASTING STAGE, Urine						
Sugar, Fasting stage	ABSENT	gms%				
Interpretation:						
(+) < 0.5						
(++) 0.5-1.0						
(+++) 1-2						

(++++) > 2

AS

View Reports on

Chandan 24x7 App

Dr.Akanksha Singh (MD Pathology)









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:27
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 22/Sep/2024 09:45:56
UHID/MR NO	: ALDP.0000149782	Received	: 22/Sep/2024 11:15:29
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 14:10:54
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

Test Name	Result	Unit E	io. Ref. Interval	Method
THYROID PROFILE - TOTAL , Serum				
T3, Total (tri-iodothyronine)	137.00	5	4.61–201.7	CLIA
T4, Total (Thyroxine)	5.51	5	.2-12.6	CLIA
TSH (Thyroid Stimulating Hormone)	31.100	μIU/mL 0	.27 - 5.5	CLIA
Interpretation:		0.3-4.5 μIU/mL 0.5-4.6 μIU/mL 0.8-5.2 μIU/mL 0.5-8.9 μIU/mL 0.7-27 μIU/mL 2.3-13.2 μIU/mL 0.7-64 μIU/mL 1-39 μIU/mL 1.7-9.1 μIU/mL	Second Trim Third Trimes Adults Premature Cord Blood Child(21 wk L Child	nester ster 55-87 Years 28-36 Week > 37Week

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

3) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

4) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

8) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

Dr.Akanksha Singh (MD Pathology)

Page 10 of 12









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:28
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 2024-09-22 10:06:06
UHID/MR NO	: ALDP.0000149782	Received	: 2024-09-22 10:06:06
Visit ID	: ALDP0228152425	Reported	: 23/Sep/2024 11:20:50
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF X-RAY

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

X-RAY DIGITAL CHEST PA

<u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Both lung field did not reveal any significant lesion.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Soft tissue shadow appears normal.
- Bony cage is normal.

Please correlare clinically.



Dr. Aishwarya Neha (MD Radiodiagnosis



View Reports on Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mrs.NISHU SINGH-97073	Registered On	: 22/Sep/2024 09:20:28
Age/Gender	: 29 Y 11 M 0 D /F	Collected	: 2024-09-22 13:58:04
UHID/MR NO	: ALDP.0000149782	Received	: 2024-09-22 13:58:04
Visit ID	: ALDP0228152425	Reported	: 22/Sep/2024 13:59:28
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA FEMALE ABOVE 40 YRS

ULTRASOUND WHOLE ABDOMEN (UPPER & LOWER)

LIVER: - Normal in size (13.1 cm), shape and echogenicity. No focal lesion is seen. No intra hepatic biliary radicle dilation is seen.

GALL BLADDER :- Well distended. Normal wall thickness is seen. No evidence of calculus/focal mass lesion/pericholecystic fluid is seen.

CBD :- Normal in calibre at porta.

PORTAL VEIN: - Normal in calibre and colour uptake at porta.

PANCREAS: - Head is visualised, normal in size & echopattern. No evidence of ductal dilatation or calcification is seen. Rest of the pancreas is obscured by bowel gases.

SPLEEN: - Normal in size (9.8 cm), shape and echogenicity. No evidence of mass lesion is seen.

RIGHT KIDNEY: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

LEFT KIDNEY: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

URINARY BLADDER :- Is adequately distended. No evidence of wall thickening/calculus is seen.

UTERUS :- Is normal in size (6.4 x 2.6 cm). No focal myometrial lesion is seen. Endometrium is normal in thickness.


OVARIES :- Bilateral ovaries are normal in size, shape and echogenicity. Right ovary - 24 x 12 mm, Left ovary - 20 x 10 mm.

ADNEXA :- No obvious adnexal pathology is seen.

HIGH RESOLUTION :- No evidence of bowel loop dilatation or abnormal wall thickening is seen. No significant retroperitoneal lymphadenopathy is seen. No free fluid is seen in the abdomen/pelvis.

IMPRESSION : No significant abnormality seen.

Please correlate clinically.



08069366666

Chandan 24x7 App