



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE**BLOOD COUNTS, EDTA WHOLE BLOOD**

HEMOGLOBIN (HB)	13.0	12.0 - 15.0	g/dL
METHOD : SLS- HEMOGLOBIN DETECTION METHOD			
RED BLOOD CELL (RBC) COUNT	4.59	3.8 - 4.8	mil/ μ L
METHOD : HYDRODYNAMIC FOCUSING BY DC DETECTION			
WHITE BLOOD CELL (WBC) COUNT	5.85	4.0 - 10.0	thou/ μ L
METHOD : FLUORESCENCE FLOW CYTOMETRY			
PLATELET COUNT	231	150 - 410	thou/ μ L
METHOD : HYDRODYNAMIC FOCUSING BY DC DETECTION			

RBC AND PLATELET INDICES

HEMATOCRIT (PCV)	40.4	36.0 - 46.0	%
METHOD : CUMULATIVE PULSE HEIGHT DETECTION METHOD			
MEAN CORPUSCULAR VOLUME (MCV)	88.0	83.0 - 101.0	fL
METHOD : CALCULATED FROM RBC & HCT			
MEAN CORPUSCULAR HEMOGLOBIN (MCH)	28.3	27.0 - 32.0	pg
METHOD : CALCULATED FROM THE RBC & HGB			
MEAN CORPUSCULAR HEMOGLOBIN CONCENTRATION (MCHC)	32.2	31.5 - 34.5	g/dL
METHOD : CALCULATED FROM THE HGB & HCT			
RED CELL DISTRIBUTION WIDTH (RDW)	13.6	11.6 - 14.0	%
METHOD : CALCULATED FROM RBC SIZE DISTRIBUTION CURVE			
MENTZER INDEX	19.2		
MEAN PLATELET VOLUME (MPV)	11.6	High 6.8 - 10.9	fL
METHOD : CALCULATED FROM PLATELET COUNT & PLATELET HEMATOCRIT			

WBC DIFFERENTIAL COUNT

NEUTROPHILS	63	40 - 80	%
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING			
LYMPHOCYTES	27	20 - 40	%
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING			
MONOCYTES	6	2 - 10	%
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING			
EOSINOPHILS	4	1 - 6	%
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING			
BASOPHILS	0	0 - 1	%
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING			
ABSOLUTE NEUTROPHIL COUNT	3.69	2.0 - 7.0	thou/ μ L



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE LYMPHOCYTE COUNT		1.60	1.0 - 3.0	thou/ μ L
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE MONOCYTE COUNT		0.34	0.2 - 1.0	thou/ μ L
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE EOSINOPHIL COUNT		0.23	0.02 - 0.50	thou/ μ L
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING				
ABSOLUTE BASOPHIL COUNT		0.00	Low 0.02 - 0.10	thou/ μ L
METHOD : FLOW CYTOMETRY WITH LIGHT SCATTERING				
NEUTROPHIL LYMPHOCYTE RATIO (NLR)		2.3		
MORPHOLOGY				
RBC		NORMOCYTIC NORMOCHROMIC		
WBC		NORMAL MORPHOLOGY		
METHOD : MICROSCOPIC EXAMINATION				
PLATELETS		ADEQUATE		
ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD				
E.S.R		22	High < 20	mm at 1 hr
GLUCOSE FASTING, FLUORIDE PLASMA				
FBS (FASTING BLOOD SUGAR)		91	Normal 75 - 99 Pre-diabetics: 100 - 125 Diabetic: > or = 126	mg/dL
METHOD : ENZYMATIC REFERENCE METHOD WITH HEXOKINASE				
GLYCOSYLATED HEMOGLOBIN (HBA1C), EDTA WHOLE BLOOD				
HBA1C		5.4	Non-diabetic Adult < 5.7 Pre-diabetes 5.7 - 6.4 Diabetes diagnosis: > or = 6.5 Therapeutic goals: < 7.0 Action suggested : > 8.0 (ADA Guideline 2021)	%
METHOD : HPLC				
ESTIMATED AVERAGE GLUCOSE (EAG)		108.3	< 116.0	mg/dL
METHOD : CALCULATED PARAMETER				
GLUCOSE, POST-PRANDIAL, PLASMA				
PPBS (POST PRANDIAL BLOOD SUGAR)		83	70 - 139	mg/dL
METHOD : ENZYMATIC REFERENCE METHOD WITH HEXOKINASE				
LIPID PROFILE, SERUM				



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002381402

CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN : RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
CHOLESTEROL, TOTAL		140	Desirable cholesterol level < 200 Borderline high cholesterol 200 - 239 High cholesterol > / = 240	mg/dL
METHOD : ENZYMATIC COLORIMETRIC ASSAY				
TRIGLYCERIDES		47	Normal: < 150 Borderline high: 150 - 199 High: 200 - 499 Very High: >/= 500	mg/dL
METHOD : ENZYMATIC COLORIMETRIC ASSAY				
HDL CHOLESTEROL		66	High Low HDL Cholesterol <40 High HDL Cholesterol >/= 60	mg/dL
METHOD : ENZYMATIC, COLORIMETRIC				
CHOLESTEROL LDL		65	Adult levels: Optimal < 100 Near optimal/above optimal: 100-129 Borderline high : 130-159 High : 160-189 Very high : = 190	mg/dL
METHOD : ENZYMATIC COLORIMETRIC ASSAY				
NON HDL CHOLESTEROL		74	Desirable : < 130 Above Desirable : 130 -159 Borderline High : 160 - 189 High : 190 - 219 Very high : > / = 220	mg/dL
VERY LOW DENSITY LIPOPROTEIN CHOL/HDL RATIO		9.4 2.1	< OR = 30.0 Low Low Risk : 3.3 - 4.4 Average Risk : 4.5 - 7.0 Moderate Risk : 7.1 - 11.0 High Risk : > 11.0	mg/dL
LDL/HDL RATIO		1.0	0.5 - 3.0 Desirable/Low Risk 3.1 - 6.0 Borderline/Moderate Risk >6.0 High Risk	



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002381402

CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
S.K. Tower, Hari Niwas, LBS Marg
THANE, 400602
MAHARASHTRA, INDIA
Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 AGE : 36 Years SEX : Female ABHA NO :

DRAWN : RECEIVED : 18/02/2023 09:04 REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

- Cholesterol levels help assess the patient risk status and to follow the progress of patient under treatment to lower serum cholesterol concentrations.
- Serum Triglyceride (TG) are a type of fat and a major source of energy for the body. Both quantity and composition of the diet impact on plasma triglyceride concentrations. Elevations in TG levels are the result of overproduction and impaired clearance. High TG are associated with increased risk for CAD (Coronary artery disease) in patients with other risk factors, such as low HDL-C, some patient groups with elevated apolipoprotein B concentrations, and patients with forms of LDL that may be particularly atherogenic.
- HDL-C plays a crucial role in the initial step of reverse cholesterol transport, this considered to be the primary atheroprotective function of HDL
- LDL -C plays a key role in causing and influencing the progression of atherosclerosis and, in particular, coronary sclerosis. The majority of cholesterol stored in atherosclerotic plaques originates from LDL, thus LDL-C value is the most powerful clinical predictor.
- Non HDL cholesterol: Non-HDL-C measures the cholesterol content of all atherogenic lipoproteins, including LDL hence it is a better marker of risk in both primary and secondary prevention studies. Non-HDL-C also covers, to some extent, the excess ASCVD risk imparted by the sdLDL, which is significantly more atherogenic than the normal large buoyant particles, an elevated non-HDL-C indirectly suggests greater proportion of the small, dense variety of LDL particles

Serum lipid profile is measured for cardiovascular risk prediction. Lipid Association of India recommends LDL-C as primary target and Non HDL-C as co-primary treatment target.

Risk Stratification for ASCVD (Atherosclerotic cardiovascular disease) by Lipid Association of India

Risk Category	
Extreme risk group	A. CAD with > 1 feature of high risk group B. CAD with > 1 feature of Very high risk group or recurrent ACS (within 1 year) despite LDL-C < or = 50 mg/dl or polyvascular disease
Very High Risk	1. Established ASCVD 2. Diabetes with 2 major risk factors or evidence of end organ damage 3. Familial Homozygous Hypercholesterolemia
High Risk	1. Three major ASCVD risk factors. 2. Diabetes with 1 major risk factor or no evidence of end organ damage. 3. CKD stage 3B or 4. 4. LDL >190 mg/dl 5. Extreme of a single risk factor. 6. Coronary Artery Calcium - CAC >300 AU. 7. Lipoprotein a >= 50mg/dl 8. Non stenotic carotid plaque
Moderate Risk	2 major ASCVD risk factors
Low Risk	0-1 major ASCVD risk factors
Major ASCVD (Atherosclerotic cardiovascular disease) Risk Factors	
1. Age > or = 45 years in males and > or = 55 years in females	3. Current Cigarette smoking or tobacco use
2. Family history of premature ASCVD	4. High blood pressure
5. Low HDL	

Newer treatment goals and statin initiation thresholds based on the risk categories proposed by LAI in 2020.

Risk Group	Treatment Goals		Consider Drug Therapy	
	LDL-C (mg/dl)	Non-HDL (mg/dl)	LDL-C (mg/dl)	Non-HDL (mg/dl)
Extreme Risk Group Category A	<50 (Optional goal <OR = 30)	< 80 (Optional goal <OR = 60)	>OR = 50	>OR = 80



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075

AGE : 36 Years

SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Extreme Risk Group Category B	<OR = 30	<OR = 60	> 30	>60
Very High Risk	<50	<80	>OR= 50	>OR= 80
High Risk	<70	<100	>OR= 70	>OR= 100
Moderate Risk	<100	<130	>OR= 100	>OR= 130
Low Risk	<100	<130	>OR= 130*	>OR= 160

*After an adequate non-pharmacological intervention for at least 3 months.

References: Management of Dyslipidaemia for the Prevention of Stroke: Clinical Practice Recommendations from the Lipid Association of India. Current Vascular Pharmacology, 2022, 20, 134-155.

LIVER FUNCTION PROFILE, SERUM

BILIRUBIN, TOTAL	0.28		Upto 1.2	mg/dL
METHOD : COLORIMETRIC DIAZO				
BILIRUBIN, DIRECT	0.19		< 0.30	mg/dL
BILIRUBIN, INDIRECT	0.09	Low	0.1 - 1.0	mg/dL
TOTAL PROTEIN	7.5		6.0 - 8.0	g/dL
METHOD : COLORIMETRIC				
ALBUMIN	4.4		3.97 - 4.94	g/dL
METHOD : COLORIMETRIC				
GLOBULIN	3.1		2.0 - 3.5	g/dL
ALBUMIN/GLOBULIN RATIO	1.4		1.0 - 2.1	RATIO
ASPARTATE AMINOTRANSFERASE (AST/SGOT)	21		< OR = 35	U/L
METHOD : UV ABSORBANCE				
ALANINE AMINOTRANSFERASE (ALT/SGPT)	20		< OR = 35	U/L
METHOD : UV ABSORBANCE				
ALKALINE PHOSPHATASE	62		35 - 104	U/L
METHOD : COLORIMETRIC				
GAMMA GLUTAMYL TRANSFERASE (GGT)	13		0 - 40	U/L
METHOD : ENZYMATIC, COLORIMETRIC				
LACTATE DEHYDROGENASE	197		125 - 220	U/L
METHOD : UV ABSORBANCE				

BLOOD UREA NITROGEN (BUN), SERUM

BLOOD UREA NITROGEN	15		6 - 20	mg/dL
METHOD : ENZYMATIC ASSAY				

CREATININE, SERUM

CREATININE	0.63		0.5 - 0.9	mg/dL
METHOD : COLORIMETRIC				

BUN/CREAT RATIO



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 S.K. Tower, Hari Niwas, LBS Marg
 THANE, 400602
 MAHARASHTRA, INDIA
 Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
 Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 **AGE :** 36 Years **SEX :** Female

ABHA NO :

DRAWN : **RECEIVED :** 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

BUN/CREAT RATIO	23.81	High	8.0 - 15.0	
URIC ACID, SERUM				
URIC ACID	6.1	High	2.4 - 5.7	mg/dL
METHOD : ENZYMATIC COLORIMETRIC ASSAY				
TOTAL PROTEIN, SERUM				
TOTAL PROTEIN	7.5		6.0 - 8.0	g/dL
METHOD : COLORIMETRIC				
ALBUMIN, SERUM				
ALBUMIN	4.4		3.97 - 4.94	g/dL
METHOD : COLORIMETRIC				
GLOBULIN				
GLOBULIN	3.1		2.0 - 3.5	g/dL
ELECTROLYTES (NA/K/CL), SERUM				
SODIUM, SERUM	140		136 - 145	mmol/L
POTASSIUM, SERUM	4.61		3.5 - 5.1	mmol/L
CHLORIDE, SERUM	104		98 - 107	mmol/L

Interpretation(s)

Sodium	Potassium	Chloride
Decreased in: CCF, cirrhosis, vomiting, diarrhea, excessive sweating, salt-losing nephropathy, adrenal insufficiency, nephrotic syndrome, water intoxication, SIADH. Drugs: thiazides, diuretics, ACE inhibitors, chlorpropamide, carbamazepine, antidepressants (SSRI), antipsychotics.	Decreased in: Low potassium intake, prolonged vomiting or diarrhea, RTA types I and II, hyperaldosteronism, Cushing's syndrome, osmotic diuresis (e.g., hyperglycemia), alkalosis, familial periodic paralysis, trauma (transient). Drugs: Adrenergic agents, diuretics.	Decreased in: Vomiting, diarrhea, renal failure combined with salt deprivation, over-treatment with diuretics, chronic respiratory acidosis, diabetic ketoacidosis, excessive sweating, SIADH, salt-losing nephropathy, porphyria, expansion of extracellular fluid volume, adrenal insufficiency, hyperaldosteronism, metabolic alkalosis. Drugs: chronic laxative, corticosteroids, diuretics.
Increased in: Dehydration (excessive sweating, severe vomiting or diarrhea), diabetes mellitus, diabetes insipidus, hyperaldosteronism, inadequate water intake. Drugs: steroids, licorice, oral contraceptives.	Increased in: Massive hemolysis, severe tissue damage, rhabdomyolysis, acidosis, dehydration, renal failure, Addison's disease, RTA type IV, hyperkalemic familial periodic paralysis. Drugs: potassium salts, potassium-sparing diuretics, NSAIDs, beta-blockers, ACE inhibitors, high-dose trimethoprim-sulfamethoxazole.	Increased in: Renal failure, nephrotic syndrome, RTA, dehydration, overtreatment with saline, hyperparathyroidism, diabetes insipidus, metabolic acidosis from diarrhea (Loss of HCO ₃ ⁻), respiratory alkalosis, hyperadrenocorticism. Drugs: acetazolamide, androgens, hydrochlorothiazide, salicylates.
Interferences: Severe lipemia or hyperproteinemia, if sodium analysis involves a dilution step can cause spurious results. The serum sodium falls about 1.6 mEq/L for each 100 mg/dL increase in blood glucose.	Interferences: Hemolysis of sample, delayed separation of serum, prolonged fist clenching during blood drawing, and prolonged tourniquet placement. Very high WBC/PLT counts may cause spurious. Plasma potassium levels are normal.	Interferences: Test is helpful in assessing normal and increased anion gap metabolic acidosis and in distinguishing hypercalcemia due to hyperparathyroidism (high serum chloride) from that due to malignancy (Normal serum chloride)



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : **SWAGF260786181**ACCESSION NO : **0181WB00075** AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

PHYSICAL EXAMINATION, URINE

COLOR PALE YELLOW
APPEARANCE SLIGHTLY HAZY

CHEMICAL EXAMINATION, URINE

PH	5.5	5.00 - 7.50
SPECIFIC GRAVITY	1.020	1.010 - 1.030
PROTEIN	NOT DETECTED	NOT DETECTED
GLUCOSE	NOT DETECTED	NOT DETECTED
KETONES	NOT DETECTED	NOT DETECTED
BLOOD	NOT DETECTED	NOT DETECTED
UROBILINOGEN	NORMAL	NORMAL
NITRITE	NOT DETECTED	NOT DETECTED
LEUKOCYTE ESTERASE	NOT DETECTED	NOT DETECTED

MICROSCOPIC EXAMINATION, URINE

RED BLOOD CELLS	NOT DETECTED	NOT DETECTED	/HPF
PUS CELL (WBC'S)	3-5	0-5	/HPF
EPITHELIAL CELLS	8-10	0-5	/HPF
CASTS	NOT DETECTED		
CRYSTALS	NOT DETECTED		
BACTERIA	NOT DETECTED	NOT DETECTED	
YEAST	NOT DETECTED	NOT DETECTED	



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 AGE : 36 Years SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

The following table describes the probable conditions, in which the analytes are present in urine

Presence of	Conditions
Proteins	Inflammation or immune illnesses
Pus (White Blood Cells)	Urinary tract infection, urinary tract or kidney stone, tumors or any kind of kidney impairment
Glucose	Diabetes or kidney disease
Ketones	Diabetic ketoacidosis (DKA), starvation or thirst
Urobilinogen	Liver disease such as hepatitis or cirrhosis
Blood	Renal or genital disorders/trauma
Bilirubin	Liver disease
Erythrocytes	Urological diseases (e.g. kidney and bladder cancer, urolithiasis), urinary tract infection and glomerular diseases
Leukocytes	Urinary tract infection, glomerulonephritis, interstitial nephritis either acute or chronic, polycystic kidney disease, urolithiasis, contamination by genital secretions
Epithelial cells	Urolithiasis, bladder carcinoma or hydronephrosis, ureteric stents or bladder catheters for prolonged periods of time
Granular Casts	Low intratubular pH, high urine osmolality and sodium concentration, interaction with Bence-Jones protein
Hyaline casts	Physical stress, fever, dehydration, acute congestive heart failure, renal diseases
Calcium oxalate	Metabolic stone disease, primary or secondary hyperoxaluria, intravenous infusion of large doses of vitamin C, the use of vasodilator naftidrofuryl oxalate or the gastrointestinal lipase inhibitor orlistat, ingestion of ethylene glycol or of star fruit (Averrhoa carambola) or its juice
Uric acid	arthritis
Bacteria	Urinary infection when present in significant numbers & with pus cells.
Trichomonas vaginalis	Vaginitis, cervicitis or salpingitis

THYROID PANEL, SERUM

T3

102.0

Non-Pregnant Women

ng/dL

80.0 - 200.0

Pregnant Women

1st Trimester: 105.0 - 230.0

2nd Trimester: 129.0 - 262.0

3rd Trimester: 135.0 - 262.0

METHOD : ELECTROCHEMILUMINESCENCE



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 S.K. Tower, Hari Niwas, LBS Marg
 THANE, 400602
 MAHARASHTRA, INDIA
 Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
 Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 **AGE :** 36 Years **SEX :** Female

ABHA NO :

DRAWN : **RECEIVED :** 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
T4		8.22	Non-Pregnant Women 5.10 - 14.10 Pregnant Women 1st Trimester: 7.33 - 14.80 2nd Trimester: 7.93 - 16.10 3rd Trimester: 6.95 - 15.70	µg/dL
METHOD : ELECTROCHEMILUMINESCENCE				
TSH (ULTRASENSITIVE)		1.300	Non Pregnant Women 0.27 - 4.20 Pregnant Women 1st Trimester: 0.33 - 4.59 2nd Trimester: 0.35 - 4.10 3rd Trimester: 0.21 - 3.15	µIU/mL
METHOD : ELECTROCHEMILUMINESCENCE				





CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
S.K. Tower, Hari Niwas, LBS Marg
THANE, 400602
MAHARASHTRA, INDIA
Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075

AGE : 36 Years

SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

Interpretation(s)

Triiodothyronine T3, **Thyroxine T4**, and **Thyroid Stimulating Hormone TSH** are thyroid hormones which affect almost every physiological process in the body, including growth, development, metabolism, body temperature, and heart rate.

Production of T3 and its prohormone thyroxine (T4) is activated by thyroid-stimulating hormone (TSH), which is released from the pituitary gland. Elevated concentrations of T3, and T4 in the blood inhibit the production of TSH.

Excessive secretion of thyroxine in the body is hyperthyroidism, and deficient secretion is called hypothyroidism.

In primary hypothyroidism, TSH levels are significantly elevated, while in secondary and tertiary hyperthyroidism, TSH levels are low.

Below mentioned are the guidelines for Pregnancy related reference ranges for Total T4, TSH & Total T3. Measurement of the serum TT3 level is a more sensitive test for the diagnosis of hyperthyroidism, and measurement of TT4 is more useful in the diagnosis of hypothyroidism. Most of the thyroid hormone in blood is bound to transport proteins. Only a very small fraction of the circulating hormone is free and biologically active. It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4.

Sr. No.	TSH	Total T4	FT4	Total T3	Possible Conditions
1	High	Low	Low	Low	(1) Primary Hypothyroidism (2) Chronic autoimmune Thyroiditis (3) Post Thyroidectomy (4) Post Radio-Iodine treatment
2	High	Normal	Normal	Normal	(1) Subclinical Hypothyroidism (2) Patient with insufficient thyroid hormone replacement therapy (3) In cases of Autoimmune/Hashimoto thyroiditis (4). Isolated increase in TSH levels can be due to Subclinical inflammation, drugs like amphetamines, Iodine containing drug and dopamine antagonist e.g. domperidone and other physiological reasons.
3	Normal/Low	Low	Low	Low	(1) Secondary and Tertiary Hypothyroidism
4	Low	High	High	High	(1) Primary Hyperthyroidism (Graves Disease) (2) Multinodular Goitre (3) Toxic Nodular Goitre (4) Thyroiditis (5) Over treatment of thyroid hormone (6) Drug effect e.g. Glucocorticoids, dopamine, T4 replacement therapy (7) First trimester of Pregnancy
5	Low	Normal	Normal	Normal	(1) Subclinical Hyperthyroidism
6	High	High	High	High	(1) TSH secreting pituitary adenoma (2) TRH secreting tumor
7	Low	Low	Low	Low	(1) Central Hypothyroidism (2) Euthyroid sick syndrome (3) Recent treatment for Hyperthyroidism
8	Normal/Low	Normal	Normal	High	(1) T3 thyrotoxicosis (2) Non-Thyroidal illness
9	Low	High	High	Normal	(1) T4 Ingestion (2) Thyroiditis (3) Interfering Anti TPO antibodies

REF: 1. TIETZ Fundamentals of Clinical chemistry 2. Guidelines of the American Thyroid association during pregnancy and Postpartum, 2011.

NOTE: It is advisable to detect Free T3, Free T4 along with TSH, instead of testing for albumin bound Total T3, Total T4. TSH is not affected by variation in thyroid - binding protein. TSH has a diurnal rhythm, with peaks at 2:00 - 4:00 a.m. And troughs at 5:00 - 6:00 p.m. With ultradian variations.

PAPANICOLAOU SMEAR

TEST METHOD

CONVENTIONAL GYNEC CYTOLOGY

METHOD : MICROSCOPIC EXAMINATION

SPECIMEN TYPE

P-268/23

METHOD : MICROSCOPIC EXAMINATION

TWO UNSTAINED CERVICAL SMEARS RECEIVED

REPORTING SYSTEM

2014 BETHESDA SYSTEM FOR REPORTING CERVICAL CYTOLOGY



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002381402

CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075

AGE : 36 Years

SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

SPECIMEN ADEQUACY

SATISFACTORY

METHOD : PAP STAIN & MICROSCOPIC EXAMINATION

MICROSCOPY

THE SMEARS SHOW MAINLY SUPERFICIAL SQUAMOUS CELLS, FEW INTERMEDIATE SQUAMOUS CELLS AND MANY CLUSTERS OF ENDOCERVICAL CELLS IN THE BACKGROUND OF MODERATE POLYMORPHS & RBC'S.

METHOD : PAP STAIN

INTERPRETATION / RESULT

NEGATIVE FOR INTRAEPITHELIAL LESION OR MALIGNANCY WITH MODERATE INFLAMMATION

METHOD : PAP STAIN & MICROSCOPIC EXAMINATION

- REACTIVE CELLULAR CHANGES ASSOCIATED WITH INFLAMMATION.

METHOD : PAP STAIN & MICROSCOPIC EXAMINATION

Comments

PLEASE NOTE PAPANICOLAU SMEAR STUDY IS A SCREENING PROCEDURE FOR CERVICAL CANCER WITH INHERENT FALSE NEGATIVE RESULTS HENCE SHOULD BE INTERPRETED WITH CAUTION. NO CYTOLOGICAL EVIDENCE OF HPV INFECTION IN THE SMEARS STUDIED. REPEAT PAP SMEAR EXAMINATION AFTER TREATMENT OF INFLAMMATION. SMEARS WILL BE PRESERVED FOR 5 YEARS ONLY.

PHYSICAL EXAMINATION, STOOL

COLOUR

SAMPLE NOT RECEIVED

METHOD : VISUAL

ABO GROUP & RH TYPE, EDTA WHOLE BLOOD

ABO GROUP

TYPE A

METHOD : GEL COLUMN AGGLUTINATION METHOD.

RH TYPE

POSITIVE

METHOD : GEL COLUMN AGGLUTINATION METHOD.

XRAY-CHEST

IMPRESSION

NO ABNORMALITY DETECTED

TMT OR ECHO

TMT OR ECHO

NEGATIVE

ECG

ECG

WITHIN NORMAL LIMITS

MEDICAL HISTORY

RELEVANT PRESENT HISTORY

NOT SIGNIFICANT

RELEVANT PAST HISTORY

NOT SIGNIFICANT

RELEVANT PERSONAL HISTORY

MARRIED / CHILD / MIXED DIET / NO ALLERGIES / NO SMOKING / NO ALCOHOL.

LMP (FOR FEMALES)

10.02.2023



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075

AGE : 36 Years

SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

OBSTETRIC HISTORY (FOR FEMALES)

LSCS 2021.

RELEVANT FAMILY HISTORY

FATHER:- HEART DISEASE / DIABETES.

HISTORY OF MEDICATIONS

NOT SIGNIFICANT

ANTHROPOMETRIC DATA & BMI

HEIGHT IN METERS

1.57

mts

WEIGHT IN KGS.

85

Kgs

BMI

34

BMI & Weight Status as follows: kg/sqmts

Below 18.5: Underweight

18.5 - 24.9: Normal

25.0 - 29.9: Overweight

30.0 and Above: Obese

GENERAL EXAMINATION

MENTAL / EMOTIONAL STATE

NORMAL

PHYSICAL ATTITUDE

NORMAL

GENERAL APPEARANCE / NUTRITIONAL STATUS

OBESE

BUILT / SKELETAL FRAMEWORK

AVERAGE

FACIAL APPEARANCE

NORMAL

SKIN

NORMAL

UPPER LIMB

NORMAL

LOWER LIMB

NORMAL

NECK

NORMAL

NECK LYMPHATICS / SALIVARY GLANDS

NOT ENLARGED OR TENDER

THYROID GLAND

NOT ENLARGED

CAROTID PULSATION

NORMAL

BREAST (FOR FEMALES)

NORMAL

TEMPERATURE

NORMAL

PULSE

72/MIN.REGULAR, ALL PERIPHERAL PULSES WELL FELT, NO CAROTID BRUIT

RESPIRATORY RATE

NORMAL

CARDIOVASCULAR SYSTEM

BP

112/88 MM HG
(SUPINE)

mm/Hg

PERICARDIUM

NORMAL

APEX BEAT

NORMAL

HEART SOUNDS

NORMAL

MURMURS

ABSENT

RESPIRATORY SYSTEM



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075

AGE : 36 Years

SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

SIZE AND SHAPE OF CHEST	NORMAL
MOVEMENTS OF CHEST	SYMMETRICAL
BREATH SOUNDS INTENSITY	NORMAL
BREATH SOUNDS QUALITY	VESICULAR (NORMAL)
ADDED SOUNDS	ABSENT

PER ABDOMEN

APPEARANCE	NORMAL
VENOUS PROMINENCE	ABSENT
LIVER	NOT PALPABLE
SPLEEN	NOT PALPABLE
HERNIA	ABSENT

CENTRAL NERVOUS SYSTEM

HIGHER FUNCTIONS	NORMAL
CRANIAL NERVES	NORMAL
CEREBELLAR FUNCTIONS	NORMAL
SENSORY SYSTEM	NORMAL
MOTOR SYSTEM	NORMAL
REFLEXES	NORMAL

MUSCULOSKELETAL SYSTEM

SPINE	NORMAL
JOINTS	NORMAL

BASIC EYE EXAMINATION

CONJUNCTIVA	NORMAL
EYELIDS	NORMAL
EYE MOVEMENTS	NORMAL
CORNEA	NORMAL
DISTANT VISION RIGHT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT
DISTANT VISION LEFT EYE WITHOUT GLASSES	REDUCED VISUAL ACUITY 6/12
NEAR VISION RIGHT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT
NEAR VISION LEFT EYE WITHOUT GLASSES	WITHIN NORMAL LIMIT
COLOUR VISION	NORMAL

SUMMARY

RELEVANT HISTORY	NOT SIGNIFICANT
RELEVANT GP EXAMINATION FINDINGS	NOT SIGNIFICANT



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd
S.K. Tower, Hari Niwas, LBS Marg
THANE, 400602
MAHARASHTRA, INDIA
Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 AGE : 36 Years SEX : Female ABHA NO :

DRAWN : RECEIVED : 18/02/2023 09:04 REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

REMARKS / RECOMMENDATIONS

GYNAC CONSULT FOR VAGINITIS & CERVICAL EROSION.
AVOID HIGH QUALITY PROTIEN DIET.
OPHTHALMOLOGY CONSULT FOR REDUCED VISUAL ACUITY.
REPEAT URIC ACID AFTER 3 MONTHS OF DIET AND EXERCISE.
WEIGHT LOSS:- LOW CALORIE, HIGH FIBRE DIET, REGULAR EXERCISE.

Interpretation(s)

BLOOD COUNTS, EDTA WHOLE BLOOD-The cell morphology is well preserved for 24hrs. However after 24-48 hrs a progressive increase in MCV and HCT is observed leading to a decrease in MCHC. A direct smear is recommended for an accurate differential count and for examination of RBC morphology.

RBC AND PLATELET INDICES-Mentzer index (MCV/RBC) is an automated cell-counter based calculated screen tool to differentiate cases of Iron deficiency anaemia(>13) from Beta thalassaemia trait

(<13) in patients with microcytic anaemia. This needs to be interpreted in line with clinical correlation and suspicion. Estimation of HbA2 remains the gold standard for diagnosing a case of beta thalassaemia trait.

WBC DIFFERENTIAL COUNT-The optimal threshold of 3.3 for NLR showed a prognostic possibility of clinical symptoms to change from mild to severe in COVID positive patients. When age = 49.5 years old and NLR = 3.3, 46.1% COVID-19 patients with mild disease might become severe. By contrast, when age < 49.5 years old and NLR < 3.3, COVID-19 patients tend to show mild disease.

(Reference to - The diagnostic and predictive role of NLR, d-NLR and PLR in COVID-19 patients A.-P. Yang, et al. International Immunopharmacology 84 (2020) 106504 This ratio element is a calculated parameter and out of NABL scope.

ERYTHROCYTE SEDIMENTATION RATE (ESR), WHOLE BLOOD-TEST DESCRIPTION :-

Erythrocyte sedimentation rate (ESR) is a test that indirectly measures the degree of inflammation present in the body. The test actually measures the rate of fall (sedimentation) of erythrocytes in a sample of blood that has been placed into a tall, thin, vertical tube. Results are reported as the millimetres of clear fluid (plasma) that are present at the top portion of the tube after one hour. Nowadays fully automated instruments are available to measure ESR.

ESR is not diagnostic it is a non-specific test that may be elevated in a number of different conditions. It provides general information about the presence of an inflammatory condition. CRP is superior to ESR because it is more sensitive and reflects a more rapid change.

TEST INTERPRETATION

Increase in: Infections, Vasculitides, Inflammatory arthritis, Renal disease, Anemia, Malignancies and plasma cell dyscrasias, Acute allergy Tissue injury, Pregnancy, Estrogen medication, Aging.

Finding a very accelerated ESR(>100 mm/hour) in patients with ill-defined symptoms directs the physician to search for a systemic disease (Paraproteinemias, Disseminated malignancies, connective tissue disease, severe infections such as bacterial endocarditis).

In pregnancy BRI in first trimester is 0-48 mm/hr(62 if anemic) and in second trimester (0-70 mm/hr(95 if anemic). ESR returns to normal 4th week post partum.

Decreased in: Polycythemia vera, Sickle cell anemia

LIMITATIONS

False elevated ESR : Increased fibrinogen, Drugs(Vitamin A, Dextran etc), Hypercholesterolemia

False Decreased : Poikilocytosis,(SickleCells,spherocytes),Microcytosis, Low fibrinogen, Very high WBC counts, Drugs(Quinine,

salicylates)

REFERENCE :

1. Nathan and Oski's Haematology of Infancy and Childhood, 5th edition 2. Paediatric reference intervals. AACCPress, 7th edition. Edited by S. Soldin 3. The reference for the adult reference range is "Practical Haematology by Dacie and Lewis, 10th edition.

GLUCOSE FASTING, FLUORIDE PLASMA-TEST DESCRIPTION

Normally, the glucose concentration in extracellular fluid is closely regulated so that a source of energy is readily available to tissues and so that no glucose is excreted in the urine.

Increased in

Diabetes mellitus, Cushing's syndrome (10 - 15%), chronic pancreatitis (30%). Drugs:corticosteroids, phenytoin, estrogen, thiazides.

Decreased in

Pancreatic islet cell disease with increased insulin, insulinoma, adrenocortical insufficiency, hypopituitarism, diffuse liver disease, malignancy (adrenocortical, stomach, fibrosarcoma), infant of a diabetic mother, enzyme deficiency diseases(e.g., galactosemia), Drugs- insulin, ethanol, propranolol sulfonyleureas, tolbutamide, and other oral hypoglycemic agents.

NOTE: While random serum glucose levels correlate with home glucose monitoring results (weekly mean capillary glucose values), there is wide fluctuation within individuals. Thus, glycosylated hemoglobin(HbA1c) levels are favored to monitor glycemic control.

High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc.

GLYCOSYLATED HEMOGLOBIN(HBA1C), EDTA WHOLE BLOOD-Used For:

1. Evaluating the long-term control of blood glucose concentrations in diabetic patients.



Scan to View Details



Scan to View Report



CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 S.K. Tower, Hari Niwas, LBS Marg
 THANE, 400602
 MAHARASHTRA, INDIA
 Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
 Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075 **AGE :** 36 Years **SEX :** Female **ABHA NO :**

DRAWN : **RECEIVED :** 18/02/2023 09:04 **REPORTED :** 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status	Final	Results	Biological Reference Interval	Units
--------------------	-------	---------	-------------------------------	-------

2.Diagnosing diabetes.

3. Identifying patients at increased risk for diabetes (prediabetes).

The ADA recommends measurement of HbA1c (typically 3-4 times per year for type 1 and poorly controlled type 2 diabetic patients, and 2 times per year for well-controlled type 2 diabetic patients) to determine whether a patient's metabolic control has remained continuously within the target range.

1. eAG (Estimated average glucose) converts percentage HbA1c to mg/dl, to compare blood glucose levels.

2. eAG gives an evaluation of blood glucose levels for the last couple of months.

3. eAG is calculated as $eAG (mg/dl) = 28.7 * HbA1c - 46.7$

HbA1c Estimation can get affected due to :

I. Shortened Erythrocyte survival : Any condition that shortens erythrocyte survival or decreases mean erythrocyte age (e.g. recovery from acute blood loss, hemolytic anemia) will falsely lower HbA1c test results. Fructosamine is recommended in these patients which indicates diabetes control over 15 days.

II. Vitamin C & E are reported to falsely lower test results. (possibly by inhibiting glycation of hemoglobin).

III. Iron deficiency anemia is reported to increase test results. Hypertriglyceridemia, uremia, hyperbilirubinemia, chronic alcoholism, chronic ingestion of salicylates & opiates addition are reported to interfere with some assay methods, falsely increasing results.

IV. Interference of hemoglobinopathies in HbA1c estimation is seen in
 a. Homozygous hemoglobinopathy. Fructosamine is recommended for testing of HbA1c.

b. Heterozygous state detected (D10 is corrected for HbS & HbC trait.)

c. HbF > 25% on alternate platform (Boronate affinity chromatography) is recommended for testing of HbA1c. Abnormal Hemoglobin electrophoresis (HPLC method) is recommended for detecting a hemoglobinopathy

GLUCOSE, POST-PRANDIAL, PLASMA-High fasting glucose level in comparison to post prandial glucose level may be seen due to effect of Oral Hypoglycaemics & Insulin treatment, Renal Glycosuria, Glycaemic index & response to food consumed, Alimentary Hypoglycemia, Increased insulin response & sensitivity etc. Additional test HbA1c LIVER FUNCTION PROFILE, SERUM-LIVER FUNCTION PROFILE

Bilirubin is a yellowish pigment found in bile and is a breakdown product of normal heme catabolism. Bilirubin is excreted in bile and urine, and elevated levels may give yellow discoloration in jaundice. Elevated levels result from increased bilirubin production (eg, hemolysis and ineffective erythropoiesis), decreased bilirubin excretion (eg, obstruction and hepatitis), and abnormal bilirubin metabolism (eg, hereditary and neonatal jaundice). Conjugated (direct) bilirubin is elevated more than unconjugated (indirect) bilirubin in Viral hepatitis, Drug reactions, Alcoholic liver disease. Conjugated (direct) bilirubin is also elevated more than unconjugated (indirect) bilirubin when there is some kind of blockage of the bile ducts like in Gallstones getting into the bile ducts, tumors & Scarring of the bile ducts. Increased unconjugated (indirect) bilirubin may be a result of Hemolytic or pernicious anemia, Transfusion reaction & a common metabolic condition termed Gilbert syndrome, due to low levels of the enzyme that attaches sugar molecules to bilirubin.

AST is an enzyme found in various parts of the body. AST is found in the liver, heart, skeletal muscle, kidneys, brain, and red blood cells, and it is commonly measured clinically as a marker for liver health. AST levels increase during chronic viral hepatitis, blockage of the bile duct, cirrhosis of the liver, liver cancer, kidney failure, hemolytic anemia, pancreatitis, hemochromatosis. AST levels may also increase after a heart attack or strenuous activity. ALT test measures the amount of this enzyme in the blood. ALT is found mainly in the liver, but also in smaller amounts in the kidneys, heart, muscles, and pancreas. It is commonly measured as a part of a diagnostic evaluation of hepatocellular injury, to determine liver health. AST levels increase during acute hepatitis, sometimes due to a viral infection, ischemia to the liver, chronic hepatitis, obstruction of bile ducts, cirrhosis.

ALP is a protein found in almost all body tissues. Tissues with higher amounts of ALP include the liver, bile ducts and bone. Elevated ALP levels are seen in Biliary obstruction, Osteoblastic bone tumors, osteomalacia, hepatitis, Hyperparathyroidism, Leukemia, Lymphoma, Paget's disease, Rickets, Sarcoidosis etc. Lower-than-normal ALP levels seen in Hypophosphatasia, Malnutrition, Protein deficiency, Wilson's disease. GGT is an enzyme found in cell membranes of many tissues mainly in the liver, kidney and pancreas. It is also found in other tissues including intestine, spleen, heart, brain and seminal vesicles. The highest concentration is in the kidney, but the liver is considered the source of normal enzyme activity. Serum GGT has been widely used as an index of liver dysfunction. Elevated serum GGT activity can be found in diseases of the liver, biliary system and pancreas. Conditions that increase serum GGT are obstructive liver disease, high alcohol consumption and use of enzyme-inducing drugs etc. Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin. Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease. Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc. Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc

BLOOD UREA NITROGEN (BUN), SERUM- Causes of Increased levels include Pre renal (High protein diet, Increased protein catabolism, GI haemorrhage, Cortisol, Dehydration, CHF Renal), Renal Failure, Post Renal (Malignancy, Nephrolithiasis, Prostatism)

Causes of decreased level include Liver disease, SIADH.

CREATININE, SERUM- Higher than normal level may be due to:

- Blockage in the urinary tract
- Kidney problems, such as kidney damage or failure, infection, or reduced blood flow
- Loss of body fluid (dehydration)
- Muscle problems, such as breakdown of muscle fibers
- Problems during pregnancy, such as seizures (eclampsia), or high blood pressure caused by pregnancy (preeclampsia)

Lower than normal level may be due to:

- Myasthenia Gravis
- Muscular dystrophy

URIC ACID, SERUM- Causes of Increased levels:- Dietary (High Protein Intake, Prolonged Fasting, Rapid weight loss), Gout, Lesch nyhan syndrome, Type 2 DM, Metabolic syndrome

Causes of decreased levels- Low Zinc intake, OCP, Multiple Sclerosis

TOTAL PROTEIN, SERUM- Serum total protein, also known as total protein, is a biochemical test for measuring the total amount of protein in serum. Protein in the plasma is made up of albumin and globulin

Higher-than-normal levels may be due to: Chronic inflammation or infection, including HIV and hepatitis B or C, Multiple myeloma, Waldenstrom's disease





CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :
 ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
 F-703, F-703, LADO SARAI, MEHRAULI
 SOUTH WEST DELHI
 NEW DELHI 110030
 DELHI INDIA
 8800465156

SRL Ltd
 S.K. Tower, Hari Niwas, LBS Marg
 THANE, 400602
 MAHARASHTRA, INDIA
 Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956
 Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA **PATIENT ID : SWAGF260786181**

ACCESSION NO : 0181WB00075 **AGE : 36 Years** **SEX : Female** **ABHA NO :**

DRAWN : **RECEIVED : 18/02/2023 09:04** **REPORTED : 22/02/2023 14:52**

REFERRING DOCTOR : SELF **CLIENT PATIENT ID :**

Test Report Status	Results	Biological Reference Interval	Units
Final			

Lower-than-normal levels may be due to: Agammaglobulinemia, Bleeding (hemorrhage), Burns, Glomerulonephritis, Liver disease, Malabsorption, Malnutrition, Nephrotic syndrome, Protein-losing enteropathy etc.
 ALBUMIN, SERUM-Human serum albumin is the most abundant protein in human blood plasma. It is produced in the liver. Albumin constitutes about half of the blood serum protein. Low blood albumin levels (hypoalbuminemia) can be caused by: Liver disease like cirrhosis of the liver, nephrotic syndrome, protein-losing enteropathy, Burns, hemodilution, increased vascular permeability or decreased lymphatic clearance, malnutrition and wasting etc.
 ABO GROUP & RH TYPE, EDTA WHOLE BLOOD-Blood group is identified by antigens and antibodies present in the blood. Antigens are protein molecules found on the surface of red blood cells. Antibodies are found in plasma. To determine blood group, red cells are mixed with different antibody solutions to give A, B, O or AB.

Disclaimer: "Please note, as the results of previous ABO and Rh group (Blood Group) for pregnant women are not available, please check with the patient records for availability of the same."

The test is performed by both forward as well as reverse grouping methods.

MEDICAL HISTORY

 THIS REPORT CARRIES THE SIGNATURE OF OUR LABORATORY DIRECTOR. THIS IS AN INVIOABLE FEATURE OF OUR LAB MANAGEMENT SOFTWARE. HOWEVER, ALL EXAMINATIONS AND INVESTIGATIONS HAVE BEEN CONDUCTED BY OUR PANEL OF DOCTORS.



Scan to View Details



Scan to View Report



Patient Ref. No. 775000002381402

CLIENT CODE : C000138394

CLIENT'S NAME AND ADDRESS :

ACROFEMI HEALTHCARE LTD (MEDIWHEEL)
F-703, F-703, LADO SARAI, MEHRAULI
SOUTH WEST DELHI
NEW DELHI 110030
DELHI INDIA
8800465156

SRL Ltd

S.K. Tower, Hari Niwas, LBS Marg

THANE, 400602

MAHARASHTRA, INDIA

Tel : 9111591115, Fax : CIN - U74899PB1995PLC045956

Email : customercare.thane@srl.in

PATIENT NAME : SWAGATA RANA

PATIENT ID : SWAGF260786181

ACCESSION NO : 0181WB00075

AGE : 36 Years

SEX : Female

ABHA NO :

DRAWN :

RECEIVED : 18/02/2023 09:04

REPORTED : 22/02/2023 14:52

REFERRING DOCTOR : SELF

CLIENT PATIENT ID :

Test Report Status

Final

Results

Units

MEDI WHEEL FULL BODY HEALTH CHECKUP BELOW 40FEMALE**ULTRASOUND ABDOMEN****ULTRASOUND ABDOMEN**

UPPER ABDOMEN :- GRADE I FATTY LIVER

****End Of Report****Please visit www.srlworld.com for related Test Information for this accession**CONDITIONS OF LABORATORY TESTING & REPORTING**

1. It is presumed that the test sample belongs to the patient named or identified in the test requisition form.
2. All tests are performed and reported as per the turnaround time stated in the SRL Directory of Services.
3. Result delays could occur due to unforeseen circumstances such as non-availability of kits / equipment breakdown / natural calamities / technical downtime or any other unforeseen event.
4. A requested test might not be performed if:
 - i. Specimen received is insufficient or inappropriate
 - ii. Specimen quality is unsatisfactory
 - iii. Incorrect specimen type
 - iv. Discrepancy between identification on specimen container label and test requisition form
5. SRL confirms that all tests have been performed or assayed with highest quality standards, clinical safety & technical integrity.
6. Laboratory results should not be interpreted in isolation; it must be correlated with clinical information and be interpreted by registered medical practitioners only to determine final diagnosis.
7. Test results may vary based on time of collection, physiological condition of the patient, current medication or nutritional and dietary changes. Please consult your doctor or call us for any clarification.
8. Test results cannot be used for Medico legal purposes.
9. In case of queries please call customer care (91115 91115) within 48 hours of the report.

SRL Limited

Fortis Hospital, Sector 62, Phase VIII,
Mohali 160062



Scan to View Details



Scan to View Report