PID No. : MED121679603 Register On : 522302708 SID No.

: MediWheel

Age / Sex : 37 Year(s) / Male

: OP

Type

Ref. Dr

Report On

: 21/02/2023 9:38 AM

Collection On : 21/02/2023 9:45 AM

: 21/02/2023 3:26 PM

Printed On

: 01/03/2023 1:52 PM



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
HAEMATOLOGY			
Complete Blood Count With - ESR			
Haemoglobin (EDTA Blood/Spectrophotometry)	14.2	g/dL	13.5 - 18.0
Packed Cell Volume(PCV)/Haematocrit (EDTA Blood)	41.5	%	42 - 52
RBC Count (EDTA Blood)	4.92	mill/cu.mm	4.7 - 6.0
Mean Corpuscular Volume(MCV) (EDTA Blood)	84.3	fL	78 - 100
Mean Corpuscular Haemoglobin(MCH) (EDTA Blood)	28.8	pg	27 - 32
Mean Corpuscular Haemoglobin concentration(MCHC) (EDTA Blood)	34.2	g/dL	32 - 36
RDW-CV	14.0	%	11.5 - 16.0
RDW-SD	41.31	fL	39 - 46
Total Leukocyte Count (TC) (EDTA Blood)	6500	cells/cu.mm	4000 - 11000
Neutrophils (Blood)	51.8	%	40 - 75
Lymphocytes (Blood)	37.0	%	20 - 45
Eosinophils (Blood)	3.0	%	01 - 06

7.7



%

APPROVED BY

Monocytes

(Blood)

01 - 10

PID No. : MED121679603 Register On : 21/02/2023 9:38 AM : 522302708 Collection On : 21/02/2023 9:45 AM SID No.

Age / Sex : 37 Year(s) / Male Report On : 21/02/2023 3:26 PM

Type : OP : 01/03/2023 1:52 PM **Printed On**

Ref. Dr : MediWheel



Investigation	Observed <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Basophils (Blood)	0.5	%	00 - 02
INTERPRETATION: Tests done on Automated Five F	Part cell counter. All a	abnormal results are	reviewed and confirmed microscopically.
Absolute Neutrophil count (EDTA Blood)	3.37	10^3 / μ1	1.5 - 6.6
Absolute Lymphocyte Count (EDTA Blood)	2.40	10^3 / μl	1.5 - 3.5
Absolute Eosinophil Count (AEC) (EDTA Blood)	0.20	10^3 / μl	0.04 - 0.44
Absolute Monocyte Count (EDTA Blood)	0.50	10^3 / μl	< 1.0
Absolute Basophil count (EDTA Blood)	0.03	10^3 / μl	< 0.2
Platelet Count (EDTA Blood)	187	10^3 / μl	150 - 450
MPV (Blood)	8.1	fL	7.9 - 13.7
PCT (Automated Blood cell Counter)	0.15	%	0.18 - 0.28
ESR (Erythrocyte Sedimentation Rate) (Citrated Blood)	4	mm/hr	< 15



PID No. : MED121679603 Register On

: 522302708 SID No.

Age / Sex : 37 Year(s) / Male Type

Ref. Dr

: OP

: 21/02/2023 9:38 AM

Collection On : 21/02/2023 9:45 AM

Report On : 21/02/2023 3:26 PM

: 01/03/2023 1:52 PM **Printed On**

: MediWheel

Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> Reference Interval
BIOCHEMISTRY			
Liver Function Test			
Bilirubin(Total) (Serum/DCA with ATCS)	1.37	mg/dL	0.1 - 1.2
Bilirubin(Direct) (Serum/Diazotized Sulfanilic Acid)	0.37	mg/dL	0.0 - 0.3
Bilirubin(Indirect) (Serum/Derived)	1.00	mg/dL	0.1 - 1.0
SGOT/AST (Aspartate Aminotransferase) (Serum/Modified IFCC)	15.29	U/L	5 - 40
SGPT/ALT (Alanine Aminotransferase) (Serum/Modified IFCC)	21.93	U/L	5 - 41
GGT(Gamma Glutamyl Transpeptidase) (Serum/IFCC / Kinetic)	28.29	U/L	< 55
Alkaline Phosphatase (SAP) (Serum/Modified IFCC)	64.4	U/L	53 - 128
Total Protein (Serum/Biuret)	6.74	gm/dl	6.0 - 8.0
Albumin (Serum/Bromocresol green)	4.40	gm/dl	3.5 - 5.2
Globulin (Serum/Derived)	2.34	gm/dL	2.3 - 3.6
A : G RATIO (Serum/Derived)	1.88		1.1 - 2.2

Sr.Consultant Pathologist Reg No: 100674 **APPROVED BY**

PID No. : MED121679603

. WLD121073000

: 522302708

Age / Sex : 37 Year(s) / Male

: OP

Ref. Dr ; MediWheel

SID No.

Type

Register On : 21/02/2023 9:38 AM

Collection On : 21/02/2023 9:45 AM

Report On : 21/02/2023 3:26 PM

Printed On : 01/03/2023 1:52 PM



<u>Investigation</u>	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
<u>Lipid Profile</u>			
Cholesterol Total (Serum/CHOD-PAP with ATCS)	145.65	mg/dL	Optimal: < 200 Borderline: 200 - 239 High Risk: >= 240
Triglycerides (Serum/GPO-PAP with ATCS)	228.59	mg/dL	Optimal: < 150 Borderline: 150 - 199 High: 200 - 499 Very High: >= 500

INTERPRETATION: The reference ranges are based on fasting condition. Triglyceride levels change drastically in response to food, increasing as much as 5 to 10 times the fasting levels, just a few hours after eating. Fasting triglyceride levels show considerable diurnal variation too. There is evidence recommending triglycerides estimation in non-fasting condition for evaluating the risk of heart disease and screening for metabolic syndrome, as non-fasting sample is more representative of the `usual_circulating level of triglycerides during most part of the day.

r · · · · · · · · · · · · · · · · · · ·			
HDL Cholesterol (Serum/Immunoinhibition)	45.74	mg/dL	Optimal(Negative Risk Factor): >= 60 Borderline: 40 - 59 High Risk: < 40
LDL Cholesterol (Serum/Calculated)	54.2	mg/dL	Optimal: < 100 Above Optimal: 100 - 129 Borderline: 130 - 159 High: 160 - 189 Very High: >= 190
VLDL Cholesterol (Serum/Calculated)	45.7	mg/dL	< 30
Non HDL Cholesterol (Serum/Calculated)	99.9	mg/dL	Optimal: < 130 Above Optimal: 130 - 159 Borderline High: 160 - 189 High: 190 - 219 Very High: >= 220



PID No. : MED121679603

Register On : 21/02/2023 9:38 AM

: 522302708 Collection On : 21/02/2023 9:45 AM

Printed On

Age / Sex : 37 Year(s) / Male Report On : 21/02/2023 3:26 PM

Type : OP

SID No.

Ref. Dr : MediWheel



Investigation	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

INTERPRETATION: 1. Non-HDL Cholesterol is now proven to be a better cardiovascular risk marker than LDL Cholesterol. 2.It is the sum of all potentially atherogenic proteins including LDL, IDL, VLDL and chylomicrons and it is the "new bad cholesterol" and is a co-primary target for cholesterol lowering therapy.

: 01/03/2023 1:52 PM

Total Cholesterol/HDL Cholesterol Ratio	3.2	Optimal: < 3.3
(Serum/Calculated)		Low Risk: 3.4 - 4.4

Average Risk: 4.5 - 7.1 Moderate Risk: 7.2 - 11.0 High Risk: > 11.0

Triglyceride/HDL Cholesterol Ratio 5 Optimal: < 2.5

(TG/HDL) Mild to moderate risk: 2.5 - 5.0

High Risk: > 5.0(Serum/Calculated)

LDL/HDL Cholesterol Ratio 1.2 Optimal: 0.5 - 3.0

Borderline: 3.1 - 6.0 (Serum/Calculated) High Risk: > 6.0

> Dr Anusha.K.S Sr.Consultant Pathologist Reg No: 100674

PID No. : MED121679603 Register On : 21/02/2023 9:38 AM

: 522302708 Collection On : 21/02/2023 9:45 AM

Age / Sex : 37 Year(s) / Male Report On : 21/02/2023 3:26 PM

Type : OP

Ref. Dr : MediWheel

SID No.

: 01/03/2023 1:52 PM **Printed On**



Investigation	<u>Observed</u> <u>Value</u>	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>
Glycosylated Haemoglobin (HbA1c)			
HbA1C (Whole Blood/HPLC)	5.6	%	Normal: 4.5 - 5.6 Prediabetes: 5.7 - 6.4 Diabetic: >= 6.5

INTERPRETATION: If Diabetes - Good control: 6.1 - 7.0 %, Fair control: 7.1 - 8.0 %, Poor control >= 8.1 %

Estimated Average Glucose 114.02 mg/dL

(Whole Blood)

INTERPRETATION: Comments

HbA1c provides an index of Average Blood Glucose levels over the past 8 - 12 weeks and is a much better indicator of long term glycemic control as compared to blood and urinary glucose determinations.

Conditions that prolong RBC life span like Iron deficiency anemia, Vitamin B12 & Folate deficiency,

hypertriglyceridemia, hyperbilirubinemia, Drugs, Alcohol, Lead Poisoning, Asplenia can give falsely elevated HbAlC values.

Conditions that shorten RBC survival like acute or chronic blood loss, hemolytic anemia, Hemoglobinopathies, Splenomegaly, Vitamin E ingestion, Pregnancy, End stage Renal disease can cause falsely low HbA1c.

> Sr.Consultant Pathologist Reg No: 100674

PID No. : MED121679603 Register On : 21/02/2023 9:38 AM

: 522302708 SID No. Collection On : 21/02/2023 9:45 AM

Age / Sex : 37 Year(s) / Male Report On : 21/02/2023 3:26 PM

Printed On

Type : OP

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
	Value		Reference Interval

: 01/03/2023 1:52 PM

IMMUNOASSAY

THYROID PROFILE / TFT

T3 (Triiodothyronine) - Total 1.28 ng/ml 0.7 - 2.04

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T3 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T3 is recommended as it is

Metabolically active.

4.2 - 12.0T4 (Tyroxine) - Total 7.85 μg/dl

(Serum/ECLIA)

INTERPRETATION:

Comment:

Total T4 variation can be seen in other condition like pregnancy, drugs, nephrosis etc. In such cases, Free T4 is recommended as it is Metabolically active.

μIU/mL 0.35 - 5.50TSH (Thyroid Stimulating Hormone) 1.94

(Serum/ECLIA)

INTERPRETATION:

Reference range for cord blood - upto 20

1 st trimester: 0.1-2.5 2 nd trimester 0.2-3.0 3 rd trimester: 0.3-3.0

(Indian Thyroid Society Guidelines)

Comment:

- 1.TSH reference range during pregnancy depends on Iodine intake, TPO status, Serum HCG concentration, race, Ethnicity and BMI.
- 2.TSH Levels are subject to circadian variation, reaching peak levels between 2-4am and at a minimum between 6-10PM. The variation can be of the order of 50%, hence time of the day has influence on the measured serum TSH concentrations.
- 3. Values & amplt 0.03 µIU/mL need to be clinically correlated due to presence of rare TSH variant in some individuals.



Type : OP **Printed On** : 01/03/2023 1:52 PM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u>	<u>Unit</u>	<u>Biological</u>
-	<u>Value</u>		Reference Interval
CLINICAL DATIOLOGY			

CLINICAL PATHOLOGY

PHYSICAL EXAMINATION (URINE COMPLETE)

Colour	Yellow	Yellow to Amber

(Urine)

PID No.

Appearance Clear Clear

(Urine)

Volume(CLU) 25

(Urine)

CHEMICAL EXAMINATION (URINE

COMPLETE)

pH 5 4.5 - 8.0

(Urine)

Specific Gravity 1.023 1.002 - 1.035

(Urine)

Ketone Negative Negative

(Urine)

Urobilinogen Normal Normal

(Urine)

Blood Negative Negative

(Urine)

Nitrite Negative Negative

(Urine)

Bilirubin Negative Negative

(Urine)

Protein Negative Negative

(Urine)



 SID No.
 : 522302708
 Collection On
 : 21/02/2023 9:45 AM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 21/02/2023 3:26 PM

 Age / Sex
 : 37 Year(s) / Male
 Report On
 : 21/02/2023 3:26 PM

 Type
 : OP
 Printed On
 : 01/03/2023 1:52 PM

Ref. Dr : MediWheel



<u>Investigation</u>	<u>Observed</u> <u>Uni</u> <u>Value</u>	t <u>Biological</u> <u>Reference Interval</u>
Glucose (Urine/GOD - POD)	Negative	Negative
Leukocytes(CP) (Urine)	Negative	
MICROSCOPIC EXAMINATION (URINE COMPLETE)		

Pus Cells (Urine) NIL (Urine)

Epithelial Cells 0-1 /hpf NIL

(Urine)

RBCs NIL /hpf NIL

(Urine)

Others

(Urine)

INTERPRETATION: Note: Done with Automated Urine Analyser & Automated urine sedimentation analyser. All abnormal reports are reviewed and confirmed microscopically.

Dr Anusha.K.S Sr.Consultant Pathologist Reg No : 100674

: MED121679603

: 522302708

: MediWheel

Age / Sex : 37 Year(s) / Male

Type : OP

PID No.

SID No.

Ref. Dr

Register On : 21/02/2023 9:38 AM

Collection On : 21/02/2023 9:45 AM

Report On : 21/02/2023 3:26 PM

Printed On : 01/03/2023 1:52 PM

Investigation <u>Unit</u> **Biological** <u>Observed</u> Reference Interval <u>Value</u>

IMMUNOHAEMATOLOGY

BLOOD GROUPING AND Rh TYPING 'O' 'Positive'

(EDTA Blood/Agglutination)

INTERPRETATION: Note: Slide method is screening method. Kindly confirm with Tube method for transfusion.

Sr.Consultant Pathologist Reg No: 100674

PID No. : MED121679603

: 522302708 SID No.

Age / Sex : 37 Year(s) / Male

Register On

Report On

21/02/2023 3:26 PM

: 21/02/2023 9:38 AM

Printed On : 01/03/2023 1:52 PM

Collection On : 21/02/2023 9:45 AM

: OP Ref. Dr

Type

: MediWheel



Investigation	Observed Value	<u>Unit</u>	<u>Biological</u> <u>Reference Interval</u>	
BIOCHEMISTRY				
BUN / Creatinine Ratio	9.6		6.0 - 22.0	
Glucose Fasting (FBS) (Plasma - F/GOD-PAP)	108.98	mg/dL	Normal: < 100 Pre Diabetic: 100 - 125 Diabetic: >= 126	

INTERPRETATION: Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level.

legative		Negative
90.97	mg/dL	70 - 140
	C	

(Plasma - PP/GOD-PAP) INTERPRETATION:

Factors such as type, quantity and time of food intake, Physical activity, Psychological stress, and drugs can influence blood glucose level. Fasting blood glucose level may be higher than Postprandial glucose, because of physiological surge in Postprandial Insulin secretion, Insulin resistance, Exercise or Stress, Dawn Phenomenon, Somogyi Phenomenon, Anti- diabetic medication during treatment for Diabetes.

Urine Glucose(PP-2 hours) (Urine - PP)	Negative		Negative
Blood Urea Nitrogen (BUN) (Serum/Urease UV / derived)	8.9	mg/dL	7.0 - 21
Creatinine (Serum/Modified Jaffe)	0.92	mg/dL	0.9 - 1.3

INTERPRETATION: Elevated Creatinine values are encountered in increased muscle mass, severe dehydration, Pre-eclampsia, increased ingestion of cooked meat, consuming Protein/ Creatine supplements, Diabetic Ketoacidosis, prolonged fasting, renal dysfunction and drugs such as cefoxitin ,cefazolin, ACE inhibitors ,angiotensin II receptor antagonists, N-acetylcyteine , chemotherapeutic agent such as flucytosine

Uric Acid 7.62 3.5 - 7.2mg/dL

(Serum/Enzymatic)



APPROVED BY

-- End of Report --



Name	Mr.KUMAR MAHESH	ID	MED121679603
Age & Gender	37/MALE	Visit Date	21/02/2023
Ref Doctor Name	MediWheel		

ABDOMINO-PELVIC ULTRASONOGRAPHY

LIVER is normal in shape, size (12.5 cm) and shows increased echogenicity. No evidence of focal lesion or intrahepatic biliary ductal dilatation. Hepatic and portal vein radicals are normal.

GALL BLADDER is partially distended.

CBD is not dilated.

PANCREAS has normal shape, size and uniform echopattern. No evidence of ductal dilatation or calcification.

SPLEEN shows normal shape, size and echopattern.

No demonstrable Para-aortic lymphadenopathy.

BOTH KIDNEYS

Right kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

Left kidney: Normal in shape, size and echopattern. Cortico-medullary differentiation is well madeout. No evidence of calculus or hydronephrosis.

The kidney measures as follows:

	Bipolar length (cms)	Parenchymal thickness (cms)
Right Kidney		
Left Kidney	10.8	1.7

URINARY BLADDER is partially distended. It has clear contents. No evidence of diverticula.

PROSTATE shows normal shape, size and echopattern.

No evidence of ascites.

IMPRESSION:

• Grade I fatty infiltration of liver.

REPORT DISCLAIMER

- 1. This is only a radiologincal imperssion. Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.
- 2. The results reported here in are subject to interpretation by qualified medical professionals only.
- 3. Customer identities are accepted provided by the customer or their representative.
- 4.information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.
- 5.If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.
- 6.Test results should be interpreted in context of clinical and other findings if any. In case of any clarification /doubt, the refrering doctor/patient can contact the respective section head of the laboratory.
- 7.Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,
- 8.If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.
- 9.Liability is limited to the extend of amount billed.
- $10. \\ Reports are subject to interpretation in their entirety, partial or selective interpretation may lead to false opinion.$
- 11.Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.



Name	Mr.KUMAR MAHESH	ID	MED121679603
Age & Gender	37/MALE	Visit Date	21/02/2023
Ref Doctor Name	MediWheel		

• No other significant abnormality detected.

DR. HEMANANDINI V.N CONSULTANT RADIOLOGISTS Hn/mj

REPORT DISCLAIMER

^{1.}This is only a radiologincal imperssion.Like other investigations, radiological investication also have limitation. Therefore radiologincal reports should be interpreted in correlation with clinical and pathological findings.

^{2.} The results reported here in are subject to interpretation by qualified medical professionals only.

^{3.} Customer identities are accepted provided by the customer or their representative.

^{4.}information about the customer's condition at the time of sample collection such as fasting, food consumption, medication, etc are accepted as provided by the customer or representative and shall not be investigated for its truthfulness.

^{5.}If any specimen/sample is received from any others laboratory/hospital,its is presumed that the sample belongs to the patient identified or named.

^{6.}Test results should be interpreted in context of clinical and other findings if any.In case of any clarification /doubt , the refrering doctor/patient can contact the respective section head of the laboratory.

^{7.}Results of the test are influenced by the various factors such as sensitivity, specificity of the procedures of the tests, quality of the samples and drug interactions etc.,

^{8.}If the test results are found not to be correlating clinically can contact the lab in charge for clarification or retesting where practicable within 24 hours from the time of issue of results.

^{9.}Liability is limited to the extend of amount billed.

^{10.}Reports are subject to interpretation in their entirety.partial or selective interpretation may lead to false opinion.

^{11.}Disputes, if any , with regard to the report findings are subject to the exclusive jurisdiction of the competent courts chennai only.