



Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:09
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 12:25:42
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF HAEM ATOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
Blood Group (ABO & Rh typing), Blood				
Blood Group	0			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Ph (Anti-D)	POSTIVE			ERYTHROCYTE MAGNETIZED TECHNOLOGY/ TUBE AGGLUTINA
Complete Blood Count (CBC), Whole Blood				
Haemoglobin	14.80	g/ dl	1 Day- 14.5-22.5 g/dl 1 Wk- 13.5-19.5 g/dl 1 Mo- 10.0-18.0 g/dl 3-6 Mo- 9.5-13.5 g/dl 0.5-2 Yr- 10.5-13.5 g/dl 2-6 Yr- 11.5-15.5 g/dl 6-12 Yr- 11.5-15.5 g/dl 12-18 Yr 13.0-16.0 g/dl Male- 13.5-17.5 g/dl Female- 12.0-15.5 g/dl	COLORIMETRICMETHOD (CYANIDE-FREE REAGENT)
TLC (WBC) DLC	4,600.00	/Qumm	4000-10000	IMPEDANCE METHOD
Polymorphs (Neutrophils)	60.00	%	40-80	FLOW CYTOMETRY
Lymphocytes	32.00	%	20-40	FLOW CYTOMETRY
Monocytes	7.00	%	2-10	FLOW CYTOMETRY
Eosinophils	1.00	%	1-6	FLOW CYTOMETRY
Basophils ESR	0.00	%	<1-2	FLOW CYTOMETRY
Observed	6.00	MM/1H	10-19 Yr 8.0 20-29 Yr 10.8 30-39 Yr 10.4 40-49 Yr 13.6 50-59 Yr 14.2 60-69 Yr 16.0 70-79 Yr 16.5 80-91 Yr 15.8	



Page 1 of 12







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:09
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 12:25:42
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF HAEM ATOLOGY


MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
			Pregnancy Early gestation - 48 (62 if anaemic) Leter gestation - 70 (95 if anaemic)	
Corrected	-	Mm for 1st hr.	<9	
PCV (HCT)	48.00	%	40-54	
Platelet count				
Platelet Count	1.10	LACS' cu mm	1.5-4.0	ELECTRONIC IMPEDANCE/MICROSCOPIC
PDW (Platelet Distribution width)	16.40	fL	9-17	ELECTRONIC IMPEDANCE
P-LOR (Platelet Large Cell Patio)	-	%	35-60	ELECTRONIC IMPEDANCE
PCT (Platelet Hematocrit)	0.14	%	0.108-0.282	ELECTRONIC IMPEDANCE
MPV (Mean Platelet Volume)	14.40	fL	6.5-12.0	ELECTRONIC IMPEDANCE
RBCCount				
RBC Count	5.01	Mill./cumm	4.2-5.5	ELECTRONIC IMPEDANCE
Blood Indices (MCV, MCH, MCHC)				
MCV	96.10	fl	80-100	CALCULATED PARAMETER
МОН	29.50	pg	27-32	CALCULATED PARAMETER
МОНС	30.70	%	30-38	CALCULATED PARAMETER
RDW-CV	15.00	%	11-16	ELECTRONIC IMPEDANCE
RDW-SD	52.70	fL	35-60	ELECTRONIC IMPEDANCE
Absolute Neutrophils Count	2,760.00	/cu mm	3000-7000	
Absolute Eosinophils Count (AEC)	46.00	/cumm	40-440	

th-











Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:10
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 11:36:22
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interva	al Method
GLUCOSE FASTING, Plasma Glucose Fasting	148.10	JC	100 Normal)0-125 Pre-diabetes 126 Diabetes	GOD POD

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

CLINICAL SIGNIFICANCE:- Glucose is the major source of energy in the body. Lack of insulin or resistance to it section at the cellular level causes diabetes. Therefore, the blood glucose levels are very high. Elevated serum glucose levels are observed in diabetes mellitus and may be associated with pancreatitis, pituitary or thyroid dysfunction and liver disease. Hypoglycaemia occurs most frequently due to over dosage of insulin.

Gucose PP	311.20	mg/ dl	<140 Normal	GOD POD
Sample:Plasma After Meal			140-199 Pre-diabetes	
			>200 Diabetes	

Interpretation:

a) Kindly correlate clinically with intake of hypoglycemic agents, drug dosage variations and other drug interactions.b) A negative test result only shows that the person does not have diabetes at the time of testing. It does not mean that the person will never get diabetics in future, which is why an Annual Health Check up is essential.c) I.G.T = Impaired Glucose Tolerance.

GLYCOSYLATED HAEM OGLOBIN (HBA1C), EDTA BLOOD

Glycosylated Haemoglobin (HbA1c)	7.60	%NGSP	HPLC (NGSP)
Glycosylated Haemoglobin (HbA1c)	59.90	mmol/mol/IFCC	
Estimated Average Glucose (eAG)	172	mg/ dl	

Interpretation:

<u>NOTE</u>:-

• eAG is directly related to A1c.



View Reports on

Chandan 24x7 App

Page 3 of 12







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:10
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 11:36:22
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name Result	Unit	Bio. Ref. Interval	Method	
------------------	------	--------------------	--------	--

- An A1c of 7% -the goal for most people with diabetes-is the equivalent of an eAG of 154 mg/dl.
- eAG may help facilitate a better understanding of actual daily control helping you and your health care provider to make necessary changes to your diet and physical activity to improve overall diabetes mnagement.

The following ranges may be used for interpretation of results. However, factors such as duration of diabetes, adherence to therapy and the age of the patient should also be considered in assessing the degree of blood glucose control.

Haemoglobin A1C (%)NGSP	mmol/mol / IFCC Unit	eAG (mg/dl)	Degree of Glucose Control Unit
> 8	>63.9	>183	Action Suggested*
7-8	53.0 -63.9	154-183	Fair Control
< 7	<63.9	<154	Goal**
6-7	42.1 -63.9	126-154	Near-normal glycemia
< 6%	<42.1	<126	Non-diabetic level

*High risk of developing long term complications such as Retinopathy, Nephropathy, Neuropathy, Cardiopathy, etc. **Some danger of hypoglycemic reaction in Type 1diabetics. Some glucose intolerant individuals and "subclinical" diabetics may demonstrate HbA1C levels in this area.

N.B.: Test carried out on Automated VARIANT II TURBO HPLC Analyser.

<u>Clinical Implications:</u>

*Values are frequently increased in persons with poorly controlled or newly diagnosed diabetes.

*With optimal control, the HbA 1c moves toward normal levels.

*A diabetic patient who recently comes under good control may still show higher concentrations of glycosylated hemoglobin. This level declines gradually over several months as nearly normal glycosylated *Increases in glycosylated hemoglobin occur in the following non-diabetic conditions: a. Iron-deficiency anemia b. Splenectomy

c. Alcohol toxicity d. Lead toxicity

*Decreases in A 1c occur in the following non-diabetic conditions: a. Hemolytic anemia b. chronic blood loss

*Pregnancy d. chronic renal failure. Interfering Factors:

*Presence of Hb F and H causes falsely elevated values. 2. Presence of Hb S, C, E, D, G, and Lepore (autosomal recessive mutation resulting in a hemoglobinopathy) causes falsely decreased values.

BUN (Blood	Urea	Nitrogen))
Sample:Serum			

10.09

mg/dL 7.0-23.0

CALCULATED



Page 4 of 12







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR - 6	6311	Registered Collected	l On	: 13/Oct/2024 09	
Age/Gender UHID/MR NO	: 51 Y 9 M 16 D /M : ALDP.0000079402		Received		: 13/Oct/2024 09 : 13/Oct/2024 09	
Visit ID	: ALDP0260932425		Reported		: 13/Oct/2024 11	
Ref Doctor	: Dr. MEDIWHEEL-ARCO CARE LTD -	FEMI HEALTH	Status		: Final Report	
		DEPARTMEN	TOFBIOCH	IEMISTR	Y	
	MEDIWH	IEEL BANK OF	BARODA M	ALE AB	OVE 40 YRS	
Test Name		Result	ι	Init	Bio. Ref. Interval	Method
Interpretation:						
Note: Elevated B	UN levels can be seen in th	e following:				
High-protein diet, I	Dehydration, Aging, Certain m	edications, Burns,	Gastrointestir	nal (GI) bl	leeding.	
Low BUN levels o	can be seen in the following	:				
Low-protein diet, o	overhydration, Liver disease.					
		4.05	<i>(</i> 1	0740		
reatinine ample:Serum		1.05	mg/ dl	0.7-1.3	O N	AODIFIED JAFFES
ample:Serum Interpretation: The significance of mass will have a hig absolute creatinine	single creatinine value must b gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma	e interpreted in lig The trend of serun ne concentrations	ht of the patien n creatinine co may increase	nts muscle oncentratio when an A	mass. A patient wit ns over time is mor CE inhibitor (ACE	th a greater muscle e important than E) is taken. The assay
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m	gher creatinine concentration.	e interpreted in lig The trend of serun ne concentrations	ht of the patien n creatinine co may increase	nts muscle oncentratio when an A	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her	th a greater muscle e important than E) is taken. The assay
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fo	e interpreted in lig The trend of serun ne concentrations lous values if serun 5.34 5.34	ht of the patien n creatinine co may increase m samples hav mg/ dl	nts muscle oncentratio when an A /e heteropl 3.4-7.0	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her	th a greater muscle e important than E) is taken. The assay nolyzed, icteric or
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma	e interpreted in lig The trend of serun ne concentrations lous values if serun 5.34 5.34	ht of the patien n creatinine co may increase m samples hav mg/ dl	nts muscle oncentratio when an A /e heteropl 3.4-7.0	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her	th a greater muscle e important than E) is taken. The assay nolyzed, icteric or
Ample:Serum Interpretation: The significance of mass will have a high absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the for protein diet, alcohol), Chronic IA GT) , Serum	e interpreted in lig The trend of serun ne concentrations lous values if serun 5.34 5.34 ollowing: kidney disease, H	ht of the patien n creatinine co may increase m samples hav mg/dl	nts muscle oncentratio when an A /e heteropl 3.4-7.0	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her	th a greater muscle te important than b) is taken. The assay nolyzed, icteric or JRICASE
Interpretation: The significance of mass will have a high absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMM SGOT / Aspartate A	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fo protein diet, alcohol), Chronic IA GT) , Serum Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if serun 5.34 5.34 ollowing: kidney disease, H	ht of the patien n creatinine co may increase m samples hav mg/dl ypertension, C	nts muscle oncentratio when an A ze heteropl 3.4-7.0 Dbesity.	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her U	th a greater muscle e important than b) is taken. The assay nolyzed, icteric or JRICASE
Interpretation: The significance of mass will have a high absolute creatinine could be affected me lipemic. Fric Acid Ample:Serum Interpretation: Note:- Elevated uric acid Drugs, Diet (high-prise) FT (WITH GAMM SGOT / Aspartate A SGPT / Alanine Am	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the for protein diet, alcohol), Chronic IA GT) , Serum	e interpreted in lig The trend of serum ne concentrations lous values if serum 5.34 ollowing: kidney disease, H 14.20 13.70	ht of the patien n creatinine co may increase m samples hav mg/ dl ypertension, C U/L U/L	nts muscle oncentratio when an A ze heteropl 3.4-7.0 Desity. <35 <40	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her U	th a greater muscle e important than E) is taken. The assay nolyzed, icteric or JRICASE
Interpretation: The significance of mass will have a high absolute creatinine could be affected main lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-present FT (WITH GAMIN SGOT / Aspartate A SGPT / Alanine Arr Gamma GT (GGT)	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fo protein diet, alcohol), Chronic IA GT) , Serum Aminotransferase (AST)	e interpreted in lig The trend of serum ne concentrations lous values if serum 5.34 bllowing: kidney disease, H 14.20 13.70 12.00	ht of the patien n creatinine co may increase m samples hav mg/dl ypertension, C U/L U/L IU/L	nts muscle oncentratio when an A /e heteropl 3.4-7.0 Desity. <35 <40 11-50	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her U	th a greater muscle e important than E) is taken. The assay nolyzed, icteric or URICASE
ample:Serum Interpretation: The significance of mass will have a hig absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMIN SGOT / Aspartate A SGPT / Alanine Am Gamma GT (GGT) Protein	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fo protein diet, alcohol), Chronic IA GT) , Serum Aminotransferase (AST)	e interpreted in lig The trend of serun ne concentrations lous values if serun 5.34 ollowing: kidney disease, H 14.20 13.70 12.00 6.67	ht of the patien n creatinine co may increase m samples hav mg/dl ypertension, C U/L U/L IU/L gm/dl	nts muscle oncentratio when an A /e heteropl 3.4-7.0 Desity. <35 <40 11-50 6.2-8.0	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her U	th a greater muscle e important than b) is taken. The assay nolyzed, icteric or JRICASE
Interpretation: The significance of mass will have a hig absolute creatinine could be affected m lipemic. Interpretation: Note:- Elevated uric acid Drugs, Diet (high-p FT (WITH GAMIN SGOT / Aspartate J SGPT / Alanine Arr Gamma GT (GGT)	gher creatinine concentration. concentration. Serum creatinin hildly and may result in anoma d levels can be seen in the fo protein diet, alcohol), Chronic IA GT) , Serum Aminotransferase (AST)	e interpreted in lig The trend of serum ne concentrations lous values if serum 5.34 bllowing: kidney disease, H 14.20 13.70 12.00	ht of the patien n creatinine co may increase m samples hav mg/dl ypertension, C U/L U/L IU/L	nts muscle oncentratio when an A /e heteropl 3.4-7.0 Desity. <35 <40 11-50	mass. A patient wit ns over time is mor CE inhibitor (ACE hilic antibodies, her U	th a greater muscle e important than E) is taken. The assay nolyzed, icteric or URICASE









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patie	nt Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:10
Age/	Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID	/ MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit	ID	: ALDP0260932425	Reported	: 13/Oct/2024 11:36:22
Ref [Ooctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report


DEPARTMENT OF BIOCHEMISTRY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	ι	Jnit	Bio. Ref. Interv	al Method
Alkaline Phosphatase (Total)	103.00	U/L	42.0-1	165.0	PNP/ AMP KINETIC
Bilirubin (Total)	0.95	mg/ dl	0.3-1.	2	JENDRASSIK & GROF
Bilirubin (Direct)	0.38	mg/ dl	< 0.30)	JENDRASSIK & GROF
Bilirubin (Indirect)	0.57	mg/ dl	<0.8		JENDRASSIK & GROF
LIPID PROFILE (MINI), Serum					
Cholesterol (Total)	141.00	mg/ dl		Desirable 39 Borderline High High	CHOD-PAP
HDL Cholesterol (Good Cholesterol)	51.80	mg/ dl	30-70		DIRECT ENZYMATIC
LDL Cholesterol (Bad Cholesterol)	73	mg/ dl	100-12 Optim 130-1 160-12	Optimal 29 Nr. nal/ Above Optima 59 Borderline High 89 High Very High	
VLDL	15.94	mg/ dl	10-33		CALCULATED
Triglycerides	79.70	mg/ dl	150-1 200-4	Normal 99 Borderline High 99 High Very High	GPO-PAP

th-

View Reports on









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:09
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 13:12:32
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 13:28:12
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 15:32:10
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
URINE EXAMINATION, ROUTINE, Urine				
Color	PALEYELOW			
Specific Gravity	1.010			
Reaction PH	Acidic (5.0)			DIPSTICK
Appearance	CLEAR			
Protein	ABSENT	mg %	<10 Absent 10-40 (+) 40-200 (++) 200-500 (+++) >500 (++++)	DIPSTICK
Sugar	ABSENT	gms%	<0.5 (+) 0.5-1.0 (++) 1-2 (+++) >2 (++++)	DIPSTICK
Ketone	ABSENT	mg/dl	Serum-0.1-3.0 Urine-0.0-14.0	BIOCHEMISTRY
Bile Salts	ABSENT			
Bile Pigments	ABSENT			
Bilirubin	ABSENT			DIPSTICK
Leucocyte Esterase	ABSENT			DIPSTICK
Urobilinogen(1:20 dilution)	ABSENT			
Nitrite	ABSENT			DIPSTICK
Blood	ABSENT			DIPSTICK
Microscopic Examination:				
Epithelial œlls	1-2/h.p.f			MICROSCOPIC EXAMINATION
Puscells	0-2/h.p.f			
RBCs	ABSENT			MICROSCOPIC EXAMINATION
Cast	ABSENT			
Crystals	ABSENT			MICROSCOPIC EXAMINATION
Others	ABSENT			

Urine Microscopy is done on centrifuged urine sediment.



Page 7 of 12







Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 CIN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:09
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 13:12:32
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 13:28:12
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 15:32:10
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF CLINICAL PATHOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
SUGAR, FASTING STAGE, Urine Sugar, Fasting stage	ABSENT	gms%		
Interpretation: (+) < 0.5				
SUGAR, PP STAGE, Urine Sugar, PP Stage	TRACE			
Interpretation: (+) < 0.5 gms%				

th-









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965.0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:09
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 11:58:46
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method
PSA (Prostate Specific Antigen), Total Sample:Serum	0.49	ng/mL	≪4.1	ALD

Interpretation:

- 1. PSA is detected in the serum of males with normal, benign hypertrophic, and malignant prostate tissue.
- 2. Measurement of serum PSA levels is not recommended as a screening procedure for the diagnosis of cancer because elevated PSA levels also are observed in patients with benign prostatic hypertrophy. However, studies suggest that the measurement of PSA in conjunction with digital rectal examination (DRE) and ultrasound provide a better method of detecting prostate cancer than DRE alone[.]
- 3. PSA levels increase in men with cancer of the prostate, and after radical prostatectomy PSA levels routinely fall to the undetectable range.
- 4. If prostatic tissue remains after surgery or metastasis has occurred, PSA appears to be useful in detecting residual and early recurrence of tumor.
- 5. Therefore, serial PSA levels can help determine the success of prostatectomy, and the need for further treatment, such as radiation, endocrine or chemotherapy, and in the monitoring of the effectiveness of therapy.

THYROID PROFILE - TOTAL, Serum

T3, Total (tri-iodothyronine)	123.00	ng/ dl	84.61–201.7	ALD
T4, Total (Thyroxine)	6.80	ug/ dl	3.2-12.6	ALD
TSH (Thyroid Stimulating Hormone)	7.800	μIU/mL	0.27 - 5.5	ALD

Interpretation:

0.3-4.5	µIU/mL	First Trimest	er	
0.5-4.6	µIU/mL	Second Trimester		
0.8-5.2	µIU/mL	Third Trimester		
0.5-8.9	µIU/mL	Adults	55-87 Years	
0.7-27	µIU/mL	Premature	28-36 Week	
2.3-13.2	µIU/mL	Cord Blood	> 37Week	
0.7-64	µIU/mL	Child(21 wk	- 20 Yrs.)	
1-39	µIU/mL	Child	0-4 Days	
1.7-9.1	µIU/mL	Child	2-20 Week	

08069366666

1) Patients having low T3 and T4 levels but high TSH levels suffer from primary hypothyroidism, cretinism, juvenile myxedema or





Page 9 of 12



33)

CHANDAN DIAGNOSTIC CENTRE

Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:09
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 13/Oct/2024 09:32:09
UHID/MR NO	: ALDP.0000079402	Received	: 13/Oct/2024 09:56:32
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 11:58:46
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF IMMUNOLOGY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

Test Name	Result	Unit	Bio. Ref. Interval	Method

autoimmune disorders.

2) Patients having high T3 and T4 levels but low TSH levels suffer from Grave's disease, toxic adenoma or sub-acute thyroiditis.

3) Patients having either low or normal T3 and T4 levels but low TSH values suffer from iodine deficiency or secondary hypothyroidism.

4) Patients having high T3 and T4 levels but normal TSH levels may suffer from toxic multinodular goiter. This condition is mostly a symptomatic and may cause transient hyperthyroidism but no persistent symptoms.

5) Patients with high or normal T3 and T4 levels and low or normal TSH levels suffer either from T3 toxicosis or T4 toxicosis respectively.

6) In patients with non thyroidal illness abnormal test results are not necessarily indicative of thyroidism but may be due to adaptation to the catabolic state and may revert to normal when the patient recovers.

7) There are many drugs for eg. Glucocorticoids, Dopamine, Lithium, Iodides, Oral radiographic dyes, etc. which may affect the thyroid function tests.

8) Generally when total T3 and total T4 results are indecisive then Free T3 and Free T4 tests are recommended for further confirmation along with TSH levels.

View Reports on

Chandan 24x7 App









Add: 49/19-B, Kamla Nehru Road, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR -66311	Registered On	: 13/Oct/2024 09:25:10
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 2024-10-13 09:49:22
UHID/MR NO	: ALDP.0000079402	Received	: 2024-10-13 09:49:22
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 13:51:49
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF X-RAY

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

X-RAY DIGITAL CHEST PA

<u>X-RAY REPORT</u> (300 mA COMPUTERISED UNIT SPOT FILM DEVICE) <u>CHEST P-A VIEW</u>

- Both lung field did not reveal any significant lesion.
- Costo-phrenic angles are bilaterally clear.
- Trachea is central in position.
- Cardiac size & contours are normal.
- Hilar shadows are normal.
- Soft tissue shadow appears normal.
- Bony cage is normal.

Please correlare clinically.





Home Sample Collection 08069366666 Page 11 of 12

View Reports on

Chandan 24x7 App







Add: 49/19-B, Kamla Nehru Poad, Katra, Prayagraj Ph: 9235447965,0532-3559261 QN: U85110UP2003PLC193493

Patient Name	: Mr.VIJENDRA KUAMR - 66311	Registered On	: 13/Oct/2024 09:25:10
Age/Gender	: 51 Y 9 M 16 D / M	Collected	: 2024-10-13 10:37:38
UHID/MR NO	: ALDP.0000079402	Received	: 2024-10-13 10:37:38
Visit ID	: ALDP0260932425	Reported	: 13/Oct/2024 10:52:20
Ref Doctor	: Dr. MEDIWHEEL-ARCOFEMI HEALTH CARE LTD -	Status	: Final Report

DEPARTMENT OF ULTRASOUND

MEDIWHEEL BANK OF BARODA MALE ABOVE 40 YRS

ULTRASOUND WHOLE ABDOM EN (UPPER & LOWER)

LIVER: - Normal in size (15.4 cm), shape and **shows diffusely raised echotexture**. No focal lesion is seen. No intra hepatic biliary radicle dilation is seen.

GALL BLADDER :- Well distended. Normal wall thickness is seen. No evidence of calculus/focal mass lesion/pericholecystic fluid is seen.

CBD :- Normal in calibre at porta.

PORTAL VEIN: - Normal in calibre and colour uptake at porta.

PANCREAS: - Head is visualised, normal in size & echopattern. No evidence of ductal dilatation or calcification is seen. Rest of the pancreas is obscured by bowel gases.

SPLEEN: - Normal in size (7.7 cm), shape and echogenicity. No evidence of mass lesion is seen.

RIGHT KIDNEY: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

LEFT KIDNEY: - Normal in size, shape and position. Cortical echogenicity is normal with maintained corticomedullary differentiation. No focal lesion or calculus is seen. Pelvicalyceal system is not dilated.

URINARY BLADDER :- Is adequately distended. No evidence of calculus is seen. **Wall is thickened** (Maximum thickness 7.2 mm) and irregular. Pre void vol - 236 cc, Post void vol - 45 cc.

PROSTATE :- Normal in size (2.3 x 2.0 x 3.1 cm vol - 7.5 cc), shape and echo pattern.

HIGH RESOLUTION :- No evidence of bowel loop dilatation or abnormal wall thickening is seen. No significant retroperitoneal lymphadenopathy is seen. No free fluid is seen in the abdomen/pelvis.

IMPRESSION :

- Grade I fatty liver.
- Chronic cystitis.

Please correlate clinically

