FINAL REPORT

Bill No.	:	APHHC240000342	Bill Date	Г	02-03-2024 10:04		
Patient Name	:	MR. NUG NARAYAN PRASAD	UHID	Г	APH000021041		
Age / Gender	:	41 Yrs 2 Mth / MALE	Patient Type	Г	OPD	If PHC	:
Ref. Consultant	:	MEDIWHEEL	Ward / Bed	Г	1		
Sample ID	:	APH24007542	Current Ward / Bed	F	1		
	:		Receiving Date & Time	F	02-03-2024 15:56		
	Г		Reporting Date & Time	Γ	02-03-2024 16:43		

BIOCHEMISTRY REPORTING

Interval	Test (Methodology)	Flag	Result	UOM	Biological Reference Interval
----------	--------------------	------	--------	-----	----------------------------------

Sample Type: EDTA Whole Blood, Plasma, Serum

MEDIWHEEL FULL BODY HEALTH CHECKUP _MALE(ABOVE 40)@2550

BLOOD UREA Urease-GLDH,Kinetic		20	mg/dL	15 - 45
BUN (CALCULATED)		9.3	mg/dL	7 - 21
CREATININE-SERUM (Modified Jaffe's Kinetic)	L	0.8	mg/dL	0.9 - 1.3
GLUCOSE-PLASMA (FASTING) (UV Hexokinase)		100.0	mg/dL	70 - 100

Note: A diagnosis of diabetes mellitus is made if fasting blood glucose exceeds 126 mg/dL. (As per American Diabetes Association recommendation)

7

GLUCOSE-PLASMA (POST PRANDIAL) (UV Hexokinase)	118.0	mg/dL	70 - 140

Note: A diagnosis of diabetes mellitus is made if 2 hour post load glucose exceeds 200 mg/dL. (As per American Diabetes Association recommendation)

LIPID PROFILE

CHOLESTROL-TOTAL (CHO-POD)	Н	178	mg/dL	0 - 160
HDL CHOLESTROL Enzymatic Immunoinhibition		48	mg/dL	>40
CHOLESTROL-LDL DIRECT Enzymatic Selective Protection	Н	107	mg/dL	0 - 100
S.TRIGLYCERIDES (GPO - POD)		123	mg/dL	0 - 160
NON-HDL CHOLESTROL	Н	130.0	mg/dL	0 - 125
TOTAL CHOLESTROL / HDL CHOLESTROL		3.7		1/2Average Risk < 3.3 Average Risk 3.3-4.4 2 Times Average Risk 4.5-7.1 3 Times Average Risk 7.2-11.0
LDL CHOLESTROL / HDL CHOLESTROL		2.2		1/2Average Risk <1.0 Average Risk 1.0-3.6 2 Times Average Risk 3.7-6.3 3 Times Average Risk 6.4-8.0
CHOLESTROL-VLDL		25	mg/dL	10 - 35

Comments:

- Disorders of Lipid metabolism play a major role in atherosclerosis and coronary heart disease.
- There is an established relationship between increased total cholesterol & LDL cholesterol and myocardial infarction.
- HDL cholesterol level is inversely related to the incidence of coronary artery disease.
- Major risk factors which adversely affect the lipid levels are:
 - 1. Cigarette smoking.
 - 2. Hypertension.
 - 3. Family history of premature coronary heart disease.
 - 4. Pre-existing coronary heart disease.

LIVER FUNCTION TESTS (LFT)

BILIRUBIN-TOTAL (DPD)		0.63	mg/dL	0.2 - 1.0
BILIRUBIN-DIRECT (DPD)		0.11	mg/dL	0 - 0.2
BILIRUBIN-INDIRECT		0.52	mg/dL	0.2 - 0.8
S.PROTEIN-TOTAL (Biuret)		6.6	g/dL	6 - 8.1

FINAL REPORT

ill No.	T	APHHC240000342			Bill Date			02-03-2024 10:04		
atient Name	1	: MR. NUG NARAYAN PRASAD			UHID			APH000021041		
ge / Gender	1	41 Yrs 2 Mth / MALE	Mth / MALE		Patient Type		:	OPD	If PHC :	
ef. Consultant	1	MEDIWHEEL	EL				:	1		
ample ID	1	APH24007542			Current Ward / Bed		:	1		
	1				Receiving Date & Time		:	02-03-2024 15:56	<u>2</u> 4 15 56	
	Т				Reporting Date & Tin	1e	:	02-03-2024 16:43		
ALBUMIN-SER	UM	1 (Dye Binding-Bromocresol Green)		4.3	3	g/dL				
S.GLOBULIN			L	2.	3	g/dL		2.8-3.8		
A/G RATIO				1.8	37			1.5 - 2	.5	
ALKALINE PHO	OSF	PHATASE IFCC AMP BUFFER		54	.6	IU/L		53 - 12	8	
ASPARTATE A	ΜI	NO TRANSFERASE (SGOT) (IFCC)		40	.3	IU/L		10 - 42		
ALANINE AMI	NO	TRANSFERASE(SGPT) (IFCC)	Н	83	3.7	IU/L		10 - 40		
GAMMA-GLUT	ΑΜ	YLTRANSPEPTIDASE (IFCC)		44	.7	IU/L		11 - 50		
LACTATE DEH	ΙYD	ROGENASE (IFCC; L-P)		17	9.1	IU/L		0 - 248	3	
S.PROTEIN-TO	OT <i>A</i>	AL (Biuret)		6.6	3	g/dL		6 - 8.1		
			1	<u> </u>						
URIC ACID Urio	ase -	Trinder		4.8	3	mg/d	L	2.6 - 7	.2	

** End of Report **

IMPORTANT INSTRUCTIONS
CL - Critical Low, CH - Critical High, H - High, L - Low

DR. ASHISH RANJAN SINGH

MBBS,MD CONSULTANT

FINAL REPORT

Bill No.	T	APHHC240000342	Bill Date	T	02-03-2024 10:04		
Patient Name	Г	MR. NUG NARAYAN PRASAD	UHID	Г	APH000021041		
Age / Gender	Г	41 Yrs 2 Mth / MALE	Patient Type	Г	OPD If PHC :		
Ref. Consultant	Г	MEDIWHEEL	Ward / Bed		1		
Sample ID	1	APH24007542	Current Ward / Bed	1	1		
	F		Receiving Date & Time	:	02-03-2024 15:56		
	Т		Reporting Date & Time		02-03-2024 16:43		

Sample Type: EDTA Whole Blood, Plasma, Serum

MEDIWHEEL FULL BODY HEALTH CHECKUP _MALE(ABOVE 40)@2550

HBA1C (Turbidimetric Immuno-inhibition)	Н	6.3	%	4.0 - 6.2

INTERPRETATION:

HbA1c %	Degree of Glucose Control
>8% Action suggested due to high risk of developing long term complications like Retinop Nephropathy, Cardiopathy and Neuropathy	
7.1 - 8.0	Fair Control
<7.0	Good Control

Note:

- 1.A three monthly monitoring is recommended in diabetics.
- 2. Since HbA1c concentration represents the integrated values for blood glucose over the preceding 6 10 weeks and is not affected by daily glucose fluctuation, exercise and recent food intake, it is a more useful tool for monitoring diabetics.

** End of Report **

IMPORTANT INSTRUCTIONS

CL - Critical Low, CH - Critical High, H - High, L - Low

DR. ASHISH RANJAN SINGH

MBBS,MD CONSULTANT